

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`


600047162Q

Math. B.

A TREATISE

ON

SPHERICAL TRIGONOMETRY,

WITH NUMEROUS EXAMPLES.

Part I.

TO THE END OF THE SOLUTION OF TRIANGLES.

BY

WILLIAM J. M'CLELLAND, Sch., B.A.,

Principal of the Incorporated Society's School, Santry, Dublin;

AND

THOMAS PRESTON, Sch., B.A.

DUBLIN: HODGES, FIGGIS, & CO., GRAFTON STREET.

LONDON: LONGMANS, GREEN, & CO., PATERNOSTER ROW.


1885.

18382 e. 1

DUBLIN :

PRINTED AT THE UNIVERSITY PRESS,

BY PONSONBY AND WELDRICK.



PREFACE.

THE object of the present Treatise is to bring Spherical Trigonometry to the standard required for University Examinations, and demanded by the impulse given to mathematical subjects by modern text-books.

Simplicity of treatment has been constantly kept in view.

Part I. treats of the subject as far as the solution of Triangles, inclusive. In the text will be found all the propositions usually contained in treatises on the subject, besides such other theorems as appeared to us to be of special importance on account of their utility.

The volume is replete with examples (in many cases worked out), the arrangement of which has been the subject of our special attention; our aim throughout being to place them in immediate connexion with the subject matter of which they are illustrative.

At the end of each Chapter, Miscellaneous Examples bearing on all the preceding matter have been added. We have not hesitated to use Determinant Notation

whenever elegance or simplicity could be gained thereby. This notation has now become so generally known as to render apology for its use early in the work quite unnecessary.

The Numerical Solution of Triangles, treated of in Chapters IV. and V., has received much attention, each Case being treated of in detail. In connexion with these Numerical Examples, we must acknowledge our obligations to Mr. ROBERT BAILE, M.A., Athlone, who carefully worked and verified them all. They have also been independently verified and tested, and we therefore trust they will all be found correct to the nearest half second.

Geometrical proofs of many Propositions have been added to those commonly given in text-books, *e.g.* the Analogies of Napier and Delambre.

Most of the examples have been taken from University and Science and Art Examination Papers. Many, however, as far as we are aware, appear now for the first time. A series of Examples have been appended in the form of Examination Papers.

Among published works on the subject, we are indebted chiefly to those of Todhunter, Snowball, and Luby.

WM. J. M'CLELLAND,
THOMAS PRESTON.

May, 1885.

CONTENTS.

PART I.

CHAPTER I.

THE SPHERE.

PAGE

Definitions. Great and Small Circles, ... 1
Axes and Poles, ... 3
Primary and Secondary Circles, ... 4
The Angle between two Great Circles, ... 5
The Lune. Theorem, ... 7
Solid Angles, ... 8
The arc of connexion of two points is the polar of the intersection of the polars of the points, ... 9
Examples, ... 11

CHAPTER II.

SPHERICAL TRIANGLES.

Definitions, ... 12
Colunar or Associated Triangles, ... 13
Parts of a Spherical Triangle, ... 14
Sphere of Infinite Radius, ... 15
Analogy of Problems *in plano* and on the Sphere, ... 17
Polar Triangles. Reciprocal and Supplemental Relations, . . . 22
Limits to sides and angles of a Spherical Triangle, ... 27
Examples, ... 28

CHAPTER III.

FUNDAMENTAL RELATIONS BETWEEN THE PARTS OF A SPHERICAL TRIANGLE.

SECTION I. :—

PAGE

Given the sides to determine the angles, ... 29

Examples, ... 32

Analogous formula *in plano*, ... 34

Legendre's Theorem, ... 34

SECTION II. :—

Expression for the sine of an angle in terms of the sides, . . . 36

Rule of Sines. *Examples*, ... 37

Geometrical proof of the Rule of Sines, ... 39

Expressions for $\sin \frac{1}{2}A$, $\cos \frac{1}{2}A$, and $\tan \frac{1}{2}A$ in terms of the sides, ... 40

Examples, ... 43

Supplemental Theorems, ... 44

Analogous formula *in plano*, ... 45

Examples, ... 45

Expressions for $\sin \frac{1}{2}a$, $\cos \frac{1}{2}a$, and $\tan \frac{1}{2}a$ in terms of the angles, 49

Examples, ... 50

SECTION III.—THEOREMS :—

1. $\cot a \sin b = \cot A \sin C + \cos b \cos C$, ... 52

Corresponding Theorem *in plano*, ... 54

Examples, ... 55

2. $\sin BC \cos AP + \sin CA \cos BP + \sin AB \cos CP = 0$, . . 58

Analogous Theorem *in plano*, ... 59

Examples, ... 60

3. $\tan \frac{1}{2}\rho \tan \frac{1}{2}\rho' = \text{const.}$, ... 61

Examples, ... 63

Table of Formulae, ... 64

Miscellaneous Examples, ... 66

CHAPTER IV.

RIGHT-ANGLED TRIANGLES.

SECTION I. :—

PAGE

Geometrical Deduction of Formulae, ... 78
Napier's Rules of Circular Parts, ... 80
Examples, ... 82

SECTION II. :—

Solution of Right-angled Triangles, ... 83
CASE I.—Having given two sides. *Examples*, . . . 84
CASE II.—Having given two angles. *Examples*, . . . 85
CASE III.—Having given the hypotenuse and a side. *Examples*, 86
CASE IV.—Having given the hypotenuse and an angle. *Examples*, 87
CASE V.—Having given a side and the adjacent angle. *Examples*, 88
CASE VI.—Having given a side and the opposite angle. *Examples*, 89

SECTION III.—THEOREMS :—

The perpendiculars of a triangle are concurrent, . . . 91
(1). $\Sigma \sin BC \sin AA' = 0$, ... 92
(2). $\Sigma \sin B'C' \tan AA' = 0$, ... 92
(3). $\Sigma \sin BC \cot A = 0$, ... 92
(4). $\Sigma \sin B'C' \cos A = 0$, ... 92
Corollaries, ... 94
Geometrical Deduction of the Analogies of Napier and Delambre, 95
Examples, ... 98
Miscellaneous Examples, ... 101

CHAPTER V.

OBLIQUE-ANGLED TRIANGLES.

CASE I.—Having given the three sides. *Examples*, . . 106
CASE II.—Having given the three angles. *Examples*, . . 108
CASE III.—Having given two sides and the included angle, . 110
Napier's Analogies, ... 111
Delambre's (or Gauss's) Analogies, ... 112
Examples of CASE III., ... 114

PAGE

CASE IV.—Having given two angles and the adjacent side. *Examples*, ... 117

CASE V.—Having given two sides and an angle opposite one of them. *Examples*, ... 120

CASE VI.—Having given two angles and a side opposite one of them. *Examples*, ... 125

Discussion of the Ambiguous Case, ... 129

Examples, ... 134

The Subsidiary Angle (Geometrical interpretation), . . . 135

Examples, ... 137

EXAMINATION PAPERS, ... 140

ANSWERS TO EXAMINATION PAPERS, ... 153

CORRIGENDA.

Page 11, lines 12 and 13, for $r \cos \theta$, read $r \sin \theta$.

,, 27, line 5 from bottom, *for* fig. 11, *read* fig. 12.

,, 33, Ex. 7, for $a = b$, read $b = c$.

,, 52, Ex. 14, *for* $\left(\frac{r}{2}\right)^2$, *read* $\left(\frac{\gamma}{2}\right)^2$.

,, 61, Ex. 5, *for* be drawn, *read* are drawn.

,, 63, Ex. 1, *for* Euclid I., *read* Euclid III.

,, 68, Ex. 7, *for* fig. 18, *read* fig. 19.

SPHERICAL TRIGONOMETRY.

CHAPTER I.

The Sphere.



1. **A Sphere** is a solid body such that all points on its surface are equally distant from a certain point within it, called the *Centre*.

Any straight line drawn from the centre of a sphere to the surface is called a *Radius*, and any straight line drawn through the centre and terminated both ways by the surface is called a *Diameter*.

A sphere may be generated by the revolution of a semicircle round its diameter.

2. **Great and Small Circles.**—Let a sphere be gene




Fig. 1.

rated by the revolution of the semicircle ACB (fig. 1)

round its diameter AB. Let C be any point on the semi-circle, and let CD be a perpendicular from C on AB.

It is obvious that as ACB revolves round AB, the point C describes a circle round D as centre; and also that O, the middle point of AB, being equally distant from all points on ACB, is the centre of the sphere.

When the plane of the circle CC' described by C passes through O it divides the sphere into two equal parts, and the curve of section is called a *Great Circle.*

When its plane does not pass through the centre it is called a *Small Circle.*

Example.

The Meridians, Equator, and Ecliptic, are great circles. The Parallels of Latitude are small circles.

The angular distance AC is called the *Spherical Radius* of the circle CC'. It is obvious that the spherical radius of a great circle is a quadrant.

3. **Only one Great Circle** can be drawn through two given points on the surface of a sphere; for its plane must also pass through the centre; and three points not in the same right line are sufficient to determine a plane completely.

If the two given points be diametrically opposite, the right line joining them passes through the centre of the sphere, and an infinite number of great circles can be drawn through them, as, for example, the meridians on the surface of the Earth.

The *shortest distance* that can be traced on the surface of a sphere between two points on it is the arc of the great circle passing through them.

For, when the radius of a sphere is indefinitely increased, its surface, at any part, may be considered as a plane (see Chap. II., Art. 18), and a great circle passing through two points on it becomes a right line joining two points in a plane; but the right line joining two points is the shortest distance between them; therefore, etc.*

It is for this reason, and because only one great circle can be drawn through two points that distances on the surface of a sphere are measured along arcs of great circles. Advantage is taken of this property of the great circle in Navigation.

4. **Axes and Poles.**—The line AB (fig. 1) is called the *Axis* of the circle CC' (described by *any* point C of the semicircle ACB during its revolution round AB), and the extremities A and B of its axis are called its *Poles*. Any point, and the great circle of which it is the pole, are termed *pole* and *polar*, with respect to one another. It is obvious, from the manner in which CC' was generated, that AB is perpendicular to the plane of CC', for CD remained perpendicular to AB during its entire revolution.

Hence the *Axis* of a circle may be defined as *the diameter*

* The following method of looking at this question is also instructive:—

If a string be stretched between two points on the surface of a sphere (or on any surface) it will evidently be the shortest distance that can be traced on the surface between the points, since by pulling the ends of the string its length, between the points, will be shortened as much as the surface will permit. Now any part of the string being acted on by two terminal tensions, and by the reaction of the surface, which is everywhere normal to it, must lie in a plane containing the normal to the surface. Hence, the plane of the string contains the normals to the surface at all points of its length; *i. e.* the string lies in the form of a great circle.

of the sphere perpendicular to the plane of the circle, or *the line joining the centre of the sphere with the pole of the circle.*

Cor.—The pole of a circle is equidistant from all points on the circumference of the circle.

For $AC^2 = AD^2 + CD^2 = \text{constant}$.

In the case of a great circle, D becomes the centre of the sphere, and hence the poles of a great circle are equidistant from its circumference.

5. **Primary and Secondary Circles.**—Any circle is called a *Primary* in relation to those *great circles* which cut it at right angles. These latter are called *Secondaries;* e.g. parallels of latitude are primaries, and the meridians are secondaries to them.

In fig. 1 regarding the circle CC' as a primary, the circle ACB during its revolution round AB is in all positions a secondary to CC'. Hence it follows that:—

(1). The plane of any secondary contains the axis of the primary.

(2). All the secondaries pass through the poles of the primary.

(3). The planes of all the secondaries have a common line of intersection, viz., the axis of the primary.

(4). If there can be drawn common secondaries to two circles, the planes of those circles are parallel. For, by (3), the two circles have the same axis.

The distance of any point on the surface of a sphere from a circle traced thereon is measured by the arc of the secondary intercepted between the point and the circle.

Example.

The latitude of a place is measured by the arc of the meridian intercepted between the place and the Equator.

6. **The Angle between two Planes** is measured by the angle between any two lines drawn, one in each plane, perpendicular to their line of intersection.

The angle between any two circles, great or small, is measured by the angle between the tangents drawn to them at their point of intersection.

The Angle between two Great Circles is equal to—

(1). *The angle between their planes.*

For the tangents to them at their point of intersection lie in their respective planes and are perpendicular to their common diameter.

(2). *The arc intercepted by them on the great circle to which they are secondaries.*

For if AC and BC (fig. 2) be secondaries to AB they




Fig. 2.

are perpendicular to AB, and are each quadrants; therefore OA and OB are perpendiculars to OC; hence the

angle between OA and OB, or the arc AB, is the angle between the planes of AC and BC.

(3). *The angular distance between their poles.*

For the angular distance between the poles of two circles is the same as the angle between their axes; but the axes are perpendicular to the planes of the circles, and the angle between the perpendiculars to two planes is equal to the angle between the planes.

Hence, by drawing a great circle parallel to a small circle, we see that the angle between the planes of a great and a small circle, or the angle between the planes of two small circles, is equal to the angular distance between the poles of the circles.

7. *To compare the length of the arc of a small circle subtending any angle at its centre, with the arc of a great circle subtending an equal angle at the centre of the sphere.*

Let ab (fig. 2) be an arc of a small circle subtending any angle at its centre D, and let AB be an arc of a great circle subtending an equal angle at its centre O, then if the circles be placed parallel, so that they may have a common pole at C, O will be the centre of the sphere, AC and BC will be quadrants, CO will be perpendicular to the planes ABO and abD; and since the perimeters of circles are to each other as their radii, we have

$$\frac{ab}{AB} = \frac{bD}{BO} = \frac{bD}{bO} = \sin bOD = \sin bC = \cos bB.$$

The above equation expresses the arc of a parallel of latitude in terms of the latitude of the place and the corresponding arc of the Equator. Thus $ab = AB \cos$ (latitude); or, the distance ab between two places in the same latitude,

measured on the parallel, is equal to the difference of longitude (AB) multiplied by the cosine of the latitude.

8. **The Lune.**—A *Lune* is a portion of the surface of a sphere enclosed by two great circles.

In fig. 1 the area $ACBEA$ is a lune, so also is the area $AEBC'A$.

The angle between the two great circles bounding the lune is called the *Angle of the Lune.*

The area of a lune is readily expressed in terms of the angle of the lune and radius of the sphere. For we have evidently—

Area of Lune : Area of Sphere : : Angle of Lune : 2π ;

or, denoting the angle of the lune by A, and assuming the area of the sphere to be $4\pi r^2$, where r is the radius of the sphere,

$$\text{Area of Lune} = 2Ar^2.$$

9. **Theorem.**—*Any section of the surface of a sphere by a plane is a circle.*

For (fig. 1) if OD be let fall from O, the centre of the sphere, perpendicular to the plane CC', we have

$$CD^2 = CO^2 - OD^2 = \text{constant},$$

since CO is constant, and OD is fixed. Hence, every point of the section of the surface is equidistant from D, and therefore lies on a circle round D as centre.

10. *The intersection of two spherical surfaces is a circle.*

Consider the triangle ABX formed by joining the centres A and B of the two spheres and any point X on their curve of intersection. The sides of this triangle are *given*, and the line of centres AB is fixed ; hence the opposite vertex X (*i. e.* the variable point on the curve of sec-

tion of the two spheres) is at a constant distance from AB, or lies on a circle having AB for axis.

This property is more general than that stated in the previous Article, and reduces to it as a particular case by supposing the radius of either sphere to become infinitely great.

11. **Solid Angles.**—When three or more planes intersect at a point they form a *solid angle* at that point. There is a solid angle, for example, at each corner of a cube, and at each vertex of a tetrahedron or pyramid. Every surface in the neighbourhood of a point subtends a solid angle at that point, the solid angle being enclosed by the lines drawn from the point to the contour of the surface.




Fig. 3.

Let $ABCD$ (fig. 3) be any surface or plane area. From any point O draw lines OA, OB, OC, OD, &c., touching the surface (or passing through the boundary of the plane area). These lines form a kind of irregular cone, having its vertex at O, and enclosing the surface $ABCD$. The solid angle of this cone is the solid angle which $ABCD$ subtends at O. To measure this angle, an expedient similar to that employed in reckoning plane angles by circular

measure is used. About O describe a sphere of unit radius, and let the cone formed by the lines from O to $ABCD$ intersect the surface of this sphere in the curve $A'B'C'D'$. The curve $A'B'C'D'$ will enclose an area on the surface of the sphere, and this area measures the solid angle at O, just as the arc of a circle of unit radius measures the angle it subtends at the centre.

Since the radius of the sphere is unity, its area is 4π; hence the solid angle subtended at O by the n^{th} part of the surface of the sphere is $\frac{4\pi}{n}$.

Example.

The solid angle at a corner of a cube is $\frac{\pi}{2}$.

12. **Theorem.**—*The point of intersection of two great circles and the arc joining their poles are pole and polar with respect to one another.*

For let A and B be the two points, LO and MO their




Fig. 4.

polars, O the point of intersection of the polars; then $OA = \frac{\pi}{2}$, $OB = \frac{\pi}{2}$; hence OA and OB are secondaries to AB; $\therefore$ &c. (Art. 5 (2)).

Examples.

1. Two great circles bisect each other.

They have a common diameter.

2. The axis of any circle is determined by the intersection of two of its secondaries (Art. 5, sec. (3)).

3. Find the locus of the centres of a system of circles having common secondaries. *Ans.* The common axis of the system.

4. The pole of any great circle is ninety degrees from the circumference (Art. 4).

5. The arc drawn through the poles of two great circles cuts both at right angles.

5*a*. Draw a great circle cutting two given circles, great or small, at right angles.

6. If three or more great circles are concurrent their poles are concyclic.

7. Any great circle is the locus of the poles of all its secondaries.

8. If from any point on the surface of a sphere *two* great circles can be drawn perpendicular to a given great circle, that point is the pole of the circle.

[For, each of the arcs drawn perpendicular are secondaries; ∴ &c. (Art. 5, sec. (2)).]

9. The external bisector of an angle passes through the pole of the internal bisector; and conversely.

10. If two great circles are equally inclined to a third, their poles are equidistant from the pole of the third.

11. Two equal small circles are drawn touching each other: show that the angle between their planes is twice the complement of their spherical radius.

—(*Science and Art Examination Papers.*)

For, join the centres of the small circles to the centre of the sphere, and also to the point of contact of the circles.

Then, the lines drawn to the centre of the sphere contain an angle equal to twice the spherical radius of either small circle, and the lines drawn to the point of contact of the small circles contain an angle equal to the inclination of the planes of the circles; therefore, &c. (*Euc.* III. 22.)

12. On a sphere whose radius is r a small circle of spherical radius, θ, is described, and a great circle is described having its pole on the small circle; show that the length of their common chord is

$$\frac{2r}{\sin\theta}\sqrt{-\cos 2\theta}.$$

(*Science and Art Examination Papers.*)

For, the angle between the axis of the small circle and the plane of the great circle is evidently $90-\theta$. Hence the distance, d, from the middle point of the common chord to the centre of the small circle is given by the equation

$$d = r\cos\theta\cot\theta.$$

Again, if c be the common chord, it follows that

$$\left(\frac{c}{2}\right)^2 = (r\sin\theta + d)(r\sin\theta - d),$$

since $r\sin\theta$ is the radius of the small circle; therefore

$$c = \frac{2r}{\sin\theta}\sqrt{-\cos 2\theta},$$

the negative sign occurring under the radical, since 2θ must be greater than $90°$ for *real* section.

Note—Hence the segments of a great circle, made by the intersection of a small circle of given radius, may be calculated on a sphere of given radius.

13. The angle subtended at the centre of a circle, great or small, by two points on it, is equal to the angle subtended by them at its pole. (Art. 6.)

14. If through the centre O of a sphere a line be drawn parallel to a chord AB of a semicircle ABC described thereon, it meets the sphere at the middle point of the arc BC.

15. Given A, B, C, D, four points on a segment of a circle; prove the relations:—

$\alpha°$. $\sin BC\sin AD + \sin CA\sin BD + \sin AB\sin CD = 0.$

$\beta°$. $\sin BC\cos AD + \sin CA\cos BD + \sin AB\cos CD = 0.$

CHAPTER II.

Spherical Triangles.

13. In the present chapter it will be shown that geometrical figures drawn on the surface of a sphere (or different spheres of equal radii) have many properties and relationships analogous to those of corresponding figures described on a plane surface; and with the latter the reader is supposed to be already familiar.

14. **Definitions.**—By a *Spherical Figure* is meant a portion of the surface of a sphere enclosed by arcs of *great** circles, *e. g.*

A *Spherical Triangle* is bounded by the arcs of three great circles; a *Spherical Quadrilateral* by four great circles; a *Spherical Polygon* by many great circles.

In the particular case, when the number of enclosing arcs becomes indefinitely great, the figure becomes a *Spherical Curve.* A great circle drawn through two indefinitely near points on a curve is a *Tangent Circle* to the curve. Great and Small Circles are examples of spherical curves.

The arcs are generally spoken of as the *Sides*, and their angles of inclination as the *Angles* of the Spherical figure.

Since, however, three great circles on a sphere intersect one another so as to form *eight* triangles, for the sake of convenience, and to avoid ambiguity, that particular triangle is selected which has two, or if possible, three sides each less than a quadrant.

* Unless otherwise stated.

The three sides and three angles are termed the *Parts* of a spherical triangle; and, as will be hereafter seen, any three whatever given in magnitude are necessary and sufficient to determine the remaining three.

15. **By the Arc connecting two points** on a sphere, unless otherwise expressed, we mean the lesser segment of the great circle passing through the points (see Chap. I. Art. 3).

Examples.

1. Each side of a spherical triangle is less than two right angles.

2. Each angle is less than two right angles. For if possible let ABC be a spherical triangle, having an angle at A greater than two right angles. Continue the arc BA to the point X on the side BC.

Then BX is a semicircle; hence $BX + CX$ is greater; therefore ABC is not a spherical triangle.




Fig. 5.

3. The area of any spherical triangle is less than $2\pi r^2$. (See Art. 8.)

16. **Colunar or Associated Triangles.**—By producing the sides of a triangle below the base, a triangle is formed, having two sides and the opposite angles respectively supplemental to two sides and the opposite angles of the original triangle, while the remaining angle of the one is equal to the remaining angle of the other (being

opposite angles of a lune). Triangles whose parts are so related are termed *Colunar*, and the three triangles colunar with the given one are sometimes termed its *Associated Triangles*. The triangle taken in the first instance is called the *Primitive Triangle*.

Examples.

1. The three associated triangles, taken in pairs, have a side and an opposite angle of the one equal to a side and an opposite angle of the other.
2. When a triangle is equilateral, the colunars are each isosceles.
3. The arcs joining the vertices of the colunar triangles are equal to the sides of the primitive triangle.

17. **Parts of a Spherical Triangle.**—Let ABC be a spherical triangle described on a sphere having its centre at O. Then $OA = OB = OC$ = the radius of the sphere. The arcs BC, CA, and AB are proportional to the angles BOC, COA, and AOB, which they subtend at the centre,




Fig. 6.

and the angles BAC, ABC, and BCA between the sides are equal to the angles between the planes containing the solid angle at O (Chap. I., Art. 6).

The notation for the parts of a spherical triangle adopted throughout will be that of Fig. 6, viz., for the three angles

the letters A, B, and C, and for the opposite sides the small letters a, b, and c, respectively; though, as will be hereafter noticed, the letters A, B, and C, may denote indifferently the number of degrees in the angles or the circular measure of the angles.

Any expression involving one or more parts of a triangle is a *Function* of those parts, and may be represented in a general form such as—for the side a, $f(a)$; for the sides a, b, and c, $f(a, b, c)$; for two sides b and c, and the angle A, $f(b, c, A)$; for all the parts, $f(a, b, c, A, B, C)$.

18. **Sphere of Infinite Radius.**—Let two tangents be drawn to two great circles in their respective planes at the point of intersection; the plane containing both tangents is, with respect to the sphere, the *Tangent Plane* at the point. Now consider the radius of the sphere to increase indefinitely. It is manifest that as the radius increases, each element of surface, and, as a consequence, the whole surface in the neighbourhood of the point, approaches nearer and nearer to the plane; and the great circles described thereon approach nearer and nearer to the tangents. In the limiting case, when the radius of the sphere becomes indefinitely great, and the curvature therefore indefinitely small, the great circles approximate so closely in position to the tangents that they may be regarded as coincident with one another.

Thus the sphere and the great circles degenerate respectively into two* planes (one of which is at infinity), and

* In accordance with analytical geometry, but it is quite sufficient to understand that any finite portion of the surface is a plane.

two pairs of lines (one pair being also at infinity).* But it is to be observed that the angle of inclination of the planes of the great circles alone remains unaltered (*cf.* Chap. I., Art. 6).

Hence the relations involving certain parts of a spherical triangle may be regarded as more general than those involving similar parts of a plane triangle; and by the aid of the expansions

$$\sin a = \frac{a}{r} - \frac{1}{\lfloor 3} \frac{a^3}{r^3} + \frac{1}{\lfloor 5} \frac{a^5}{r^5} + \ldots \text{ to infinity,} \qquad (1)$$

and

$$\cos a = 1 - \frac{1}{\lfloor 2} \frac{a^2}{r^2} + \frac{1}{\lfloor 4} \frac{a^4}{r^4} - \ldots \text{ to infinity,} \qquad (2)$$

* Reference to the Earth, considered as a large sphere, will elucidate this statement.

The surface of a lake or any small portion of water is apparently a perfect plane, although it partakes of the general curvature of the Earth, which may be observed in the case of the ocean, where a distinct curvature is exhibited. It can therefore be readily imagined that if the radius of the Earth were greatly increased, even the surface of the ocean would present no perceptible curvature.

Now take any point on the surface, and draw a plane passing through it and through the centre of the Earth. This plane will cut the surface of the Earth in a great circle (Chap. I., Art. 9). But the surface, at any point, being sensibly a plane, the trace of this great circle on it will be approximately a right line, since the intersection of two planes is a right line. If another great circle be drawn through the point, the two will not differ sensibly from a pair of right lines intersecting at it; and since they both pass through the diametrically opposite point, they will intersect at it also as a pair of right lines, the surface of the Earth being at this point also approximately a plane. Now if the radius of the Earth were indefinitely increased, no curvature whatever would be observed at any part of its surface, and the point diametrically opposite to any selected one would be infinitely distant from it, and the results stated above would follow.

(see Todhunter's *Plane Trig.*), any function of A, B, C, a, b, c the parts of the former, can be transformed into a function of $A, B, C, \alpha, \beta, \gamma$, the parts of the latter. It should, however, be clearly understood that α represents the *length* of an arc subtending an *angle* a, at the centre of a sphere of radius r; and when r becomes infinite the arc α becomes a right line, viz., a side of the corresponding plane triangle.

19. **The Analogy between Problems in Plane and Spherical Trigonometry** referred to in the previous Article may be more clearly illustrated as follows:—

(1). *Any two sides of a triangle are together greater than the third side.* Cf. Euc., Book I., Prop. xx., with Chap. i., Art. 3. See also Euc., XI. xx.

(2). *The angles at the base of an isosceles triangle are equal to one another.* (Euc., I. v.)

For, let tangents be drawn to the equal sides from the extremities of the base of the triangle in the planes of those great circles. It is manifest that the intersection of the tangents is a point on both planes, viz., on their common diameter of intersection; and they are equal to each other (Euc., I. xlvii.), since either $= \sqrt{x^2 - r^2}$, where x is the distance of the point of intersection from the centre of the sphere of radius r. Again, since the tangents to the base at its extremities are equal; therefore, &c. (Euc., I. viii.).

(3). *If two angles of a triangle are equal, the triangle is isosceles.*

(4). *The sum of one pair of opposite angles of a quadrilateral inscribed in a circle is equal to the sum of the remaining pair.*

For, join the pole of the circle to the angles of the quadrilateral forming four isosceles triangles; therefore, &c. (Cf. Prop. 2.)

May this property be regarded as the criterion of a cyclic spherical quadrilateral? *Ans.* Yes.

[Any polygon on the sphere will be cyclic if its vertices are co-planar (Art. 9, chap. i.)]

(5). *The sum of the sides of a quadrilateral is greater than the sum of the diagonals.* (See Prop. I.).

(6). *Only one triangle can be constructed with three given arcs* (*any two of which being supposed greater than the third*). (Cf. Euc. I. XXII.)

(7). *If two triangles have two sides and the included angle of the one equal to two sides and the included angle of the other, each to each, the triangles are equal in every respect.*

For, since two sides of given length intersecting at a given angle determine the vertices of the triangle, therefore the triangle formed by the chords of the sides is determined.




Fig. 7.

The identity of *the chordal triangles** is self-evident, and hence the bases of the spherical triangles are equal since they are the arcs of equal chords; therefore, &c., by the aid of Prop. 6.

* Or, the triangles formed by the chords of the arcs.

It may be remarked that the proof of the identity of the triangles by superposition fails in certain cases on the sphere, *e.g.*, when the non-corresponding sides are equal, as in the accompanying figure, in which case the triangles are *symmetrically* equal.

NOTE.—It may be seen from the foregoing that the vertical angle of the chordal triangle is a function of the two sides and included angle of the spherical triangle (v., chap. viii.).

(8). *The greater side of every triangle is subtended by the greater angle.*

For, let ABC (fig. 8) be the given triangle having the angle B greater than the angle A. Draw a great circle




Fig. 8.

BX making the angle ABX = angle A. Then $BX = AX$. But $BX + CX > BC$; therefore $AC > BC$.

It may be easily shown that the theorems given in Euc. I., Props. XV., XIX., XXI., XXIV., XXV., XXVI., are more generally true for the sphere.

(9). *The internal bisectors of the angles of a triangle, or two external and one internal, are concurrent.* (Euc. IV. v.) [The points of concurrence are the poles of the small circles touching the sides of the triangle.]

(10). *The arcs of great circles bisecting the sides of a triangle at right angles are concurrent.* (Euc. IV. VI.). [The point of concurrence is the pole of a small circle passing through the vertices of the triangle.]

(11). *Given the base and sum of base angles, the external bisector of the vertical angle always passes through a fixed point.* (*Dublin University Exam. Papers.*)

For, let ABC (fig. 9) be a triangle having the given




Fig 9.

base AB. Upon AB construct an isosceles triangle AOB, having each of the base angles OAB and $OBA = \frac{A+B}{2}$. Draw OX and OY secondaries to BC and AC, respectively.

In the two right-angled triangles BOX and AOY, $AO = BO$, $< OAY = < OBX = \frac{B-A}{2}$; therefore $AY = BX$, and $OX = OY$.

Again, compare triangles COX and COY, $OX = OY$, OC is common, and the angles at X and Y are right angles: hence CO is the bisector of the external vertical angle passing through the fixed point O above determined.

(12). Given of a triangle the base and *difference* of base angles, the *internal bisector* of the vertical angle passes through a fixed point. (Cf. Prop. (11)).

20. **Theorem.**—*The great circle bisecting the sides of a triangle intersects the base* 90° *distant from its middle point.*

Let ABC (fig. 10) be the triangle; X, Y, Z, the middle




Fig. 10.

points of the sides; AL, BM, and CN, secondaries drawn through the vertices of the triangle to the great circle XY, passing through the middle points of the sides. Let XY meet the base in the points D and E, which are, therefore, diametrically opposite. O is the pole of XY. (Chap. i. Ex. 8.)

Now in the two triangles ALY and CYN we have $AY = CY$, $\angle AYL = \angle CYN$, and $\angle ALY = \angle CNY$; therefore the triangles are equal in every respect. Hence $AL = CN$. Similarly it can be shown that $BM = CN$.

Therefore

$$AL = BM = CN.$$

Again, in the two triangles ALD and BME it follows easily that $DL = ME$ and $AD = BE$; hence

$$DZ = ZE = 90°.$$

Examples.

1. The arcs DL and XY are complementary.

2. The angle $LAY = S - A$ where $2S = A + B + C$.

3. The triangle AOB is isosceles.

4. The arcs OC and OA are supplementary.

5. Given the base and the sum of the three angles, the locus of the vertex C is a small circle having its pole at O.

For under the given conditions the triangle AOB is fixed; hence also OC, by the aid of Ex. 4; therefore, &c.

Polar Triangles.

Definition.—Two triangles so related that the vertices of the one are the poles of the sides of the other are called, with respect to one another, *Polar Triangles.*

21. **Reciprocal Relations of the Sides and Angles of Polar Triangles.**—*If two triangles be so related that the vertices of the one are the poles of the corresponding sides of the other; then, conversely, the vertices of the latter shall be the poles of the corresponding sides of the former.*

Let the points A, B, and C (fig. 11) be respectively the poles* of $B'C'$, $C'A'$, and $A'B'$ the sides of the triangle

* The poles of the sides of a triangle may be conveniently obtained by drawing great circles through the vertices perpendicular to the opposite sides

$A'B'C'$. Produce the side $B'C'$ (if necessary) to meet AC in the point M, and AB in the point N.




Fig. 11.

Now, since A is the pole of $B'C'$, the arc AB' is a quadrant; and since C is the pole of $A'B'$, the arc CB' is a quadrant; hence B' is the pole of AC: similarly, A' is the pole of BC, and C' the pole of AB; therefore, &c. (Art. 12).

The arcs connecting the corresponding vertices of a triangle and its polar pass through a point.

NOTE.—[For the arcs at X, Y, Z (fig. 10), perpendicular to the sides of the triangle ABC, are concurrent (Art. 19 (10)). Hence, by Art. 20, the perpendiculars of the triangle XYZ, and, therefore, of any spherical triangle, are concurrent; and since AA', BB', CC', are the perpendiculars of the triangle $A'B'C'$; therefore, &c.]

22. **Theorem.**—*The angles of the primitive triangle are the supplements of the corresponding sides of the polar; and conversely.*

Since B' (fig. 11) is the pole of AC, $B'M$ is a quadrant, and for a similar reason $C'N$ is also a quadrant (Chap. i., Ex. 4).

and by taking points on them (measured from the sides towards the vertices) 90° distant from the sides. These points are evidently the required poles.

Hence $MN + B'C' =$ two right angles.

But MN is equal to the angle A; therefore the angle A and the side $B'C'$ are supplemental.

Similarly, the angles B and C are the supplements of the sides $C'A'$ and $A'B'$ respectively; therefore, &c.

NOTE. From the above property Polar Triangles are also termed *Supplemental Triangles;* and if a', b', c', A', B', C', represent the parts of the triangle $A'B'C'$ (fig. 11), it follows that

$$A + a' = B + b' = C + c' = \pi.$$

Also

$$A' + a = B' + b = C' + c = \pi.$$

These results are of the greatest importance, inasmuch as any theorem which holds good between the sides and angles (e.g. a, b, c, A, B) of a spherical triangle necessarily involves a reciprocal or supplemental theorem, involving the opposite angles and sides (viz. A, B, C, a, b), and which may be derived from it by changing the sides into the supplements of the corresponding angles, and the angles into the supplements of the corresponding sides.

23. The results obtained in the previous Article may be further exemplified, as follows:—

(1). It has been seen that two sides of a triangle are greater than the third: thus $b + c > a$.

Applying this inequality to the sides of the supplemental triangle, it follows that

$$\pi - B + \pi - C > \pi - A.$$

Hence, $B + C - A < \pi$, i. e. $S - A < \frac{\pi}{2}$. (1)

Similarly, $C + A - B < \pi$, i. e. $S - B < \frac{\pi}{2}$. (2)

And $A + B - C < \pi$, i. e. $S - C < \frac{\pi}{2}$. (3)

Or, *the difference between any two angles of a spherical triangle is less than the supplement of the third angle.*

Particular cases of the inequalities (1), (2), and (3) are worthy of notice.

a. Suppose, for example, that the primitive triangle is right-angled at C, it follows from (3) that

$$A + B < \frac{3\pi}{2};$$

or, *the sum of the angles of a right-angled triangle is less than four right angles.*

β. It likewise follows, by the aid of (2), that

$$A - B < \frac{\pi}{2};$$

or, *the difference of the oblique angles of a right-angled triangle is less than a right angle.*

(2). Prop. (2), Art. 19, involves the converse theorem, viz., if two angles of a triangle are equal, the sides subtending them are also equal. [See Prop. (3), Art. 19.]

(3). Prop. (8), Art. 19, involves the converse theorem, viz., the greater angle is subtended by the greater side.

(4). Prop. (7), Art. 19, involves the supplemental theorem, viz., If two triangles have two angles and a side of the one equal to two angles and a side of the other, the sides

being similarly situated with respect to the equal angles, the triangles are equal in every respect.

(5). In any triangle if $C = A + B$, then $C - A$ and $C - B$ are each less than a right angle.

For $C - A = B$, and $C - B = A$; but $B + C - A < \pi$; hence $B < \frac{\pi}{2}$, and $A < \frac{\pi}{2}$; therefore, &c.

Examples.

1. Given two angles of a triangle to be 45° and 120°, find the maximum value of the third angle.

Let A be the third angle, then by aid of (1),

$$A + 120 - 45 < 180°.$$

Hence,

$$A < 105°.$$

2. The angles of a triangle are A, 30° and 150°; find the maximum value of A. *Ans.* $A < 60°$.

[Proof as before.]

3. If the angles are A, 20° and 110°, find the maximum value of A. *Ans.* $A < 90°$.

4. If the difference between any two angles of a triangle is equal to 90°, the remaining angle is less than 90°. [Cf. Art. 23, β.]

5. If the primitive triangle be equilateral or isosceles, the supplemental triangle is equilateral or isosceles.

24. **Theorem.**—*If two sides of a triangle are supplemental, the opposite angles are supplemental.*

For in the triangle ABC (fig. 12), if $AC + BC = \pi$, since the arc ABA' is a semicircle, then will

$$AC = A'B, \text{ and } A'C = AB.$$

Therefore the two triangles ABC and $A'BC$ are equal in every respect; therefore, &c.

Examples.

1. If two sides of a triangle are supplemental, two sides of the polar triangle are likewise supplemental.

2. In fig. 12, the arc joining the middle point M of BC with A or A' is a quadrant.

[The triangles $A'BM$ and ACM are equal in every respect.]

3. Given the base of a spherical triangle, and the sum of the sides equal to two right angles, find the locus of the vertex. *Ans.* A great circle, having the middle point of the base as pole (Ex. 2).

25. **Angular Limits of the Sides and Angles of a Spherical Triangle.**—It has been said that each side and each angle of a spherical triangle may have any values between 0° and π. It is, therefore, manifest that the sum of the angles cannot exceed 3π; and it will be here shown that the sum of the sides cannot exceed 2π. For in the triangle $A'BC$ (fig. 12),




Fig. 12.

$$BC < A'C + A'B.$$

To each add $AC + AB$.

Hence,

$$BC + CA + AB < AC + A'C + AB + A'B < 2\pi.$$

Hence $a + b + c$ *can have any values between* $0°$ *and* 2π, *and* $A + B + C$ *can have any values between* π *and* 3π.

The latter may be seen otherwise, thus :—

Since $0°$, $a + b + c$, 2π, are in ascending order of magnitude; hence, for the supplemental triangle,

$$0°, \quad \pi - A + \pi - B + \pi - C, \quad 2\pi$$

are likewise in ascending order of magnitude; therefore, &c.

Examples.

1. Any side of a triangle is greater than the difference between the other two sides.

2. When does the primitive triangle coincide with the supplemental?

Ans. When its sides and angles are each $\frac{\pi}{2}$.

NOTE.—Such a triangle is called a *Quadrantal Triangle*.

3. Assuming the earth to be a sphere, is the area included between two meridians and a parallel of latitude a spherical triangle? Give a reason for your answer. (*Sc. and Art Exam. Papers.*)

4. The sides a and b of a triangle are produced both ways to points x and x', y and y', $90°$ distant from the middle point of the sides. Show that, if secondaries to the sides from x and y intersect in P, and from x' and y' in P', the points P and P' are diametrically opposite.

5. The triangle $P'Q'R'$ is the polar triangle of that formed by joining the middle points of the sides of ABC, where Q' and R' are formed by similarly producing the sides c and a, a and b, through the extremities of the remaining side.

6. If a, b, c, d be the sides of a spherical quadrilateral inscribed in a circle, δ and δ' its diagonals, show that

$$\sin \tfrac{1}{2}a \sin \tfrac{1}{2}c + \sin \tfrac{1}{2}b \sin \tfrac{1}{2}d = \sin \tfrac{1}{2}\delta \sin \tfrac{1}{2}\delta'.$$

[Apply Ptolemy's Theorem to the chordal quadrilateral, and notice that the chord of the side a is proportional to $\sin \frac{1}{2}a$, &c.]

CHAPTER III.

FUNDAMENTAL RELATIONS BETWEEN THE PARTS OF A SPHERICAL TRIANGLE.

SECTION I.

26. *Having given the sides of a spherical triangle, to determine the cosines of the angles.*




Fig. 13.

Let ABC be a spherical triangle; a, b, and c, its sides; O the centre of the sphere. Draw CD and CE tangents to a and b at C.*

Since these tangents lie in the planes of the circles to which they are drawn, they will meet the radii OB and

* The same result will be obtained if, instead of drawing tangents at C, any point C' be taken on OC, and lines $C'D'$ and $C'E'$ be drawn perpendicular to it, meeting OB and OA in D' and E', respectively.

OA in D and E respectively, and the angle between EC and CD is equal to the angle C (Chap. i., Art. 6).

Now, from the two triangles ECD and EOD we have

$$DE^2 = CD^2 + CE^2 - 2CD \,.\, CE \,.\, \cos C \;\ldots \qquad (\alpha)$$

Also $$DE^2 = OD^2 + OE^2 - 2OD \,.\, OE \,.\, \cos c \;\ldots; \qquad (\beta)$$

and since the angles OCE and OCD are each 90°,

$$OC^2 = OD^2 - CD^2 = OE^2 - CE^2.$$

Hence, on subtracting (α) from (β),

$$0 = OC^2 + CD \,.\, CE \cos C - OD \,.\, OE \,.\, \cos c;$$

or,

$$\cos c = \frac{OC}{OD} \cdot \frac{OC}{OE} + \frac{CD}{OD} \cdot \frac{CE}{OE} \cdot \cos C;$$

$$\left.\begin{aligned} &\text{therefore,} && \cos c = \cos a \cos b + \sin a \sin b \cos C. \\ &\text{Similarly,} && \cos b = \cos c \cos a + \sin c \sin a \cos B, \\ &\text{and} && \cos a = \cos b \cos c + \sin b \sin c \cos A. \end{aligned}\right\} \qquad (1)$$

From these formulæ we obtain the angles in terms of the sides. Thus:

$$\left.\begin{aligned} \cos A &= \frac{\cos a - \cos b \cos c}{\sin b \sin c} \;\ldots \\ \cos B &= \frac{\cos b - \cos c \cos a}{\sin c \sin a} \;\ldots \\ \cos C &= \frac{\cos c - \cos a \cos b}{\sin a \sin b} \;\ldots \end{aligned}\right\} \qquad (2).$$

These results are of special importance, and the student should make himself perfectly familiar with them at the outset.

They apply to all spherical triangles, whether the arcs be greater or less than quadrants.

27. **Generalization.**—In the preceding Article it is plain, on referring to the figure, that the proof given only applies to a spherical triangle having the sides a and b each less than $\frac{\pi}{2}$. For, in the two triangles OCE and OCD, each right-angled at C, the angles COE and COD are acute, and therefore the arcs a and b subtended by them on the sphere are less than $\frac{\pi}{2}$. No such restriction has been placed on the limits of the remaining side c: it may, therefore, be greater, equal to, or less than $\frac{\pi}{2}$.

It therefore remains to be shown that the foregoing theorem is applicable when *either of* or *both* the sides a and b are greater than quadrants.

(1). *In the former case* let a be greater than, and b less than, a quadrant.




Fig. 14.

Let a and c (fig. 14) be produced to meet in B'. Then in the triangle $AB'C$, by the preceding it follows that

$$\cos AB' = \cos B'C \cos AC + \sin B'C \sin AC \cos ACB';$$

therefore,

$$\cos a = \cos a \cos b + \sin a \sin b \cos C.$$

(2). *In the latter case* let a and b (fig. 15) be each greater than a quadrant




Fig. 15.

Then in the triangle ABC' we have

$$\cos c = \cos BC' \cos AC' + \sin BC' \sin AC' \cos C;$$

or,

$$\cos c = \cos a \cos b + \sin a \sin b \cos C.$$

The remaining cases, when either of or both the sides a and b are quadrants, are left as easy exercises for the student. The formulæ referred to are, therefore, applicable to all triangles drawn on the surface of a sphere.

Exercises on Fundamental Formulæ (1) *and* (2).

$$\text{Let } C = \frac{\pi}{2}.$$

(1). Hence $\cos c = \cos a \cos b.$

(2). Again, from the formula,

$$\cos A = \frac{\cos a - \cos b \cos c}{\sin b \sin c}.$$

On substituting the value of $\cos a$ from Ex. (1) we have

$$\cos A = \frac{\tan b}{\tan c}.$$

Similarly,

$$\cos B = \frac{\tan a}{\tan c}.$$

(3). On substituting the value of $\cos b$ from Ex. (1) we have from (2), after reduction,

$$\sin A = \frac{\sin a}{\sin c}.$$

Similarly,

$$\sin B = \frac{\sin b}{\sin c}.$$

(4). From Examples (2) and (3) we have, by division,

$$\tan a = \tan A \sin b,$$

and

$$\tan b = \tan B \sin a.$$

(5). From (4) we have

$$\tan a \tan b = \tan A \tan B \cdot \sin a \sin b;$$

or,

$$\cot A \cot B = \cos a \cos b = \cos c.$$

(6). From (2) and (3) we have

$$\cos A = \cos a \sin B.$$

The above six formulæ, connecting the sides and angles of a right-angled triangle, are expressed in general terms thus:—

(1). cos (hypotenuse) = product of cosines of sides.
(2). cos (angle) = tangent (adjacent side) ÷ tan (hypotenuse).
(3). sin (angle) = sin (opposite side) ÷ sin (hypotenuse).
(4). tan (side) = tan (opposite angle) × sin (remaining side).
(5). cos (hypotenuse) = product of cotangents of base angles.
(6). cos (angle) = cos (opposite side) × sin (remaining angle).

(7). Let $b = c = \frac{\pi}{2}$, and the expression

$$\cos a = \cos b \cos c + \sin b \sin c \cos A$$

reduces to

$$\cos a = \cos A, \qquad (\textit{Vide} \text{ Art. 6, fig. 2.})$$

as is otherwise evident, since in this case C is the pole of AB.

(8). Let $$a = b = c;$$

then $$\cos a = \cos^2 a + \sin^2 a \cos A;$$

or, $$\cos a (1 - \cos a) = (1 - \cos^2 a) \cos A.$$

Hence, $$\sec A = 1 + \sec a,$$

a constant relation between the angle and side of an equilateral spherical triangle.

(9). Given a side or angle of an equilateral triangle; solve it completely, and deduce the limiting values of the sides and angles of an equilateral triangle. Apply Ex. (8).

(10) Given the latitudes and difference of longitude of two places on the Earth's surface, compute the distance between them.

(*Science and Art Exam. Papers*).

28. **Analogous Formula in Plano.**—Supposing the radius of the sphere to be *indefinitely great*, show that the formula

$$\frac{\cos a - \cos b \cos c}{\sin b \sin c} = \cos A$$

degenerates into

$$\frac{\beta^2 + \gamma^2 - a^2}{2\beta\gamma} = \cos A,$$

where A is an angle, and a, β, and γ the sides of a plane triangle.

Expanding $\cos a$, &c., by the aid of formulæ, Chap. II., Art. 18, and neglecting all powers of the radius above the second, we have

$$\frac{\left(1 - \frac{a^2}{\lfloor 2r^2}\right) - \left(1 - \frac{\beta^2}{\lfloor 2r^2}\right)\left(1 - \frac{\gamma^2}{\lfloor 2r^2}\right)}{\frac{\beta}{r} \cdot \frac{\gamma}{r}} = \cos A,$$

which reduces to

$$\frac{\beta^2 + \gamma^2 - a^2}{2\beta\gamma} = \cos A.$$

29. **Legendre's Theorem.**—*If the sides of a spherical triangle be small compared with the radius of the sphere, then each angle of the spherical triangle exceeds by one-third of the spherical excess the corresponding angle of the plane triangle, the sides of which are of the same length as the arcs of the spherical triangle.*

In the foregoing example the powers of the radius higher than the *second* were neglected. Suppose, however, a closer approximation to be made, and the powers of r above the *fourth* neglected.

Then

$$\cos A = \frac{1 - \frac{a^2}{2r^2} + \frac{a^4}{24r^4} - \left(1 - \frac{\beta^2}{2r^2} + \frac{\beta^4}{24r^4}\right)\left(1 - \frac{\gamma^2}{2r^2} + \frac{\gamma^4}{24r^4}\right)}{\frac{\beta}{r}\left(1 - \frac{\beta^2}{6r^2}\right)\frac{\gamma}{r}\left(1 - \frac{\gamma^2}{6r^2}\right)}$$

$$= \frac{\frac{\beta^2 + \gamma^2 - a^2}{2r^2} + \frac{a^4 - \beta^4 - \gamma^4 - 6\beta^2\gamma^2}{24r^4}}{\frac{\beta\gamma}{r^2}\left(1 - \frac{\beta^2 + \gamma^2}{6r^2}\right)}.$$

But, by the aid of the Binomial Theorem we have

$$\frac{1}{1 - \frac{\beta^2 + \gamma^2}{6r^2}} = \left(1 - \frac{\beta^2 + \gamma^2}{6r^2}\right)^{-1} = 1 + \frac{\beta^2 + \gamma^2}{6r^2} \text{ nearly.}$$

Hence

$$\cos A = \frac{\left\{\frac{\beta^2+\gamma^2-\alpha^2}{2r^2} + \frac{\alpha^4-\beta^4-\gamma^4-6\beta^2\gamma^2}{24r^4}\right\}\left\{1+\frac{\beta^2+\gamma^2}{6r^2}\right\}}{\frac{\beta\gamma}{r^2}}$$

$$= \frac{\beta^2+\gamma^2-\alpha^2}{2\beta\gamma} - \frac{2\beta^2\gamma^2+2\gamma^2\alpha^2+2\alpha^2\beta^2-\alpha^4-\beta^4-\gamma^4}{24\beta\gamma r^2}.$$

Now consider a plane triangle of sides α, β, and γ; angles A', B', and C'; and area $= \Delta = \frac{1}{2}\beta\gamma \sin A'$.

The above equation reduces to the form (Plane Trig.)

$$\cos A = \cos A' - \frac{\sin^2 A' \cdot \beta\gamma}{6r^2}.$$

Let the excess of A over A' be denoted by θ, θ being a very small quantity; then $A = A' + \theta$, and

$$\cos A = \cos A' - \theta \sin A' \text{ nearly};$$

therefore,

$$\cos A' - \frac{\sin^2 A'}{6r^2}\beta\gamma = \cos A' - \theta \sin A';$$

or,

$$\theta = \frac{\Delta}{3r^2}.$$

We have therefore,

$$A = A' + \frac{\Delta}{3r^2},$$

$$B = B' + \frac{\Delta}{3r^2},$$

$$C = C' + \frac{\Delta}{3r^2}.$$

BIBLIOTHECA BODLEIANA

Whence

$$A + B + C - A' - B' - C' = \frac{\Delta}{r^2}.$$

or,

$$A + B + C - \pi = \frac{\Delta}{r^2}.$$

But the expression $A + B + C - \pi$ denotes the excess of the sum of the angles of a spherical triangle over the sum of the angles of a plane triangle (both being expressed in circular measure), and is therefore called the *Spherical Excess*; therefore, &c.

Section II.

30. *Expression for the sine of an angle of a spherical triangle in terms of the trigonometrical functions of the sides.*

We have (Art. 26),

$$\cos A = \frac{\cos a - \cos b \cos c}{\sin b \sin c};$$

therefore

$$\sin^2 A = 1 - \left(\frac{\cos a - \cos b \cos c}{\sin b \sin c}\right)^2$$

$$= \frac{\sin^2 b \sin^2 c - (\cos a - \cos b \cos c)^2}{\sin^2 b \sin^2 c}$$

$$= \frac{(1 - \cos^2 b)(1 - \cos^2 c) - (\cos a - \cos b \cos c)^2}{\sin^2 b \sin^2 c}$$

$$= \frac{1 - \cos^2 a - \cos^2 b - \cos^2 c + 2 \cos a \cos b \cos c}{\sin^2 b \sin^2 c}$$

$$= \frac{4n^2}{\sin^2 b \sin^2 c},$$

where

$$4n^2 = 1 - \cos^2 a - \cos^2 b - \cos^2 c + 2 \cos a \cos b \cos c;$$ *

hence $$\sin A = \frac{2n}{\sin b \sin c} \quad \ldots$$

Similarly $$\sin B = \frac{2n}{\sin c \sin a} \quad \ldots \qquad (3)$$

and $$\sin C = \frac{2n}{\sin a \sin b} \quad \ldots$$

* This function has been called the sine of the solid angle that the triangle subtends at the centre of the sphere (*vide* Salmon's *Geometry of Three Dimensions*, Art. 54), and may be written in the determinant form

$$\begin{vmatrix} 1, & \cos c, & \cos b \\ \cos c, & 1, & \cos a \\ \cos b, & \cos a, & 1 \end{vmatrix}$$

31. **Rule of Sines.**—From the value of $\sin A$ in the previous Article we have at once

$$\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c} = \frac{2n}{\sin a \sin b \sin c}. \qquad (4)$$

Hence * *the sines of the angles of a spherical triangle are proportional to the sines of the opposite sides.*

The corresponding theorem *in plano* is, "The Sines of the Angles of a Plane Triangle are proportional to the opposite Sides" (Griffin's *Plane Trig.*, p. 30).

Examples.

1. Given the base c and the function n, find the locus of the vertex. (*Q. U. I., Exam. Papers.*)

Since $$\sin A = \frac{2n}{\sin b \sin c},$$

we have given $\sin b \sin A$, which, by the aid of Art. 31, or 27, Ex. 3, is equal to $\sin P$, where P is the perpendicular from the vertex on the base. The locus is, therefore, a small circle having the same poles as the base.

2. If α, β, γ, be the perpendiculars of a triangle, prove that

$$\sin a \sin \alpha = \sin b \sin \beta = \sin c \sin \gamma = 2n.$$

For (Art. 27, Ex. 3),

$$\sin \alpha = \sin b \sin C, \quad \sin \beta = \sin c \sin A, \quad \sin \gamma = \sin a \sin B;$$

therefore, &c. (Art. 30 (3)).

Remark.—The rule of sines follows at once from the equation

$$\sin a = \sin b \sin C = \sin c \sin B; \therefore \frac{\sin B}{\sin C} = \frac{\sin b}{\sin c}.$$

Therefore, &c.

* Or thus:—take P, the pole of the side c, and apply equation (1) Art. 26, to the triangles ACP and BCP. Then $\cos CP = \sin a \sin B = \sin b \sin A$. Therefore, &c.

3. In Ex. 2 show that

$$\sin a \cos \alpha = \sqrt{\cos^2 b + \cos^2 c - 2 \cos a \cos b \cos c},$$

with similar expressions for $\cos \beta$ and $\cos \gamma$.

Since

$$\sin \alpha = \sin b \sin C = \frac{\sqrt{1 - \cos^2 a - \cos^2 b - \cos^2 c + 2 \cos a \cos b \cos c}}{\sin a}.$$

Hence $\sin^2 a \sin^2 \alpha$ or $\sin^2 a - \sin^2 a \cos^2 \alpha$

$$= 1 - \cos^2 a - \cos^2 b - \cos^2 c + 2 \cos a \cos b \cos c;$$

therefore, &c.

4. The sines of the segments of the base of a triangle made by the bisectors of the internal and external vertical angle are to one another as the sines of the adjacent sides. (Cf. Euc. VI., III.)

5. The bisector of the base divides the vertical angle into two parts, the ratio of whose sines is the inverse ratio of the sines of the adjacent sides.

6. If M be the middle point of the base AB of a triangle ABC and M' * a point on the base such that the angle ACM is equal to the angle BCM' prove that

$$\sin AM' : \sin BM' :: \sin^2 b : \sin^2 a.$$

7. If p_1, p_2, be the perpendiculars drawn from the mid-point M of the base of a spherical triangle on the great circle bisectors of the vertical angle A, and p_3 the perpendicular from A on the great circle perpendicular to the base through M, prove—

(1). $$\sin p_1 \sin p_2 = \tfrac{1}{2} \sin p_3 \sin \frac{a}{2} \sin (B + C).$$

(2). Give the analogous theorem for a plane triangle.

(*Educational Times*, August, 1884.)

Let y and $\overset{\curvearrowleft}{x}$ be the points where the external and internal bisectors meet the perpendicular through M; then

$$\frac{\sin p_1 \sin p_2}{\sin p_3} = \frac{\sin Mx \sin x \,.\, \sin My \sin y}{\sin Ax \,.\, \sin x} = \frac{\sin Mx \,.\, \sin My}{\sin xy};$$

therefore, &c.

(2). *Ans.* $\dfrac{4 p_1 p_2}{p_3} = a \sin A.$

* The arc CM' is called a *Symmedian* of the triangle ABC.

32. **Geometrical Proof of the Rule of Sines.**— *To show geometrically that the sines of the angles of a spherical triangle are proportional to the sines of the opposite sides.*

Let ABC (fig. 16) be a spherical triangle, O the centre of the sphere. From any point P in OC let fall a perpen-




Fig. 16.

dicular PD on the plane AOB. Join OD, and let fall DE perpendicular on OA. Join PE. PD is perpendicular to all lines drawn through D in the plane AOB.

Now

$$OP^2 = OD^2 + PD^2 = OE^2 + DE^2 + PD^2 = OE^2 + PE^2;$$

therefore PEO is a right angle.

Now, since PE and DE are drawn in the planes OAC and OAB perpendicular to their line of intersection OA,

$$\angle PED = \angle A;$$

therefore

$$\sin A = \frac{PD}{PE}, \quad \sin b = \frac{PE}{OP},$$

and

$$\sin A \sin b = \frac{PD}{OP}.$$

Similarly, by letting fall a perpendicular DF on OB we get

$$\sin B \sin a = \frac{PD}{OP} = \sin A \sin b;$$

hence

$$\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c}.$$

33. *Expressions for the sine, cosine, and tangent of half an angle of a triangle as functions of the sides.*

We have

$$2\cos^2\frac{A}{2} = 1 + \cos A$$

$$= 1 + \frac{\cos a - \cos b \cos c}{\sin b \sin c} \qquad \text{(Art. 26 (2)).}$$

$$= \frac{\cos a - \cos(b+c)}{\sin b \sin c} = \frac{2\sin\frac{(a+b+c)}{2}\sin\frac{(b+c-a)}{2}}{\sin b \sin c}$$

$$= \frac{2\sin s \sin(s-a)}{\sin b \sin c}, \quad \text{where } 2s = a + b + c;$$

hence

$$\cos\frac{A}{2} = \sqrt{\frac{\sin s \sin(s-a)}{\sin b \sin c}}$$

Similarly,

$$\cos\frac{B}{2} = \sqrt{\frac{\sin s \sin(s-b)}{\sin c \sin a}}$$

and

$$\cos\frac{C}{2} = \sqrt{\frac{\sin s \sin(s-c)}{\sin a \sin b}}$$

(5)

Again,

$$2\sin^2\frac{A}{2} = 1 - \cos A$$

$$= 1 - \frac{\cos a - \cos b \cos c}{\sin b \sin c}$$

$$= \frac{\cos(b-c) - \cos a}{\sin b \sin c}$$

$$= \frac{2\sin(s-b)\sin(s-c)}{\sin b \sin c};$$

hence

$$\sin\frac{A}{2} = \sqrt{\frac{\sin(s-b)\sin(s-c)}{\sin b \sin c}}$$

Similarly,

$$\sin\frac{B}{2} = \sqrt{\frac{\sin(s-c)\sin(s-a)}{\sin c \sin a}} \qquad (6)$$

and

$$\sin\frac{C}{2} = \sqrt{\frac{\sin(s-a)\sin(s-b)}{\sin a \sin b}}$$

From the above results we obtain

$$\tan\frac{A}{2} = \frac{\sin\frac{A}{2}}{\cos\frac{A}{2}} = \sqrt{\frac{\sin(s-b)\sin(s-c)}{\sin s \sin(s-a)}},$$

with similar expressions for $\tan\frac{B}{2}$ and $\tan\frac{C}{2}$. (7)

NOTE.—

$$\sin A = 2\sin\frac{A}{2}\cos\frac{A}{2} = \frac{2}{\sin b \sin c}\sqrt{\sin s \sin(s-a)\sin(s-b)\sin(s-c)};$$

(From (5) and (6)).

hence

$$\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c} = \frac{2}{\sin a \sin b \sin c}\sqrt{\sin s \sin(s-a)\sin(s-b)\sin(s-c)}.$$

Comparing this expression for $\sin A$ with that derived in Art. 30, we see that

$$\begin{aligned}&\sin s . \sin(s-a) . \sin(s-b) . \sin(s-c)\\ &= \tfrac{1}{4}(1-\cos^2 a - \cos^2 b - \cos^2 c + 2\cos a \cos b \cos c)\\ &= n^2,\end{aligned}$$

an *identity* which can be readily proved by direct multiplication of the four factors, $\sin s$, $\sin(s-a)$, $\sin(s-b)$, and $\sin(s-c)$.

Thus,

$$2\sin s \sin(s-a) = \cos a - \cos(b+c) \text{ and } 2\sin(s-b)\sin(s-c)$$
$$= \cos(b-c) - \cos a;$$

therefore, &c.

Examples.

1. Find the conditions that

$$1 - \cos^2 a - \cos^2 b - \cos^2 c + 2\cos a \cos b \cos c \equiv 0.$$

2. Prove that

$$8n^3 = \sin^2 a \sin^2 b \sin^2 c \sin A \sin B \sin C.$$

3. Prove that

$$\cot\frac{A}{2} : \cot\frac{B}{2} : \cot\frac{C}{2} = \sin(s-a) : \sin(s-b) : \sin(s-c).$$

4. Prove that

$$\tan\frac{B}{2}\tan\frac{C}{2} = \frac{\sin(s-a)}{\sin s}.$$

5. Prove that

$$\sin(s-a) = \frac{\sin\frac{B}{2}\sin\frac{C}{2}}{\sin\frac{A}{2}}\sin a,$$

with similar expressions for $\sin(s-b)$ and $\sin(s-c)$.

6. Prove that

$$\sin s = \frac{\cos\frac{B}{2}\cos\frac{C}{2}}{\sin\frac{A}{2}}\sin a.$$

7. Prove that

$$\sin a \sin b \sin c \,.\, \sin\frac{A}{2} \,.\, \sin\frac{B}{2} \,.\, \sin\frac{C}{2} = \frac{n^2}{\sin s}.$$

8. Prove that

$$\sin\frac{A}{2}\cos\frac{B}{2} = \frac{\sin(s-b)}{\sin c}\cos\frac{C}{2}.$$

NOTE.*—By the aid of a similar equation,

$$\cos\frac{A}{2}\sin\frac{B}{2} = \frac{\sin(s-a)}{\sin c}\cos\frac{C}{2}.$$

We have, on adding and subtracting,

$$\sin\frac{A+B}{2}\cos\frac{c}{2} = \cos\frac{a-b}{2}\cos\frac{C}{2}. \qquad (\alpha)$$

$$\sin\frac{A-B}{2}\sin\frac{c}{2} = \sin\frac{a-b}{2}\cos\frac{C}{2}. \qquad (\beta)$$

Again, from Examples 5 and 6, we have

$$\cos\frac{A+B}{2}\cos\frac{c}{2} = \cos\frac{a+b}{2}\sin\frac{C}{2}. \qquad (\gamma)$$

$$\cos\frac{A-B}{2}\sin\frac{c}{2} = \sin\frac{a+b}{2}\sin\frac{C}{2}. \qquad (\delta)$$

four equations which will be afterwards proved geometrically.

* The equations α, β, γ, δ, are known as Gauss's Formulae, and the equations α', β', γ', δ' (Ex. 9), are known as Napier's Analogies. They will all be further discussed later on.

9. Prove that

$$(\alpha'). \quad \tan\frac{A+B}{2} = \frac{\cos\frac{a-b}{2}}{\cos\frac{a+b}{2}} \cot\frac{C}{2}.$$

$$(\beta'). \quad \tan\frac{A-B}{2} = \frac{\sin\frac{a-b}{2}}{\sin\frac{a+b}{2}} \cot\frac{C}{2}.$$

$$(\gamma'). \quad \tan\frac{a+b}{2} = \frac{\cos\frac{A-B}{2}}{\cos\frac{A+B}{2}} \tan\frac{c}{2}.$$

$$(\delta'). \quad \tan\frac{a-b}{2} = \frac{\sin\frac{A-B}{2}}{\sin\frac{A+B}{2}} \tan\frac{c}{2}.$$

[These results follow from Ex. 8.]

Supplemental Theorems.

34. *Having given the angles of a spherical triangle, to determine the cosines of the sides* (cf. Art. 26).

In the formula

$$\cos a = \cos b \cos c + \sin b \sin c \cos A,$$

change the sides into the supplements of the corresponding angles, and the angle into the supplement of the corresponding side (Note, Art. 22), and we get

$$-\cos A = \cos B \cos C - \sin B \sin C \cos a.$$

Therefore,

$$\cos a = \frac{\cos A + \cos B \cos C}{\sin B \sin C}$$

Similarly,

$$\cos b = \frac{\cos B + \cos C \cos A}{\sin C \sin A}$$

and

$$\cos c = \frac{\cos C + \cos A \cos B}{\sin A \sin B}$$

(8)

All the results obtained in Art. 26, as particular cases of formula (1) of this Chapter, may also be deduced as particular cases of formula (8), by supposing the triangle to be equilateral or right-angled, as the case may require.

35. **Analogous Formula in Plane.**—If we suppose the radius of the sphere to be indefinitely great, we have (Chap. II., Art. 18) $\cos a = 1$.

Therefore,

$$\cos A + \cos B \cos C = \sin B \sin C,$$

from which it follows at once that

$$A + B + C = \pi.$$

Examples.

1. In any triangle show that

$$\frac{\cos A + \cos B}{1 - \cos C} = \frac{\sin(a+b)}{\sin c}.$$

If the internal bisector of the angle C make an angle θ with the opposite side, we have

$$\cos\theta + \cos A \cos\frac{C}{2} = \sin A \sin\frac{C}{2} \cos b;$$ by Art. 26 (1).

and also

$$-\cos\theta + \cos B \cos\frac{C}{2} = \sin B \sin\frac{C}{2} \cos a.$$

Adding these equations, we eliminate θ, and deduce the above expression.

2. In any triangle show that

$$\Sigma \frac{\cos A + \cos B}{1 - \cos C} \sin(a - b) \sin c \equiv 0.$$

[This result may be obtained at once from Ex. 1].

3. Prove the relation

$$\frac{\cos A - \cos B}{1 + \cos C} = \frac{\sin(a \sim b)}{\sin c}.$$

[This follows, as in Ex. 1, by drawing the external, instead of the internal, bisector of C].

4. Prove the relation

$$\Sigma \frac{\cos A - \cos B}{1 + \cos C} \sin(a + b) \sin c \equiv 0.$$

5. If a great circle passes through the vertex C, making angles α and β with the sides a and b, the angle θ, which it makes with the side c, is given by the equation

$$\cos A \sin \alpha - \cos B \sin \beta = \pm \cos \theta \sin C.$$

For

$$\cos A + \cos \theta \cos \beta = \sin \theta \sin \beta \cos \delta \text{ (Art. 26 (1))}.$$

where δ is the intercept between the vertex and the base; also

$$\cos B - \cos \theta \cos \alpha = \sin \theta \sin \alpha \cos \delta,$$

eliminate δ between these equations; therefore, &c.

6. If through any point P on a sphere three great circles be drawn, cutting the sides of a triangle at angles X, Y, Z; X_1, Y_1, Z_1; X_2, Y_2, Z_2, respectively; prove the following *determinant** relation :—

$$\begin{vmatrix} \cos X & \cos Y & \cos Z \\ \cos X_1 & \cos Y_1 & \cos Z_1 \\ \cos X_2 & \cos Y_2 & \cos Z_2 \end{vmatrix} \equiv 0.$$

Let the three concurrent arcs make angles α, β, and $\alpha + \beta$ with each other.

* For a knowledge of Determinants, *vide* Burnside and Panton's *Theory of Equations*, Chap. xi.

Since the side a is cut at angles X, X_1, and X_2; by Ex. 5,

$$\cos X \sin a - \cos X_2 \sin \beta = \cos X_1 \sin (a + \beta).$$

Similarly,

$$\cos Y \sin a - \cos Y_2 \sin \beta = \cos Y_1 \sin (a + \beta),$$

and

$$\cos Z \sin a - \cos Z_2 \sin \beta = \cos Z_1 \sin (a + \beta).$$




Fig. 17.

Eliminating a and β from the three equations, the above result easily follows.

7. Having given that the sides of a triangle are each $\frac{\pi}{3}$, find the sides of the supplemental triangle.

[Since the angles of the latter triangle are each $\frac{2\pi}{3}$, $\cos a = -\frac{1}{3}$].

8. Find a relation connecting the angles of a triangle if one side a is a quadrant. *Ans.* $\cos A + \cos B \cos C = 0$.

9. Given A, B, C, find the angle θ which the bisector of the vertical angle makes with the base.

If i be the length of the internal bisector, we have

$$\cos i = \frac{\cos A + \cos \theta \cos \frac{C}{2}}{\sin \theta \sin \frac{C}{2}} = \frac{\cos B - \cos \theta \cos \frac{C}{2}}{\sin \theta \sin \frac{C}{2}};$$

therefore,

$$\cos \theta = \frac{\cos A - \cos B}{2 \cos \frac{C}{2}}$$

10. If θ' denote the angle made by the external bisector of the vertical angle with the base, show that

$$\cos\theta' = \frac{\cos A + \cos B}{2\sin\frac{C}{2}}. \qquad \text{(Cf. Ex. 9.)}$$

11. From any three points A, B, and C, on a great circle, secondaries AA', BB', and CC' are drawn to another great circle; prove that the *algebraic* sum

$$\sin B'C' \cos A + \sin C'A' \cos B + \sin A'B' \cos C$$

is equal to zero. (Apply Ex. 5.)

12. The extremities of the diameter AB of a small circle are joined with a point C on the circle; prove that the angles subtended at the pole by the joining arcs AC and BC are $\cos^{-1}\frac{\sin(A \sim B)}{\sin C}$. (Apply Ex. 5.)

13. Given the base c and $\frac{\cos A}{\cos B} = -\cos C$, find the locus of the vertex [$a = \frac{\pi}{2}$ by (Ex. 8). Hence the locus is a great circle, having the vertex B for pole].

36. *Expression for the side of a Spherical Triangle in terms of the Trigonometrical Functions of the Angles* (cf. Art. 33).

We have

$$2\sin^2\frac{a}{2} = 1 - \cos a = 1 - \frac{\cos A + \cos B\cos C}{\sin B\sin C} \qquad \text{(Art. 34).}$$

$$= -\frac{\cos A + \cos(B+C)}{\sin B\sin C} = -\frac{2\cos S.\cos(S-A)}{\sin B\sin C},$$

where

$$2S = A + B + C.$$

Therefore,

$$\sin\frac{a}{2} = \sqrt{\frac{-\cos S\cos(S-A)}{\sin B\sin C}}, \qquad (9)$$

with similar values for $\sin\frac{b}{2}$ and $\sin\frac{c}{2}$.

Again,

$$2\cos^2\frac{a}{2} = 1 + \cos a = 1 + \frac{\cos A + \cos B \cos C}{\sin B \sin C}$$

$$= \frac{\cos A + \cos(B - C)}{\sin B \sin C} = \frac{2\cos(S - B)\cos(S - C)}{\sin B \sin C}.$$

therefore,

$$\cos\frac{a}{2} = \sqrt{\frac{\cos(S - B)\cos(S - C)}{\sin B \sin C}}, \qquad (10)$$

with similar values for $\cos\frac{b}{2}$ and $\cos\frac{c}{2}$.

Also

$$\tan\frac{a}{2} = \frac{\sin\frac{a}{2}}{\cos\frac{a}{2}} = \sqrt{\frac{-\cos S\cos(S - A)}{\cos(S - B)\cos(S - C)}}, \qquad (11)$$

with similar values for $\tan\frac{b}{2}$ and $\tan\frac{c}{2}$.

Moreover,

$$\sin a = 2\sin\frac{a}{2}\cos\frac{a}{2}$$

$$= \frac{2}{\sin B \sin C}\sqrt{-\cos S\cos(S - A)\cos(S - B)\cos(S - C)}. \qquad (12)$$

We shall use the symbol N to denote the radical in this expression, so that we have

$$\sin B \sin C \sin a = \sin C \sin A \sin b = \sin A \sin B \sin c = 2N.$$

Remark.—The positive sign has been given to the radicals in the formulæ for $\sin\frac{a}{2}$, $\cos\frac{a}{2}$, and $\tan\frac{a}{2}$, since $\frac{a}{2}$ is necessarily less than $\frac{\pi}{2}$. See also Chap. II., Art. 23.

Examples.

1. $4N^2 = 1 - \cos^2 A - \cos^2 B - \cos^2 C - 2\cos A \cos B \cos C.$

2. If p, q, r be the perpendiculars from the vertices on the opposite sides, show that—

(α). $\sin a \sin p = \sin b \sin q = \sin c \sin r = 2n.$

(β). $\sin A \sin p = \sin B \sin q = \sin C \sin r = 2N;$

and hence

$$\frac{\sin a}{\sin A} = \text{etc.} = \frac{n}{N}.$$

3. Prove that

$$n = \frac{2N^2}{\sin A \sin B \sin C}.$$

4. Prove that

$$N = \frac{2n^2}{\sin a \sin b \sin c}.$$

5. Prove that

$$2N = (\sin a \sin b \sin c \sin^2 A \sin^2 B \sin^2 C)^{\frac{1}{2}}.$$

6. Prove that

$$\tan \frac{b}{2} \tan \frac{c}{2} = \frac{-\cos S}{\cos(S-A)}.$$

Geometrically thus (see Chap. ii, fig. 10),

$$\frac{-\cos S}{\cos(S-A)} = \frac{\tan AL}{\tan AD} : \frac{\tan AL}{\tan AY} = \cot AD \tan AY = \tan \frac{b}{2} \tan \frac{c}{2}.$$

7. Prove that

$$\frac{\sin^2 A + \sin^2 B + \sin^2 C}{\sin^2 a + \sin^2 b + \sin^2 c} = \frac{1 + \cos A \cos B \cos C}{1 - \cos a \cos b \cos c}.$$

(*Dublin Univ. Exam. Papers.*)

We have

$$\frac{N^2}{n^2} = \frac{\sin^2 A}{\sin^2 a} = \frac{\sin^2 A + \sin^2 B + \sin^2 C}{\sin^2 a + \sin^2 b + \sin^2 a} = \frac{N^2 - \sin^2 A - \sin^2 B - \sin^2 C}{n^2 - \sin^2 a - \sin^2 b - \sin^2 c};$$

therefore, &c. (see Ex. 1 and 2).

8. Given the vertical angle C of a triangle, and the function N, find the *envelope** of the base.

By formula (12)

$$2N = \sin B \sin C \sin a = \sin C \sin r,$$

where r is the perpendicular from the vertex on the base. The envelope is, therefore, a small circle round C as pole.

9. If $A = a$, then B and C are respectively equal or supplemental to b and c, and $N = n$.

10. Show that

$$4Nn = \sin a \sin b \sin c \sin A \sin B \sin C.$$

11. Verify the following formulæ by the expansions of $\tan \frac{a \pm b}{2}$:—

$$\alpha^\circ. \quad \tan \frac{a+b}{2} = \frac{\cos \frac{A-B}{2}}{\cos \frac{A+B}{2}} \tan \frac{c}{2}.$$

$$\beta^\circ. \quad \tan \frac{a-b}{2} = \frac{\sin \frac{A-B}{2}}{\sin \frac{A+B}{2}} \tan \frac{c}{2}.$$

12. Write out and otherwise establish the corresponding formulæ (to α° and β°, Ex. 11) for the supplemental triangle.

13. If l be the length of the arc joining the middle point of the base to the vertex, find an expression for its length in terms of the sides.

Let the arc l make an angle θ with the base c. Then, Art. 26,

$$\cos\theta = \frac{\cos a - \cos l \cos \frac{c}{2}}{\sin l \sin \frac{c}{2}} = -\frac{\cos b - \cos l \cos \frac{c}{2}}{\sin l \sin \frac{c}{2}};$$

therefore,

$$\cos l = \frac{\cos a + \cos b}{2 \cos \frac{c}{2}}.$$

* The envelope is the figure which it always touches.

14. What is the corresponding theorem in plano to Ex. 13?

Let α, β, γ, and λ be the *lengths* of the arcs a, b, c, and l respectively. Expanding by Chap. II., Art. 18, and neglecting the powers of the radius higher than the second, we have

$$1 - \frac{\lambda^2}{2r^2} = \frac{\left(1 - \frac{\alpha^2}{2r^2}\right)\left(1 - \frac{\beta^2}{2r^2}\right)}{2\left(1 - \frac{\gamma^2}{8r^2}\right)}.$$

Hence, by reduction,

$$\alpha^2 + \beta^2 = 2\left(\frac{\gamma}{2}\right)^2 + 2\lambda^2$$

a result well known in Geometry.

15. Having given the base and the sum of the cosines of the sides, find the locus of the vertex. (Cf. Ex. 13.)

Section III.

Three Important Theorems.

37. **Theorem I.**—In any spherical triangle

$$\cot a \sin b = \cot A \sin C + \cos b \cos C.$$




Fig. 18.

Let ABC (fig. 18) be a spherical triangle. Produce AC through C to O, so that $AO = \frac{\pi}{2}$. Join BO.

Applying Art. 26 (1), to the triangles BOC and BOA, we have

$$\cos BO = \cos a \sin b - \sin a \cos b \cos C \ . \ . \ .$$

from triangle BOC; and

$$\cos BO = \sin c \cos A \ . \ . \ .$$

from triangle BOA, since $\cos AO = 0$.

Hence, equating these expressions for $\cos BO$,

$$\cos a \sin b = \sin a \cos b \cos C + \sin c \cos A.$$

Dividing each side of this equation by $\sin a$, and substituting $\frac{\sin C}{\sin A}$ for $\frac{\sin c}{\sin a}$, and we have

$$\cot a \sin b = \cot A \sin C + \cos b \cos C \ . \ . \ . \qquad (1)$$

Similarly, by producing the side BC through C to a point $\frac{\pi}{2}$ from B, we have

$$\cot b \sin a = \cot B \sin C + \cos a \cos C \ . \ . \ . \qquad (2)$$


Similarly, by producing the sides of the triangle through the angles A and B, it follows that

$$\cot b \sin c = \cot B \sin A + \cos c \cos A. \qquad (3)$$
$$\cot c \sin b = \cot C \sin A + \cos b \cos A. \qquad (4)$$
$$\cot c \sin a = \cot C \sin B + \cos a \cos B. \qquad (5)$$
$$\cot a \sin c = \cot A \sin B + \cos c \cos B. \qquad (6)$$

The same results will be obtained directly from the equations

$$\cos a = \cos b \cos c + \sin b \sin c \cos A$$
$$\cos c = \cos a \cos b + \sin a \sin b \cos C$$
$$. \quad . \quad . \quad . \quad . \quad . \quad . \quad .$$

by eliminating the sides a or c.



The above formulæ, as they enable us, when given two sides and an angle or two angles and a side of a triangle, to determine the remaining parts, are of great importance, and are thus expressed in general terms:—

Cotangent of any side × *sine of another side* = *cotangent of the angle opposite to the first side* × *sine of included angle* + *cosine of second side* × *cosine of included angle.*

38. Applying equation (1) of preceding Article to the polar triangle, we have

$$-\cot A \sin B = -\cot a \sin c + \cos B \cos c;$$

or, by transposing,

$$\cot a \sin c = \cot A \sin B + \cos c \cos B,$$

an equation identical with (6), Art. 37.

Moreover, it will be seen that each of the formulæ just proved contains four parts of a spherical triangle, which are connected thus:—

(α) Two sides and included angle with one of the base angles; or,

(β) Two angles and an adjacent side with one of the remaining sides.

39. **Corresponding Theorem in Plane.**—Using equation (1), and putting

$$\frac{r}{a} = \cot a, \quad \frac{\beta}{r} = \sin b, \quad \text{and } \cos b = 1,$$

we have

$$\frac{\beta}{a} = \cot A \sin C + \cos C = \frac{\sin(A + C)}{\sin A}.$$

Hence

$$\frac{a}{\beta} = \frac{\sin A}{\sin B}.$$

(*Science and Art Math. Honors Exam.*)

Examples.

1. Having given two sides of a spherical triangle, find the length of the bisector of the vertical angle intercepted between the vertex and the base.




Fig. 19.

Let ι denote the internal bisector of the angle C, and θ (fig. 19) the angle it makes with the base.

Then

$$\cot a \sin \iota = \cot \theta \sin \frac{C}{2} + \cos \iota \cos \frac{C}{2},$$

and

$$\cot b \sin \iota = -\cot \theta \sin \frac{C}{2} + \cos \iota \cos \frac{C}{2};$$

by addition,

$$(\cot a + \cot b) \sin \iota = 2 \cos \iota \cos \frac{C}{2};$$

or

$$\cot \iota = \frac{\cot a + \cot b}{2 \cos \frac{C}{2}}.$$

Similarly, if η denote the external bisector of C,

$$\cot \eta = \frac{\cot a - \cot b}{2 \sin \frac{C}{2}}.$$

2. What are the analogous formulae in plano?

Ans. $\dfrac{1}{\text{Internal bisector}} = \dfrac{\dfrac{1}{a}+\dfrac{1}{b}}{2\cos\dfrac{C}{2}}$, or Internal Bisector $= \dfrac{2ab}{a+b}\cos\dfrac{C}{2}$.

and External Bisector $= \dfrac{2ab}{a\sim b}\sin\dfrac{C}{2}$.

3. If ι_1, ι_2, and ι_3, denote the bisectors of the internal angles of a triangle, prove that

$$\cot\iota_1\cos\frac{A}{2}+\cot\iota_2\cos\frac{B}{2}+\cot\iota_3\cos\frac{C}{2}=\cot a+\cot b+\cot c;$$

and hence, in a plane triangle (*vide* Hudson's *Plane Trig.*, p. 154),

$$\frac{\cos\frac{A}{2}}{\iota_1}+\frac{\cos\frac{B}{2}}{\iota_2}+\frac{\cos\frac{C}{2}}{\iota_3}=\frac{1}{a}+\frac{1}{b}+\frac{1}{c};$$

and if η_1, η_2, and η_3, denote the bisectors of the external angles,

$$\cot\eta_1\sin\frac{A}{2}+\cot\eta_2\sin\frac{B}{2}+\cot\eta_3\sin\frac{C}{2}\equiv 0.$$

4. Using the preceding notation, prove that

$$\cot\eta_1\sin\frac{A}{2}=\cot\iota_2\cos\frac{B}{2}\sim\cot\iota_3\cos\frac{C}{2}.$$

5. Find the angles θ and θ' (fig. 19) made by the bisectors of the angle C with the opposite side c.

From the equations of Example 1 we have at once

$$\cot\theta=\frac{(\cot a-\cot b)}{2\sin\dfrac{C}{2}}\sin\iota,$$

and

$$\cot\theta'=\frac{(\cot a+\cot b)}{2\cos\dfrac{C}{2}}\sin\eta.$$

6. Find the arc intercepted on the base by the bisectors of the vertical angle

$$\cos XY = \cot\theta \cot\theta' \quad \text{(fig. 19)}.$$

$$= \frac{(\cot^2 a - \cot^2 b)}{2 \sin C} \sin\iota \sin\eta\,; \text{ (Ex. 5)}$$

but $\sin\iota \sin\eta = \sin XY \sin r$ (Art. 31, Ex. 2), where r is the perpendicular from the vertex C on the base AB.

Therefore

$$\cot XY = \frac{(\cot^2 a - \cot^2 b) \sin a \sin b}{2 \sin c} \quad \text{(Art. 31, Exs. 1 and 2)}.$$

$$= \frac{\sin(a+b) \sin(a-b)}{2 \sin a \sin b \sin c};$$

or, in terms of the angles,

$$\cot XY = \frac{\sin^2 A - \sin^2 B}{2 \sin A \sin B \sin c}.$$

7. Show that the sum of the cotangents of the intercepts made by the internal and external bisectors of the angles of a spherical triangle on the opposite sides is equal to zero. (*Educational Times.*)

By the preceding example

$$\Sigma \cot XY = \Sigma \frac{\sin^2 a - \sin^2 b}{2 \sin a \sin b \sin c} \equiv 0.$$

8. Give the analogous property in plano.

(See Casey's *Sequel to Euclid*, sec. viii., Ex. 17.)

9. If ι_1, ι_2, and ι_3, the internal bisectors of the angles of a triangle, make angles respectively θ_1, θ_2, and θ_3, with the opposite sides, prove that

$$\frac{\sin\frac{A}{2}}{\sin\iota_1 \tan\theta_1} + \frac{\sin\frac{B}{2}}{\sin\iota_2 \tan\theta_2} + \frac{\sin\frac{C}{2}}{\sin\iota_3 \tan\theta_3} \equiv 0;$$

(See Ex. 5)

and that if θ_1', θ_2', and θ_3', are the angles made by the external bisectors with the opposite sides,

$$\frac{\cos\frac{A}{2}}{\sin\eta_1 \tan\theta_1'} + \frac{\cos\frac{B}{2}}{\sin\eta_2 \tan\theta_2'} + \frac{\cos\frac{C}{2}}{\sin\eta_3 \tan\theta_3'} = \cot a + \cot b + \cot c.$$

(See Ex. 5.)

10. From the vertex C of a spherical triangle an arc of a great circle is drawn to meet the base AB in D; if C_1 and C_2 are the segments of the angle C, respectively, opposite AD and BD, show that

$$\cot CD \sin C = \cot a \sin C_1 + \cot b \sin C_2 *.$$

(*Science and Art Math. Honors.*)

Let CD make an angle θ with the base, and we have

$$\cot a \sin CD = \cot \theta \sin C_2 + \cos CD \cos C_2 \ . \ . \ . ; \qquad (1)$$

also

$$\cot b \sin CD = -\cot \theta \sin C_1 + \cos CD \cos C_1 \ . \ . \ . \qquad (2)$$

Eliminate θ from equations (1) and (2); therefore, &c.

11. If ϕ be the angle between the base c of a spherical triangle and its bisector,

$$\cot A - \cot B = 2 \cos \frac{c}{2} \cot \phi.$$

12. Prove that

$$\frac{\cos^2 b - \cos^2 c}{\cos b \cot B - \cos c \cot C} = \frac{\cos^2 c - \cos^2 a}{\cos c \cot C - \cos a \cot A}$$

$$= \frac{\cos^2 a - \cos^2 b}{\cos a \cot A - \cos b \cot B},$$

and express each of the three quantities as a symmetric function of the sides of the triangle.

Multiplying equation (1) by $\cos a$ and (2) by $\cos b$, and subtracting the results; each of the expressions becomes $\sin b \sin c \sin A$, &c.

$$= 2n.$$

13. Write out the corresponding theorem for the supplemental triangle.

40. **Theorem II.**—*Relation connecting the angular distances of three points, A, B, and C, on the same great circle, with any other point, P, on the sphere.*

$$\text{Sin } BC \cos AP + \sin CA \cos BP + \sin AB \cos CP = 0.$$

* This relation connects the latitudes and *difference* of longitudes of three places in the same great circle on the surface of the Earth.

We have (fig. 20),

$$\cos AP = \cos AB \cos BP + \sin AB \sin BP \cos ABP \quad (1)$$

and

$$\cos CP = \cos BC \cos BP - \sin BC \sin BP \cos ABP. \quad (2)$$

Multiply—(1) by sin BC, and (2) by sin AB, and add the results. Then

$$\sin BC \cos AP + \sin AB \cos CP = \sin AC \cos BP;$$

or writing $\sin AC = -\sin CA$, we have the algebraic sum

$$\sin BC \cos AP + \sin CA \cos BP + \sin AB \cos CP$$

equal to zero.




Fig. 20.

41. **Analogous Theorem in Plane.**—By the method similar to that used in Art. 39, we get

$$BC.AP^2 + CA.BP^2 + AB.CP^2 = -BC.CA.AB.$$

(Townsend's *Modern Geometry*, vol. i.)

42. When the point B (fig. 20) bisects the base AC of the triangle ACP, the formula of preceding Article readily reduces to

$$\cos AP + \cos CP = 2 \cos \frac{AC}{2} \cos BP.$$

Or,

The sum of the cosines of the sides of a triangle is equal to twice the cosine of half the base into cosine of the bisector of the base. See also Ex. 13, Art. 36.

Again, when the point B on the base AC is ninety degrees distant from the middle point of AC, we have

$$\cos AP - \cos CP = 2 \sin \frac{AC}{2} \cos BP,$$

an equation which furnishes an obvious solution to the problem—*Given the base and difference of cosines of the sides, find the locus of the vertex.*

Examples.

1. Having given the base c of a triangle, and $l \cos a \pm m \cos b$, find the locus of the vertex, l and m being constants.

Ans. A circle, the pole of which divides the base internally (or externally) into segments, the sines of which are to one another as l is to m.

NOTE.—Particular cases of this example, viz., when $l = \pm m$, have been considered in Arts. 42 and 36.

2. Having given the base and ratio of cosines of the sides; find the locus of the vertex.

Let the arc CP (fig. 20) be a quadrant. Then in the triangle ABP, the formula of Art. 40 reduces thus:

$$\frac{\cos AP}{\cos BP} = \frac{\sin AC}{\sin BC};$$

hence, according to the given conditions, C is a fixed point on the base, and the vertex therefore describes a great circle having C for pole.

3. Show that the points A, B, C (fig. 20) will lie on the same great circle if

$$\sin BC \sin AL + \sin CA \sin BL + \sin AB \sin CL = 0,$$

where AL, BL, CL, are secondaries to any great circle L.

What is the corresponding theorem in plano?

(*Vide* Townsend's *Modern Geometry*, vol. i., chap. 5.)

4. If the base c of a triangle be a quadrant, show that if any point X be taken on it,

$$\cos CX = \cos a \sin AX + \cos b \cos AX.$$

5. From any three points on a great circle, secondaries x, y, z; x', y', z'; x'', y'', z'' be drawn to the sides of a triangle, prove the determinant relation,

$$\begin{vmatrix} \sin x & \sin y & \sin z \\ \sin x' & \sin y' & \sin z' \\ \sin x'' & \sin y'' & \sin z'' \end{vmatrix} = 0.$$

(Apply Ex. 3.)

43. **Theorem III.**—*If from a fixed point P on the surface of a sphere an arc of a great circle be drawn, meeting a small circle at A and B, and if PA and PB be denoted by ρ and ρ', then*

$$\tan \tfrac{1}{2}\rho \tan \tfrac{1}{2}\rho' = \text{constant}.$$

Let C (fig. 21) be the pole of the small circle. Join




Fig. 21.

CA, CB, and CP by arcs of great circles. Let the angle APC be denoted by θ, the arc CP by δ, and the arc CA by a.

From the triangle APC we have

$$\cos a = \cos\rho \cos\delta + \sin\rho \sin\delta \cos\theta.^*$$

Let $\tan\frac{1}{2}\rho = t$. Then $\cos\rho = \frac{1-t^2}{1+t^2}$, and $\sin\rho = \frac{2t}{1+t^2}$.

Substituting these values for $\cos\rho$ and $\sin\rho$ in the above equation, and arranging in powers of t, we get

$$t^2(\cos a + \cos\delta) - 2t \sin\delta \cos\theta + \cos a - \cos\delta = 0.$$

By proceeding in identically the same manner we obtain from the triangle BPC an equation in t' (where $t' = \tan\frac{1}{2}\rho'$) in all respects the same as that in t, and hence it follows that $\tan\frac{1}{2}\rho$ and $\tan\frac{1}{2}\rho'$ are the roots of the above quadratic, and therefore their product

$$\tan\tfrac{1}{2}\rho \tan\tfrac{1}{2}\rho' = \frac{\cos a - \cos\delta}{\cos a + \cos\delta} = \tan\tfrac{1}{2}(\delta - a)\tan\tfrac{1}{2}(\delta + a),$$

which value is independent of θ, and depends only on the fixed quantities a and δ; therefore, &c.

Otherwise thus:—

Draw CM perpendicular to AB. Then $AM = BM$; and since (Art. 27, Ex. 1) $\cos\delta = \cos PM \cos CM$, and $\cos a = \cos AM \cos CM$, we have by division

$$\frac{\cos AM}{\cos PM} = \frac{\cos a}{\cos\delta}.$$

Therefore,

$$\frac{\cos AM - \cos PM}{\cos AM + \cos PM} = \frac{\cos a - \cos\delta}{\cos a + \cos\delta}.$$

* This may be regarded as the polar equation of the small circle, and its analogue in plano, $\rho^2 - 2\rho d\cos\theta + d^2 - a^2 = 0$, can be deduced at once by aid of the formulæ of Art. 18.

which reduces at once to

$$\tan \tfrac{1}{2}\rho \tan \tfrac{1}{2}\rho' = \tan \tfrac{1}{2}(\delta - a) \tan \tfrac{1}{2}(\delta + a).$$

Remark.—In fig. 21 it is to be observed that we have *two* triangles CPA and CPB, having the same *three* parts a, δ, and θ. Whenever more than one triangle can be constructed from three given parts the triangle is said to be ambiguous (see Chap. v.) However, it should be noted that the remaining sides ρ and ρ' of the triangles are not independent of each other, but are connected by the relation of this Article.

Examples.

1. Deduce the *corresponding theorem in plano.*

Ans.—Euclid I. XXXVI.

2. If the vertical angle C of a triangle be equal to the sum of the base angles, and if the perpendicular p from the vertex on the base divides it into segments c_1 and c_2, show that

$$\tan \tfrac{1}{2}c_1 \tan \tfrac{1}{2}c_2 = \tan^2 \tfrac{1}{2}p.$$

Let p meet the base in D. Produce CD through D to C', so that $DC' = DC$. Then the triangles ABC and ABC' are equal in all respects, and the vertices of the quadrilateral $ACBC'$ are concyclic, being equidistant from the middle point of AB. Therefore, by Theorem III.,

$$\tan \tfrac{1}{2}c_1 \tan \tfrac{1}{2}c_2 = \tan \tfrac{1}{2}CD \tan \tfrac{1}{2}C'D = \tan^2 \tfrac{1}{2}p.$$

3. Given the base AB, and that $C = A + B$, find the locus of the vertex C.

Ans.—A circle having its pole at the middle point of the base.

4. In Theorem III., fig. 21, show that $\tan \frac{1}{2}\rho + \tan \frac{1}{2}\rho'$ is a maximum, when PAB passes through C, and determine the maximum value. (Use the quadratic in t.)

$$Ans.—\frac{2 \sin \delta}{\cos a + \cos \delta}.$$

5. Show that $\sin \frac{1}{2}\rho \sin \frac{1}{2}\rho' \sin \frac{1}{2}(\rho + \rho')$ varies as $\cos \theta$, and explain when $\theta = 90°$.

TABLE OF FORMULAE.

FORMULAE.	SUPPLEMENTAL FORMULAE.	ANALOGOUS FORMULAE IN PLANO.
$\cos a = \cos b \cos c + \sin b \sin c \cos A$ (Art. 26.)	$\cos A + \cos B \cos C = \sin B \sin C \cos a$ (Art. 34.)	$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ (Art. 28.)
$\cos b = \cos c \cos a + \sin c \sin a \cos B$,,	$\cos B + \cos C \cos A = \sin C \sin A \cos b$,,	and
$\cos c = \cos a \cos b + \sin a \sin b \cos C$,,	$\cos C + \cos A \cos B = \sin A \sin B \cos c$,,	$A + B + C = \pi$ (Art. 35.)
$\sin \frac{1}{2}A = \sqrt{\frac{\sin(s-b)\sin(s-c)}{\sin b \sin c}}$ (Art. 33.)	$\sin \frac{1}{2}a = \sqrt{-\frac{\cos S \cos(S-A)}{\sin B \sin C}}$ (Art. 36.)	$\sin \frac{1}{2}A = \sqrt{\frac{(s-b)(s-c)}{bc}}$.
$\cos \frac{1}{2}A = \sqrt{\frac{\sin s \sin(s-a)}{\sin b \sin c}}$,,	$\cos \frac{1}{2}a = \sqrt{\frac{\cos(S-B)\cos(S-C)}{\sin B \sin C}}$,,	$\cos \frac{1}{2}A = \sqrt{\frac{s(s-a)}{bc}}$.
$\tan \frac{1}{2}A = \sqrt{\frac{\sin(s-b)\sin(s-c)}{\sin s \sin(s-a)}}$,,	$\tan \frac{1}{2}a = \sqrt{-\frac{\cos S \cos(S-A)}{\cos(S-B)\cos(S-C)}}$,,	$\tan \frac{1}{2}A = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$.
With similar expressions for $\sin \frac{1}{2}B$, etc., and $\sin \frac{1}{2}C$, etc.		and $A + B + C = \pi$.

$\sin A = \dfrac{2n}{\sin b \sin c}$ (Art. 30.)	$\sin a = \dfrac{2N}{\sin B \sin C}$ (Art. 36.)	$\sin A = \dfrac{2\Delta}{bc}$
$2n = 2\sqrt{\sin s \sin(s-a) \sin(s-b) \sin(s-c)}$ $= \sqrt{1-\cos^2 a - \cos^2 b - \cos^2 c + 2\cos a \cos b \cos c}$	$2N = 2\sqrt{-\cos S \cos(S-A) \cos(S-B) \cos(S-C)}$ $= \sqrt{1-\cos^2 A - \cos^2 B - \cos^2 C - 2\cos A \cos B \cos C}$	$2\Delta = 2\sqrt{s(s-a)(s-b)(s-c)}$. $A + B + C = \pi$.
$\dfrac{\sin A}{\sin a} = \dfrac{\sin B}{\sin b} = \dfrac{\sin C}{\sin c} = \dfrac{2n}{\sin a \sin b \sin c}$ (Art. 31.)	$\dfrac{\sin a}{\sin A} = \dfrac{\sin b}{\sin B} = \dfrac{\sin c}{\sin C} = \dfrac{2N}{\sin A \sin B \sin C}$ (Art. 31.)	$\dfrac{\sin A}{a} = \dfrac{\sin B}{b} = \dfrac{\sin C}{c} = \dfrac{2\Delta}{abc} = \dfrac{1}{D}$, Δ being the area of the triangle, and D the diameter of its circumscribed circle.
$\cot a \sin b = \cot A \sin C + \cos b \cos C$ (Art. 37.) $\cot b \sin c = \cot B \sin A + \cos c \cos A$,, $\cot c \sin a = \cot C \sin B + \cos a \cos B$,,	$\cot a \sin c = \cot A \sin B + \cos c \cos B$ (Art. 38.) $\cot b \sin a = \cot B \sin C + \cos a \cos C$,, $\cot c \sin b = \cot C \sin A + \cos b \cos A$,,	$\dfrac{\sin A}{a} = \dfrac{\sin B}{b} = \text{etc.}$ (Art. 39.)

Miscellaneous Examples.

1. If s and s' are the segments of the base made by the perpendicular from the vertex, and σ and σ' those made by the bisector of the vertical angle, show that

$$\tan\frac{s-s'}{2}\tan\frac{\sigma-\sigma'}{2}=\tan^2\frac{a-b}{2}.$$

(*Dub. Univ. Exam. Papers.*)

We have

$$\frac{\cos a}{\cos b}=\frac{\cos s}{\cos s'} \quad \text{and} \quad \frac{\sin a}{\sin b}=\frac{\sin \sigma}{\sin \sigma'}.$$

Taking the difference to the sum in each equation, we get

$$\tan\frac{a-b}{2}\tan\frac{a+b}{2}=\tan\frac{s-s'}{2}\tan\frac{c}{2}, \tag{1}$$

and

$$\tan\frac{a-b}{2}\cot\frac{a+b}{2}=\tan\frac{\sigma-\sigma'}{2}\cot\frac{c}{2}. \tag{2}$$

Multiplying (1) and (2) together, we have

$$\tan\frac{s-s'}{2}\tan\frac{\sigma-\sigma'}{2}=\tan^2\frac{a-b}{2}.$$

2. Prove that

$$\sin b\sin c+\cos b\cos c\cos A=\sin B\sin C-\cos B\cos C\cos a.$$

[If from the extremities of the base AB quadrants AA' and BB' be measured on the sides towards the vertex, the left-hand side of this equation is obviously $\cos A'B'$.

Again, if secondaries be drawn to the sides at A and B, the angle between them is equal to the arc $A'B'$ (Chap. I., Art. 6), and the cosine of this angle is equal to the expression on the right-hand side of the equation by Chap. III., Art. 34; therefore, &c.]

Otherwise thus:—

Multiplying together the equations

$$\cos a=\cos b\cos c+\sin b\sin c\cos A,$$

and

$$\sin B\sin C\cos a-\cos B\cos C=\cos A,$$

and observing that

$$\sin B\sin C\sin^2 a=\sin b\sin c\sin^2 A, \qquad \text{(Art. 31.)}$$

the above result is obtained.

3. Given a, b and c, the sides of a triangle, find γ, the arc joining the middle points of a and b.

We have

$$\cos\gamma = \cos\frac{a}{2}\cos\frac{b}{2} + \sin\frac{a}{2}\sin\frac{b}{2}\cos C$$

$$= \cos\frac{a}{2}\cos\frac{b}{2} + \frac{\cos c - \cos a\cos b}{4\cos\frac{a}{2}\cos\frac{b}{2}}$$

(by Art. 26.)

$$= \frac{(1+\cos a)(1+\cos b) + \cos c - \cos a\cos b}{4\cos\frac{a}{2}\cos\frac{b}{2}}$$

$$= \frac{1+\cos a+\cos b+\cos c}{4\cos\frac{a}{2}\cos\frac{b}{2}},$$

with similar expressions for the arcs joining the middle points of b and c, c and a.

Note.—In the colunar triangle

$$\cos\gamma' = \frac{1+\cos c-\cos a-\cos b}{4\sin\frac{a}{2}\sin\frac{b}{2}}.$$

4. If α, β and γ be the arcs joining the middle points of the sides, prove

$$\frac{\cos\alpha}{\cos\frac{a}{2}} = \frac{\cos\beta}{\cos\frac{b}{2}} = \frac{\cos\gamma}{\cos\frac{c}{2}} = \frac{1+\cos a+\cos b+\cos c}{4\cos\frac{a}{2}\cos\frac{b}{2}\cos\frac{c}{2}} = \sin S;$$

and hence, if one of the arcs be a quadrant, the triangle whose sides are α, β, and γ is quadrantal. (See Ex. 3.)

5. If the side c of a spherical triangle be a quadrant, and δ the arc drawn at right angles to it from the opposite vertex, show that

$$\cot^2\delta = \cot^2 A + \cot^2 B.$$

(*Science and Art Exam. Papers.*)

[Let x be the foot of the perpendicular δ on the base AB; therefore

$$\tan\delta = \tan A \,.\, \sin AX = \tan B \,.\, \cos AX;$$

therefore, &c.]

6. Given the vertical angle in position and magnitude, and the ratio of the cosines of the base angles, find the locus of the pole of the base.

[Let the perpendicular on the base divide the vertical angle into parts α and β; then by Art. 26, Ex. 6,

$$\frac{\cos A}{\cos B} = \frac{\sin \alpha}{\sin \beta};$$

therefore the perpendicular is fixed in direction, and it obviously contains the pole of the base; therefore, &c.]

7. Prove that in fig. 18

$$2 \cot XY = \cot AY + \cot BY.*$$

For

$$\frac{\sin AY}{\sin BY} = \frac{\sin AX}{\sin XB} = \frac{\sin (AY - XY)}{\sin (XY - BY)};$$

therefore, &c.

8. If a, b, c, d be the sides of a spherical quadrilateral taken in order, and δ and δ' the diagonals; the arc ϕ joining the middle points of the diagonals is given by the equation

$$\cos \phi = \frac{\cos a + \cos b + \cos c + \cos d}{4 \cos \frac{\delta}{2} \cos \frac{\delta}{2}},$$

and give the corresponding theorem in plano. (See Art. 36, Ex. 14.)

[The sum of the squares of the sides of a quadrilateral = the sum of the squares of its diagonals + four times the square of the line joining the middle points of the diagonals. (See M'Dowell's *Geom. Exercises.*)].

9. The arc ϕ' joining the middle points of the opposite sides c and d of a quadrilateral is given by the equation

$$\cos \phi' = \frac{\cos a + \cos b + \cos \delta + \cos \delta'}{4 \cos \frac{c}{2} \cos \frac{d}{2}},$$

* An arc AB divided internally in X and externally in Y, such that sin AY : sin BY : : sin AX : sin XB, is said to be cut harmonically by the segment XY; and the cotangents of either triad of arcs, viz. AY, XY, BY, or AX, AB, AY, measured from either extremity are therefore in arithmetical progression.

with a similar expression for the arc joining the middle points of a and b (See Ex. 8.)

10. If the arcs joining the middle points of opposite sides of a quadrilateral be each quadrants, the arc ϕ joining the middle points of the diagonals δ and δ' is given by the equation

$$-\cos\phi = \frac{\cos\delta + \cos\delta'}{2\cos\frac{\delta}{2}\cos\frac{\delta'}{2}}.$$


11. Prove that the angle ϕ between the perpendicular from the vertex on the base and the bisector of the vertical angle is thus given :

$$\tan\phi = \frac{\cos\frac{1}{2}(a-b)}{\cos\frac{1}{2}(a+b)} \cdot \tan\frac{A-B}{2}.^*$$

(*Dub. Univ. Exam. Papers.*)

For

$$\frac{\cos A}{\cos B} = \frac{\sin\left(\frac{C}{2}+\phi\right)}{\sin\left(\frac{C}{2}-\phi\right)};$$



therefore, taking the difference to sum,

$$\tan\frac{A+B}{2}\tan\frac{A-B}{2} = \cot\frac{C}{2}\tan\phi.$$

Now substitute the value of $\tan\frac{A+B}{2}$ in Art. 33, Ex. 9; therefore, &c.

12. Given the base of a triangle and the bisectors of the vertical angle, construct it.

Using the notation, fig. 18,

$$\cos XY = \cos i \cos\eta;$$

also

$$\cot AX + \cot AY = 2\cot AB, \qquad \text{(Ex. 7)}$$

two equations which determine the points X and Y.

* Tan ϕ is also equal to

$$\frac{\sin\frac{1}{2}(a-b)}{\sin\frac{1}{2}(a+b)}\tan\tfrac{1}{2}(A+B).$$

13. Given the base AB and the sum of the sides of a triangle, find the locus of a point O, where the external bisector of the vertical angle C meets the secondary to BC at the point B. (Cf. Galbraith and Haughton's *Euclid*, I. II. III., Ex. 200,)




Fig. 22.

Produce AC (fig. 22) through C to D, such that $CD = BC$. Therefore AD = given sum of the sides.

Since CO bisects the angle BCD, it bisects BD at right angles.

Since BD is bisected at right angles, $BO = OD$, and the triangles CDO and CBO are equal in every respect; hence the angle CDO is a right angle. Therefore

$$\cos AD = \frac{\cos AO}{\cos OD} = \frac{\cos AO}{\cos BO}.$$

In the triangle ABO we have thus given the base AB and ratio of the cosines of the sides, to find the locus of the vertex, for a solution of which, see Art. 42, Ex. 2.

14. Given the base AB and difference of sides, find the locus of the point O, where the secondary to BC at the point B meets the internal bisector of the vertical angle. (See Ex. 13.)

15. Two places on the Earth's surface are distant, one θ' from the Pole, the other θ' from the Equator, and their difference of longitude is ϕ; find the angular distance between them. (*Science and Art Exam. Papers.*)

Ans $\cos \delta = \sin 2\theta \cos^2 \frac{\phi}{2}$.

16. In an isosceles triangle, if each of the base angles be double the vertical angle, prove that

$$\cos a \cos \frac{a}{2} = \cos\left(c + \frac{a}{2}\right).$$

If x denote one of the equal bisectors of the sides, we have

$$\cos x = \cos a \cos \frac{a}{2} + \sin a \sin \frac{a}{2} \cos A$$

$$= \cos a \cos \frac{a}{2} + \sin a \sin \frac{a}{2} \left(2 \cos^2 \frac{A}{2} - 1\right),$$

where A is the vertical angle.

Again,

$$\frac{\sin a}{\sin c} = 2 \cos A,$$

eliminating $\cos A$; therefore, &c.

17. If a side c of a spherical triangle be a quadrant, show that

(α). $\operatorname{Cot} a \cot b + \cos C = 0.$

(β). $\operatorname{Cos} S \cos(S - C) + \cos(S - A) \cos(S - B) = 0.$

(*Science and Art Exam. Papers.*)

(See Art. 26.) Also

$$\tan \frac{c}{2} = 1;$$

therefore, &c., Art. 36.

18. If λ, μ and ν be the segments towards the angles of the internal bisectors of the angles of a triangle ABC from the point of concurrence, show that

$$\begin{vmatrix} \cos\lambda & \cos\mu & \cos\nu \\ \sin a & \sin b & \sin c \\ \cos a & \cos b & \cos c \end{vmatrix} \equiv 0.$$

We have

(1). $\cos\lambda = \cos b \cos\nu + \sin b \sin\nu \cos \frac{C}{2}.$

(2). $\cos\mu = \cos a \cos\nu + \sin a \sin\nu \cos \frac{C}{2}.$

Eliminate $\cos \frac{C}{2}$. Therefore,

$$\cos\lambda \sin a - \cos\mu \sin b = \cos\nu \sin(a - b).$$

Similarly,

$$\cos\mu \sin b - \cos\nu \sin c = \cos\lambda \sin(b - c),$$

and

$$\cos\nu \sin c - \cos\lambda \sin a = \cos\mu \sin(c - a);$$

whence, on adding

$$\Sigma \cos\lambda \sin(b - c),$$

or its equivalent determinant form, given above, is equal to zero.

19. If λ, μ and ν be the segments towards the angles of two external bisectors (of angles B and C), and one internal bisector of the angles of a triangle from the point of concurrence, show that

$$\begin{vmatrix} -\cos\lambda & \cos\mu & \cos\nu \\ \sin a & \sin b & \sin c \\ -\cos a & \cos b & \cos c \end{vmatrix} \equiv 0.$$

20. Using the notation of Example 18, prove that

$$\Sigma^* \sin\lambda \sin(b - c) \cos\frac{A}{2} \equiv 0.$$

Multiplying Ex. 18, (1) by $\cos a$, and (2) by $\cos b$, and subtracting, we get

$$\cos a \cos\lambda - \cos b \cos\mu + \sin(a - b) \sin\nu \cos\frac{C}{2} = 0,$$

which, added to similar equations, involving b and c, and c and a, gives the required result.

21. Using the notation of Ex. 19, show that

$$\begin{vmatrix} \sin\lambda \cos\frac{A}{2} & \sin\mu \sin\frac{B}{2} & \sin\nu \sin\frac{C}{2} \\ \sin a & \sin b & \sin c \\ -\cos a & \cos b & \cos c \end{vmatrix} \equiv 0.$$

* The equivalent determinant form is

$$\begin{vmatrix} \sin\lambda \cos\frac{A}{2} & \sin\mu \cos\frac{B}{2} & \sin\nu \cos\frac{C}{2} \\ \sin a & \sin b & \sin c \\ \cos a & \cos b & \cos c \end{vmatrix} \equiv 0.$$

22. If θ, ϕ, and ψ denote the arcs connecting the middle points of pairs of opposite sides and diagonals of a quadrilateral, viz. a and b, c and d, and δ and δ' respectively, prove that

$$\cos\theta\cos\frac{a}{2}\cos\frac{b}{2}+\cos\phi\cos\frac{c}{2}\cos\frac{d}{2}-\cos\psi\cos\frac{\delta}{2}\cos\frac{\delta'}{2}=\tfrac{1}{2}(\cos\delta+\cos\delta').$$

(See Exs. 8 and 9.)

32. A, B, and C are three concyclic points on a sphere; X, Y and Z are three other points; show that the relation

$$\begin{vmatrix} \cos AX & \cos AY & \cos AZ \\ \cos BX & \cos BY & \cos BZ \\ \cos CX & \cos CY & \cos CZ \end{vmatrix} \equiv 0.$$

[By aid of Art. 40 we can eliminate sin BC, sin CA, and sin AB from three given equations; therefore, &c.]

24. If the sides of a quadrilateral taken in order are a, b, c, d, show that the diagonals δ and δ' intersect at an angle θ, given by the equation

$$\cos\theta=\frac{\cos a\cos c-\cos b\cos d}{\sin\delta\sin\delta'}.$$




Fig. 23.

For, representing the segments of the diagonals by x, $\delta-x$, and x', $\delta'-x'$ respectively, we have

$$\cos a=\cos x'\cos(\delta-x)+\sin x'\sin(\delta-x)\cos\theta \quad (1)$$
$$\cos c=\cos x\cos(\delta'-x')+\sin x\sin(\delta'-x')\cos\theta. \quad (2)$$
$$\cos b=\cos(\delta-x)\cos(\delta'-x')-\sin(\delta-x)\sin(\delta'-x')\cos\theta. \quad (3)$$
$$\cos d=\cos x\cos x'-\sin x\sin x'\cos\theta. \quad (4)$$

By subtracting the product of equations (3) and (4) from the product of (1) and (2), the above expression for θ is obtained.

25. By a method similar to that used in Ex. 24, show that if ϕ and ψ are the angles between a and c, b and d, respectively,

$$\cos\psi = \frac{\cos c \cos a - \cos\delta \cos\delta'}{\sin b \sin d}. \qquad (a)$$

$$\cos\phi = \frac{\cos b \cos d - \cos\delta \cos\delta'}{\sin c \sin a}. \qquad (b)$$

26. An arc δ drawn from the vertex of a triangle to meet the base at an angle θ may be expressed by the equation

$$\sin\delta = \frac{\cos a \cos y - \cos b \cos x}{\sin c \cos\theta},$$

where x and y are the segments of the base adjacent to the sides a and b respectively.

[In the figure of Ex. 24 suppose three vertices of the quadrilateral to be concyclic].

27. In any triangle prove that

$$\tan\frac{a}{2} : \tan\frac{b}{2} : \tan\frac{c}{2} = \cos(S-A) : \cos(S-B) : \cos(S-C).$$

(See p. 42, Ex. 3.)

28. Prove the following expression* for $\cos S$:—

$$-\cos S = \frac{\sin\frac{a}{2}\sin\frac{b}{2}\sin C}{\cos\frac{c}{2}}.$$

Also Apply Art. 36, (9) and (10).

$$\sin s = \frac{\cos\frac{A}{2}\cos\frac{B}{2}\sin c}{\sin\frac{C}{2}}.$$

* The equation for calculating the sum of the angles, having given two sides and the included angle, or by the aid of Art. 30, the three sides of a spherical triangle will be afterwards proved as *Cagnoli's Theorem*.

29. Prove that

$$\sin(S - A) = \frac{1 + \cos a - \cos b - \cos c}{4 \cos \frac{a}{2} \sin \frac{b}{2} \sin \frac{c}{2}},$$

and similar expressions for $\sin(S - B)$ and $\sin(S - C)$.

For,

$$\sin(S - A) = \sin \frac{B + C}{2} \cos \frac{A}{2} - \cos \frac{B + C}{2} \sin \frac{A}{2};$$

therefore, &c., by the aid of note, p. 43.

Or thus :—

Take the colunar triangle, whose angles are $\pi - B$, $\pi - C$, A, and sides $\pi - b$, $\pi - c$, a (Art. 16). Let

$$2S' = \pi - B + \pi - C + A = 2\pi - (B + C - A);$$

hence

$$S' = \pi - (S - A),$$

and the formula adapted to the colunar triangle becomes

$$\sin S' = \frac{1 + \cos a + \cos b + \cos c}{4 \cos \frac{a}{2} \cos \frac{b}{2} \cos \frac{c}{2}};$$

therefore, &c. (Cf. Ex. 4.)

30. Prove that

$$\cos(s - a) = \frac{1 + \cos B + \cos C - \cos A}{4 \cos \frac{B}{2} \cos \frac{C}{2} \sin \frac{A}{2}}.$$

31. Prove that

$$\cos(S - A) = \frac{\cos \frac{b}{2} \cos \frac{c}{2}}{\cos \frac{a}{2}} \sin A.$$

32. Prove that

$$\sin(S - A) = \frac{\sin \frac{b}{2} \sin \frac{c}{2} + \cos \frac{b}{2} \cos \frac{c}{2} \cos A}{\cos \frac{a}{2}}.$$

(See Ex. 29.)

[It is evident that the numerator of the fraction is equal to the cosine of the arc joining the middle points of the sides $\pi - b$, $\pi - c$, of the colunar triangle; therefore, &c. (Art. 26, and Ex. 4.)]

33. In any triangle

$$\frac{\cos a - \cos b}{1 - \cos c} + \frac{\sin (A - B)}{\sin C} \equiv 0.$$

(See p. 46, Ex. 3.)

34. Given the formula, Art. 36, for $\tan \frac{a}{2}$; prove that

$$-\tan S = \frac{\cot \frac{a}{2} \cot \frac{b}{2} + \cos C.}{\sin C}.$$

35. Show that

$$\cot \frac{a}{2} \cos (S - A) = \cot \frac{b}{2} \cos (S - B) = \cot \frac{c}{2} \cos (S - C)$$

$$= -\cot \frac{a}{2} \cot \frac{b}{2} \cot \frac{c}{2} \cos S;$$

and hence

$$\Sigma \cos^2 (S - A) = \cos^2 S \Sigma \cot^2 \frac{a}{2} \cot^2 \frac{b}{2}.$$

36. If

$$\tan \frac{a}{2} \tan \frac{b}{2} = \tan^2 \frac{c}{2},$$

show that

$$\cos (S - A) \cos (S - B) = \cos^2 (S - C).$$

37. If a spherical triangle is equal and similar to its polar triangle,

$$\sec a = \sec b \sec c + \tan b \tan c.$$

Give a particular example of such a triangle.

(*Science and Art Exam. Papers.*)

38. Prove that by a closer approximation, employing the notation used in Legendre's Theorem (Art. 29),

$$\cos A = \cos A' - \frac{\beta\gamma \sin^2 A'}{6r^2} + \frac{\beta\gamma (\alpha^2 - 3\beta^2 - 3\gamma^2) \sin^2 A'}{180r^4}.$$

39. Deduce the following formula for $a+b$ in terms of the opposite angles A and B, and the base c:—

$$\frac{a+b}{2} = \frac{c}{2} + \tan\frac{A}{2}\tan\frac{B}{2}\sin c + \frac{1}{2}\tan^2\frac{A}{2}\tan^2\frac{B}{2}\sin 2c$$

$$+ \frac{1}{3}\tan^3\frac{A}{2}\tan^3\frac{B}{2}\sin 3c, + \ldots$$

(*Bishop Law's Exam., Trin. Coll., Dub.*)

For
$$\tan\frac{a+b}{2} = \frac{\cos\dfrac{A-B}{2}}{\cos\dfrac{A+B}{2}}\tan\frac{c}{2},$$

(Art. 33, Ex. 9.)

being of the form $\tan\dfrac{a+b}{2} = n\tan\dfrac{c}{2}$, can be expanded in series for $\dfrac{a+b}{2}$.

(See Todhunter's *Plane Trig.*, p. 238, Art. 299.)

40. From any three points on a sphere are drawn to each of three points on a great circle arcs, a_1, b_1, c_1; a_2, b_2, c_2; a_3, b_3, c_3; making angles with the great circle, respectively, A_1, A_2, A_3, etc. . . . ; prove the determinant relation

$$\begin{vmatrix} \sin a_1\cos A_1 & \sin a_2\cos A_2 & \sin a_3\cos A_3 \\ \sin b_1\cos B_1 & \sin b_2\cos B_2 & \sin b_3\cos B_3 \\ \sin c_1\cos C_1 & \sin c_2\cos C_2 & \sin c_3\cos C_3 \end{vmatrix} \equiv 0.$$

41. If a and b are the sides, and c the base of a triangle, prove that if θ be the angle which β, the bisector of the base, makes with the base,

$$\tan^2\tfrac{1}{2}a\,\tan^2\tfrac{1}{2}b = \frac{\tan^4\frac{1}{4}c - 2\tan^2\frac{1}{4}c\tan^2\frac{1}{2}\beta\cos 2\theta + \tan^4\frac{1}{2}\beta}{1 - 2\tan^2\frac{1}{4}c\tan^2\frac{1}{2}\beta\cos 2\theta + \tan^4\frac{1}{4}c\tan^4\dfrac{\beta}{2}}.$$

(*Dublin Univ. Exam. Papers.*)

[Find $\tan^2\frac{1}{2}a$, and $\tan^2\frac{1}{2}b$, and multiply the results].

42. Find the locus of the vertex, having given the base and—

(1) Ratio of cotangents of the base angles.

(2) Sum or difference of cotangents.

(3) $l\cot A + m\cot B$. (Cf. p. 58, Ex. 10.)

(*Educational Times*, April, 1885.)

CHAPTER IV.

RIGHT-ANGLED TRIANGLES.

SECTION I.

44. **Geometrical Deduction of Formulæ connecting the Parts of a Right-angled Triangle.**— The formulæ necessary for the solution of right-angled triangles have been deduced in Chapter III. as particular cases of the general formula

$$\cos a = \cos b \cos c + \sin b \sin c \cos A.$$

We, however, add the following geometrical method of obtaining them, as it is both simple and instructive :—

Let ABC (fig. 24) be a triangle right-angled at C, and let O be the centre of the sphere on which it is described. Since C is a right angle, the planes OCA and OCB are perpendicular to each other (Art. 6). Take P any point on the radius OA, and draw PR perpendicular to OC, and RQ perpendicular to OB. Join PQ. Since the planes COA and COB are at right angles, the angle PRQ is right, and, as in Art. 32, the angle OQP is also right. Hence we have

$$\frac{OQ}{OP} = \frac{OQ}{OR} \cdot \frac{OR}{OP};$$

that is,

$$\cos c = \cos a \cos b. \qquad (1)$$

$$\frac{QR}{QO} = \frac{QR}{QP} \cdot \frac{QP}{QO};$$

that is, $\tan a = \cos B \tan c.$

Similarly, $\tan b = \cos A \tan c.$ (2)




Fig. 24

$$\frac{PR}{PO} = \frac{PR}{PQ} \cdot \frac{PQ}{PO};$$

that is, $\sin b = \sin B \sin c.$

Similarly, $\sin a = \sin A \sin c.$ (3)

$$\frac{RP}{RO} = \frac{RP}{RQ} \cdot \frac{RQ}{RO};$$

that is, $\tan b = \tan B \sin a.$

Similarly, $\tan a = \tan A \sin b.$ (4)

From R draw RP' perpendicular to OA, and draw $P'Q'$ also perpendicular to OA, in the plane AOB, and meeting OB in Q'. Then (Art. 6) the angle $Q'P'R$ is equal to the angle A. Join $Q'R$.

Now $PQ'^2 = P'Q'^2 + PP'^2 = PR^2 + RQ'^2$; (Art. 32.) therefore

$P'Q'^2 = PR^2 + RQ'^2 - PP'^2$. But $PR^2 - PP'^2 = RP^2$.

Hence $P'Q'^2 = RP'^2 + RQ'^2$; and therefore $P'RQ'$ is right, and we have

$$\cot A \cot B = \frac{RP'}{RQ'} \cdot \frac{RQ}{RP} = \frac{RP'}{RP} \cdot \frac{RQ}{RQ'}.$$

But $$\frac{RP'}{RP} = \sin RPP' = \cos ROP = \cos b,$$

and $$\frac{RQ}{RQ'} = \sin RQ'Q = \cos ROQ' = \cos a;$$

therefore $$\cot A \cot B = \cos a \cos b = \cos c. \qquad (5)$$

$$\frac{\cos B}{\cos b} = \frac{QR}{QP} \div \frac{RP'}{RP} = \frac{QR}{RP'} \cdot \frac{RP}{QP} = \frac{QR}{OR} \cdot \frac{OR}{RP'} \cdot \frac{RP}{OP} \cdot \frac{OP}{QP}$$

$$= \frac{\sin a}{\sin b} \cdot \frac{\sin b}{\sin c} = \frac{\sin a}{\sin c} = \sin A;$$

therefore $$\cos B = \sin A \cos b.$$

Similarly, $$\cos A = \sin B \cos a. \qquad (6)$$

45. **Napier's Rules of Circular Parts.**—The formulæ of the foregoing Article enable us, being given any two parts other than the right angle, to determine all the remaining parts of the triangle. They are learned by most students separately and in general terms. Napier, however, has given two rules, called after him, which embrace them all, and which are found by some to be useful aids to the memory; their utility, however, is much questioned by others. We give them here without any investigation, considering it sufficient to merely state them. The student may test them by deducing all the preceding formulæ from them, and, if anxious for further information regarding them, Napier's *Mirifici Logarithmorum*,

Canonis Descriptio, or Todhunter's *Spherical Trigonometry*, may be consulted.

Omitting the right angle, the two sides which include it and the complements of the remaining parts are termed the *Circular Parts* of the triangle. Thus the five circular parts are a, b, $\frac{\pi}{2} - A$, $\frac{\pi}{2} - c$, $\frac{\pi}{2} - B$. The student may remark that the order in which the circular parts are here written is that in which they naturally occur in the triangle.

Divide a circle (fig. 25) into five sectors, and on each sector write one of the circular parts, taking care to write them round the circle in the order they are written above; that is, in the order in which they occur in the triangle.






Fig. 25.

Any of the parts being selected and called the *middle parts*, the two parts contiguous to it are called the *adjacent parts*, and the remaining two are called the *opposite parts*. Thus, for example, if $\frac{\pi}{2} - c$ be selected for middle part,

then $\frac{\pi}{2} - A$ and $\frac{\pi}{2} - B$ will be the adjacent parts, and a and b will be the opposite parts.

This being premised, Napier's Rules are the following:—

I. Sine (middle part) = product of tangents of adjacent parts.

II. Sine (middle part) = product of cosines of opposite parts.

Examples.

1. Show that formula (1), Art. 44, reduces to Euc. I. XLVII. when the radius of the sphere is infinitely great.

[This appears at once on substituting the formulæ of Art. 18, and reducing.]

2. Show that formula (5) reduces to an identity *in plano*. Similarly, also formula (6) reduces to an identity.

3. Formulæ (2), (3), and (4) reduce to the ordinary expressions for the cosine, sine, and tangent of a plane angle respectively.

4. If c be less than 90°, show that a and b are of like affection.*

[In this case cos c is positive; and hence cos a and cos b must have like signs, as appears from formula (1).]

5. If c be greater than 90°, a and b are of opposite affection.

6. The sides of a right-angled triangle must be each less than a quadrant, or two of them must be each greater than a quadrant.

[This follows on combining Examples 4 and 5.]

7. Show that a side and the hypotenuse of a right-angled triangle are of the same or opposite affection, according as the included angle is less than, or greater than, a right angle.

[Apply formula (2), Art. 44.]

* Two angles are said to be of the same affection when they are each less than, or each greater than, a right angle.

8. The sides of a right-angled triangle are of the same affection as the opposite angles.

[By formula (6), $\cos a = \frac{\cos A}{\sin B}$; and therefore $\cos a$ is positive or negative, according as $\cos A$ is positive or negative.]

9. If one of the sides of a right-angled triangle be equal to the opposite angle, the remaining parts are each equal to 90°.

[See p. 51, Ex. 9.]

10. If the angle A of a right-angled triangle be acute, show that the difference of the sides which contain it is less than 90°.

[The opposite side (a) is less than 90°, but it is greater than the difference between the other two; therefore $c - b < 90°$.]

Section II.

Solution of Right-angled Triangles—Numerical Examples.

46. **Solution of Triangles.** — It has been already stated that every spherical triangle has six *parts*—namely, the three angles and the three sides, and that when any three of the parts are given, the remaining three can be determined.

The process by which the remaining three are determined is called the *solution* of the triangle.

In the practical solution of triangles the numerical calculations are performed by aid of a table of logarithms. Hence if we have at our disposal two or more formulæ, from which any required part can be determined in terms of other known parts, we should use that formula which is most readily adapted to logarithmic computation. If it should happen that a required part can only be obtained from a formula not adapted to logarithmic compu-

tation, it can be found by throwing the formula into logarithmic form by the use of a subsidiary angle. Any notice of this process is postponed, however, for the present.

We now proceed to the solution of all the cases which can be presented. They are six in number, and as before remarked, we require to know only two parts, the right angle forming a constant known part.

Remark.—In determining any of the parts, the use of the parts previously determined should be carefully avoided, lest they should be erroneous. Each part should be determined from the given parts alone.

It is always well to test the accuracy of the results by using the parts obtained to determine one of the given parts.

47. Case I.—*Having given the two sides a and b,* to determine A, B, and c, we have—

For A, $\cot A = \cot a \sin b$ [Formula (4), Art. 44.]
For B, $\cot B = \cot b \sin a$ „ „ „
For c, $\cos c = \cos a \cos b$ „ (1), „

In this case the solution is unique, as there is manifestly no ambiguity, none of the parts being determined from their sines.

Examples.

1. Given $a = 54° 16'$ and $b = 33° 12'$.

For A, $10 + L\cot A = L\cot(54° 16') + L\sin(33° 12')$;
therefore $L\cot A = 9{\cdot}5954382$,
and $A = 68° 29' 53''$.

For B, $10 + L\cot B = L\cot(33° 12') + L\sin(54° 16')$;
therefore $L\cot B = 10{\cdot}0935879$,
and $B = 38° 52' 26''$.

For c, $10 + L\cos c = L\cos a + L\cos b$;
therefore $L\cos c = 9{\cdot}6890261$,
and $c = 60° 44' 46''$.

2. Given $a = 55° 18'$ and $b = 39° 27'$.

$L\cot A = 9{\cdot}6434280$; $\therefore A = 66° 15' 6''$.
$L\cot B = 9{\cdot}9996157$; $\therefore B = 45° 1' 31''$.
$L\cos c = 9{\cdot}6430438$; $\therefore c = 63° 55' 21''$.

3. Given $a = 56° 20'$ and $b = 78° 40'$.

$L\cot A = 9{\cdot}8149722$; $\therefore A = 56° 51' 7''$.
$L\cot B = 9{\cdot}2222192$; $\therefore B = 80° 31' 48''$.
$L\cos c = 9{\cdot}0371914$; $\therefore c = 83° 44' 44\frac{1}{2}''$.

48. Case II.—*Having given the two angles A and B,* to determine a, b, and c, we have—

For a, $\cos A = \cos a \sin B$ [Formula (6), Art. 44.]
For b, $\cos B = \cos b \sin A$ „ „ „
For c, $\cos c = \cot A \cot B$ „ (5), „

In this case, as in Case I., there is no ambiguity.

Examples.

1. Given $A = 74° 15'$ and $B = 32° 10'$.

For a, $L\cos a = 10 + L\cos(74° 15') - L\sin(32° 10')$
$= 9{\cdot}7074497$;
therefore $a = 59° 20' 44''$.
For b, $L\cos b = 10 + L\cos(32° 10') - L\sin(74° 15')$
$= 9{\cdot}9442480$;
therefore $b = 28° 24' 54''$.

For c, $10 + L\cos c = L\cot A + L\cot B$.

$L\cos c = 9{\cdot}6516976$;

therefore $c = 63°\ 21'\ 24\frac{1}{2}''$.

2. Given $A = 52°\ 26'$ and $B = 49°\ 15'$.

$L\cos a = 9{\cdot}9056850$; $\therefore\ a = 36°\ 24'\ 34{\cdot}5''$.

$L\cos b = 9{\cdot}9156750$; $\therefore\ b = 34°\ 33'\ 40''$.

$L\cos c = 9{\cdot}8213599$; $\therefore\ c = 48°\ 29'\ 20''$.

3. Given $A = 64°\ 15'$ and $B = 48°\ 24'$.

$L\cos a = 9{\cdot}7641507$; $\therefore\ a = 54°\ 28'\ 53''$.

$L\cos b = 9{\cdot}8675405$; $\therefore\ b = 42°\ 30'\ 47''$.

$L\cos c = 9{\cdot}6316912$; $\therefore\ c = 64°\ 38'\ 38''$.

49. Case III.—*Having given the hypotenuse c and a side a*, to determine A, B, and b, we have—

For A, $\sin A = \sin a \div \sin c$ [Formula (3), Art. 44.]
For B, $\cos B = \tan a \div \tan c$ „ (2), „
For b, $\cos b = \cos c \div \cos a$ „ (1), „

In this case there is an apparent ambiguity in the value of A, but this is removed by the consideration that A and a are always of the same affection. (See Ex. 8, Art. 45).

Examples.

1. Given $c = 72°\ 30'$ and $a = 45°\ 15'$.

For A, $L\sin A = 10 + L\sin(45°\ 15') - L\sin(72°\ 30')$

$= 9{\cdot}8719522$;

therefore $A = 48°\ 7'\ 44{\cdot}5''$.

For B, $L\cos B = L\tan(45°\ 15') - L\tan(72°\ 30') + 10$

$= 9{\cdot}5025123$;

therefore $B = 71°\ 27'\ 15''$.

For b, $L\cos b = L\cos(72° 30') - L\cos(45° 15') + 10$
$= 9{\cdot}6305601.$
therefore $b = 64° 42' 52''.$

2. Given $c = 76° 42'$ and $a = 47° 18'$.

$L\sin A = 9{\cdot}8780442$; $\therefore A = 49° 2' 24{\cdot}5''.$
$L\cos B = 9{\cdot}4085340$; $\therefore B = 75° 9' 24{\cdot}75''.$
$L\cos b = 9{\cdot}5304897$; $\therefore b = 70° 10' 13''.$

3. Given $c = 91° 18'$ and $a = 72° 27'$.

$L\sin A = 9{\cdot}9794116$; $\therefore A = 72° 29' 48''.$
L cos sup. of $B = 8{\cdot}8558531$; $\therefore B = 94° 6' 53{\cdot}3''.$
L cos sup. of $b = 8{\cdot}8764415$; $\therefore b = 94° 18' 53{\cdot}8''.$

50. Case IV.—*Having given the hypotenuse c and an angle A,* to determine a, b, and B, we have—

For a, $\sin a = \sin A \sin c$ [Formula (3), Art. 44.]
For b, $\tan b = \cos A \tan c$ „ (2), „
For B, $\cot B = \tan A \cos c$ „ (5), „

The apparent ambiguity in the value of a is removed, as in Case III., by the consideration that a and A are of the same affection.

Examples.

1. Given $c = 84° 20'$ and $A = 35° 25'$.

For a, $L\sin a = L\sin(35° 25') + L\sin(84° 20') - 10$
$= 9{\cdot}7609396$;
therefore $a = 35° 13' 4''.$

For b, $L\tan b = L\cos(35° 25') + L\tan(84° 20') - 10$
$= 10{\cdot}9145116$;
therefore $b = 83° 3' 29''.$

For B, $L\cot B = L\tan(35° 25') + L\cos(84° 20') - 10$
$- 8{\cdot}8464279$;
therefore $B = 85° 59' 1''.$

2. Given $c = 67° 54'$ and $A = 43° 28'$.

$L \sin a = 9{\cdot}8044046$; $\therefore a = 39° 35' 51''$.
$L \tan b = 10{\cdot}2522138$; $\therefore b = 60° 46' 25\frac{1}{2}''$.
$L \cot B = 9{\cdot}5521908$; $\therefore B = 70° 22' 24''$.

3. Given $c = 22° 18' 30''$ and $A = 47° 39' 36''$.

$L \sin a = 9{\cdot}4480545$; $\therefore a = 16° 17' 41''$.
$L \tan b = 9{\cdot}4414574$; $\therefore b = 15° 26' 53''$.
$L \cot B = 10{\cdot}0065973$; $\therefore B = 44° 33' 53{\cdot}4''$.

51. Case V.—*Having given a side a and the adjacent angle B*, to determine A, b, and c, we have—

For A $\cos A = \cos a \sin B$ [Formula (6), Art. 44.]
For b, $\tan b = \tan B \sin a$ „ (4), „
For c, $\cot c = \cot a \cos B$ „ (2), „

In this case there is evidently no ambiguity.

Examples.

1. Given $a = 31° 20' 45''$ and $B = 55° 30' 30''$.

For A, $L \cos A = L \cos (31° 20' 45'') + L \sin (55° 30' 30'') - 10$
$= 9{\cdot}8475168$;
therefore $A = 45° 15' 30{\cdot}6''$.

For b, $L \tan b = L \tan (55° 30' 30'') + L \sin (31° 20' 45'') - 10$
$= 9{\cdot}8791734$;
therefore $b = 37° 7' 50''$.

For c, $L \cot c = L \cot (31° 20' 45'') + L \cos (55° 30' 30'') - 10$
$= 9{\cdot}9683434$;
therefore $c = 47° 5' 11''$.

2. Given $a = 82° 6'$ and $B = 43° 28'$.

$L \cos A = 8{\cdot}9756733$; $\therefore A = 84° 34' 28''$.

$L \tan b = 9{\cdot}9726026$; $\therefore b = 43° 11' 38''$.

$L \cot c = 9{\cdot}0030707$; $\therefore c = 84° 14' 57''$.

3. Given $a = 42° 30' 30''$ and $B = 53° 10' 30''$.

$L \cos A = 9{\cdot}7709180$; $\therefore A = 53° 50' 12''$.

$L \tan b = 9{\cdot}9554001$; $\therefore b = 42° 3' 47''$.

$L \cot c = 9{\cdot}8155179$; $\therefore c = 56° 49' 8''$.

52. Case VI.—*Having given a side a and the opposite angle A*, to determine *b*, *c*, and *B*, we have—

For b, $\sin b = \tan a \cot A$ [Formula (4), Art. 44.]
For c, $\sin c = \sin a \div \sin A$ „ (3), „
For B, $\sin B = \cos A \div \cos a$ „ (6), „

In this case an ambiguity arises as the parts are determined from their sines, and two values for each are in general admissible. However, for each value of b there will, in general, be only one value for c, since c and b are connected by the relation $\cos c = \cos a \cos b$; and at the same time only one admissible value for B, since $\cos c = \cot A \cot B$. Hence there will in general be only *two* triangles having the given parts, except when the side a is a quadrant and the angle A also 90°, in which case the solution becomes *indeterminate*.

When $a = A$, the formulæ, and also the figure, show that b, c, and B are right angles, and the solution is therefore *unique*.

It is easily seen that a is less than A when both are acute (they must be of the same affection), and greater than A when both are obtuse; and when the data do not satisfy these conditions the solution is impossible.

It appears at once from the figure that there are in general two solutions, for the *colunar* triangle $A'BC$ has also the given parts A' and a, and therefore satisfies the given conditions.

Examples.

1. Given $a = 46° 45'$ and $A = 59° 12'$.

For b, $L \sin b = L \tan (46° 45') + L \cot (59° 12') - 10$

$= 9{\cdot}8018795$;

therefore $b = 39° 19' 23\frac{1}{2}''$, or $140° 40' 36\frac{1}{2}''$.

For c, $L \sin c = L \sin a - L \sin A + 10$

$= 9{\cdot}9283797$;

therefore $c = 57° 59' 29''$, or $122° 0' 31''$.

For B, $L \sin B = L \cos A - L \cos a + 10$

$= 9{\cdot}8734997$;

therefore $B = 48° 21' 28''$, or $131° 38' 32''$.

2. Given $a = 21° 39'$ and $A = 42° 10' 10''$.

$L \sin b = 9{\cdot}6417030 = 25° 59' 27{\cdot}8''$, or $154° 0' 32{\cdot}2'$.

$L \sin c = 9{\cdot}7400177 = 33° 20' 13{\cdot}4''$, or $146° 39' 46{\cdot}6''$.

$L \sin B = 9{\cdot}9016853 = 52° 23' 2{\cdot}8''$, or $127° 36' 57{\cdot}2'$.

3. Given $a = 25° 18' 45''$ and $A = 15° 58' 15''$.

[*Ans.* Impossible by Art. 52.]

Section III.

Theorems.

53. In this section we propose to investigate some theorems which apply to spherical triangles in general, but which are immediately deducible from a knowledge of the properties of right-angled triangles.

54. *The arcs drawn from the vertices of a spherical triangle perpendicular to the opposite sides are concurrent.*

Let AP, BQ (fig. 26), be arcs drawn from A, B, per-




Fig. 26.

pendicular to BC and CA, respectively, and let O be their point of intersection.

Now
$$\frac{\sin OQ}{\sin OR} = \frac{\sin OAQ}{\sin OAR} = \frac{\sec B}{\sec C}$$

(formula (6), Art. 44); and it is obvious that the sines of the perpendiculars from any other point of AP on the sides b and c bear the same ratio to each other.

Similarly, $$\frac{\sin OP}{\sin OR} = \frac{\sec A}{\sec C};$$

and hence, on dividing this equation by the former, we get

$$\frac{\sin OP}{\sin OQ} = \frac{\sec A}{\sec B}$$

and therefore the arc joining C to O is perpendicular to the side AB. Thus the perpendiculars meet in a point; and if α, β, γ be the lengths intercepted on them between their point of concurrence and the sides a, b, and c, respectively, we have

$$\sin \alpha : \sin \beta : \sin \gamma = \sec A : \sec B : \sec C.$$

In the same manner, it may be shown that the arcs joining the vertices to the middle points of the opposite sides are concurrent; and if α', β', γ' be the perpendiculars from their point of concurrence on the sides a, b, c, we have

$$\sin \alpha' : \sin \beta' : \sin \gamma' = \operatorname{cosec} A : \operatorname{cosec} B : \operatorname{cosec} C.$$

Similarly for point of intersection of perpendicular at middle points of sides—

$$\sin \alpha'' : \sin \beta'' : \sin \gamma'' = \cos (S-A) : \cos (S-B) : \cos (S-C).$$

55. *If from three points A, B, C on a great circle perpendiculars AA', BB', CC' be drawn to another great circle—*

(1) $\sin BC \sin AA' + \sin CA \sin BB' + \sin AB \sin CC' = 0,$
(2) $\sin B'C' \tan AA' + \sin C'A' \tan BB' + \sin A'B' \tan CC' = 0,$
(3) $\sin BC \cot A + \sin CA \cot B + \sin AB \cot C = 0,$
(4) $\sin B'C' \cos A + \sin C'A' \cos B + \sin A'B' \cos C = 0,$

where A, B, C are the angles the perpendiculars AA', BB', CC' make with the great circle ABC.

To prove (1)—

$$\frac{\sin AA'}{\sin AD} = \frac{\sin BB'}{\sin BD} = \frac{\sin CC'}{\sin CD} = \sin D \text{ (fig. 27).} \qquad (\alpha)$$

But since A, B, C are concyclic (p. 11, Ex. 15),

$$\sin BC \sin AD + \sin CA \sin BD + \sin AB \sin CD = 0.$$




Fig. 27.

Substituting in this expression the values of $\sin AD$, $\sin BD$, and $\sin CD$, obtained from the equations (α), we obtain (1).

To prove (2)—

$$\frac{\tan AA'}{\sin A'D} = \frac{\tan BB'}{\sin B'D} = \frac{\tan CC'}{\sin C'D} = \tan D. \qquad (\beta)$$

But, as before,

$$\sin B'C' \sin A'D + \sin C'A' \sin B'D + \sin A'B' \sin C'D = 0.$$

Substituting in this expression the values of $\sin A'D$, $\sin B'D$, and $\sin C'D$, obtained from the equations (β), we obtain (2).

To prove (3)—

$$\frac{\cos AD}{\cot A} = \frac{\cos BD}{\cot B} = \frac{\cos CD}{\cot C} = \cot D.$$

But

$$\sin BC \cos AD + \sin CA \cos BD + \sin AB \cos CD = 0;$$

therefore, &c. [See p. 77, Ex. 42.]

To prove (4)—

$$\frac{\cos A}{\cos A'D} = \frac{\cos B}{\cos B'D} = \frac{\cos C}{\cos C'D} = \sin D.$$

[Formula (6), Art. 44.]

But

$$\sin B'C' \cos A'D + \sin C'A' \cos B'D + \sin A'B' \cos C'D = 0;$$

therefore, &c.

By observing that AA', BB', and CC' meet at P, the pole of $A'B'C'$, we are enabled to write down the following corollaries:—

Cor. 1.—If B be a point on the base AC of a triangle APC, we have

$$\sin BC \cos AP + \sin CA \cos BP + \sin AB \cos CP = 0.$$

[Cf. Art. 40.]

Cor. 2.—

$$\cot AP \sin BPC + \cot BP \sin CPA + \cot CP \sin APB = 0.$$

[Cf. p. 58, Ex. 10.]

Cor. 3.—

$$\cos A \sin BPC + \cos B \sin CPA + \cos C \sin APB = 0.$$

[Cf. p. 46, Ex. 5.]

NOTE.—It may be noticed that (1) and (4) are supplemental theorems, as are also (2) and (3). When an arc is drawn from the vertical angle of a triangle to meet the base, (1) connects it with the segments of the base; (2) connects it with the segments of the vertical angle; (3) connects the angle it makes with the base with the segments of the base; and (4) connects the angle it makes with the base with the segments of the vertical angle, the parts of the triangle being supposed known.

56. **Geometrical Deduction of the Analogies of Napier and Delambre.**—*Construction.* On the base AB of the triangle ABC (fig. 28) construct an isosceles triangle AOB, having each base angle $= \frac{1}{2}(A + B)$. From its vertex O draw OX, OY, OZ perpendiculars to the sides a, b, and c of the given triangle. Join OC. Draw AP and BP perpendicular arcs to AO and BO. The arc OZ, on being produced, evidently passes through P.

Now the triangles BOX and AOY are equal in all respects; for the side BO and angle X are equal to the side AO and the angle Y, respectively; and the angle $OBX = B - \frac{1}{2}(A + B) = \frac{1}{2}(B - A) =$ angle OAY. Therefore the triangles are equal in all respects, and as a consequence the triangles COX and COY are equal in all respects.

Hence, $$AY = BX = \tfrac{1}{2}(a + b),$$

and $$CX = CY = \tfrac{1}{2}(a - b).$$

But since the angles AOY and BOX are equal, it follows

that the angles AOB and XOY are equal, and hence their halves AOZ and COY are equal.




Fig. 28.

Again, if from P perpendiculars PX', PY', and PZ be let fall on the sides, it will follow, as before,

that $AY' = BX' = \frac{1}{2}(a - b)$,

and $CX' = CY' = \frac{1}{2}(a + b)$;

also P is on the internal bisector of the angle C, and the angle APY' is equal to the angle BPX'.

Firstly—*Napier's Analogies:*

1. $$\frac{\cos AP}{\cos CP} = \frac{\cos AY'}{\cos CY'}$$

[From the triangles PAY' and PCY', Art. 44 (1).]

$$= \frac{\cot PAZ}{\cot PCY}$$

[From the triangles PAZ and PCY', Art. 44 (5)].

Hence,

$$\tan \tfrac{1}{2}(A+B) = \frac{\cos \frac{1}{2}(a-b)}{\cos \frac{1}{2}(a+b)} \cot \frac{C}{2}.$$

2. $\tan OY = \tan OAY \sin AY = \tan OCY \sin CY.$

[Art. 44 (4).]

Hence,

$$\tan \tfrac{1}{2}(A-B) = \frac{\sin \frac{1}{2}(a-b)}{\sin \frac{1}{2}(a+b)} \cot \frac{C}{2}.$$

3. $$\tan AO = \frac{\tan AZ}{\cos OAZ} = \frac{\tan AY}{\cos OAY}.$$

[Art. 44 (2).]

Hence,

$$\tan \tfrac{1}{2}(a+b) = \frac{\cos \frac{1}{2}(A-B)}{\cos \frac{1}{2}(A+B)} \tan \frac{c}{2}.$$

4. $$\tan AP = \frac{\tan AY'}{\cos PAY'} = \frac{\tan AZ}{\cos PAZ}.$$

Hence,

$$\tan \tfrac{1}{2}(a-b) = \frac{\sin \frac{1}{2}(A-B)}{\sin \frac{1}{2}(A+B)} \tan \frac{c}{2}.$$

Secondly—*Delambre's* Analogies:*


1. $\cos APZ = \cos AZ \sin PAZ = \cos CY' \sin PCY'.$

[Art. 44 (6).]

Hence,

$$\cos \tfrac{1}{2}(A+B) \cos \frac{c}{2} = \cos \tfrac{1}{2}(a+b) \sin \frac{C}{2}.$$

* Delambre's Analogies are commonly called Gauss's Formulæ. (See foot note, p. 43.)



3. Prove that

$$\sin^2 a \sin^2 b = \sin^2 a + \sin^2 b - \sin^2 c. \qquad \text{[Art. 44 (1).]}$$

4. Prove that

$$\tan^2 \frac{A}{2} = \frac{\sin (c-b)}{\sin (c+b)}.$$

[This is obviously another form of (2), Art. 44.]

5. What is the Great Circle course and distance between two places in latitude 45° North, with 90° difference of longitude?

Ans. Course makes 60° with meridian. Distance $= \cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$.

6. If p be the perpendicular from the right angle on the hypotenuse, prove that—

(*a*) $\cos^2 p = \cos^2 A + \cos^2 B$. [Art. 44 (6.)]

(*b*) $\cot^2 p = \cot^2 a + \cot^2 b$. [Art. 44 (4).]

(*c*) $\sin^2 p \sin^2 c = \sin^2 a + \sin b^2 - \sin^2 c$. [See Ex. 3.]

7. Express the angle A in terms of—(1) b and c; (2) c and a; (3) a and b.

8. Show that

$$2 \sin^2 \tfrac{1}{2} c = \sin^2 \tfrac{1}{2} (a + b) + \sin^2 \tfrac{1}{2} (a - b).$$

[Reduces to (1), Art. 44.]

9. In a spherical triangle, if $c = 90°$, prove that

$$\tan a \tan b + \sec C = 0.$$

[Supplemental to (5), Art. 44.]

10. If the side c of a spherical triangle be a quadrant, show that

$$\sin^2 p = \cot \theta \cot \phi,$$

where p is the perpendicular on c, and θ and ϕ are the segments of the vertical angle.

11. Show that the ratio of the cosines of the segments of the base made by the perpendicular from the vertex is equal to the ratio of the cosines of the sides.

[Apply (1), Art. 44.]

12. Given the base and the ratio of the cosines of the sides of any spherical triangle; find the locus of the vertex, and state the analogue *in plano*.

[See Ex. 11.]

13. A station is in latitude λ (North), and longitude l (East); find the longitudes of the places on the Equator distant δ from the station.

[*Science and Art Exam. Papers.*]

Ans. Longitudes $= l \pm \cos^{-1}\left(\frac{\cos\delta}{\cos\lambda}\right)$.

14. Given the base and base angles of any spherical triangle; calculate the segments x and $c - x$ of the base made by the perpendicular on it from the vertex.

Ans. $\tan A \sin x = \tan B \sin (c - x)$.

15. What is the shortest distance between two places on the earth's surface—one in latitude 45° North and longitude 45° West, the other in latitude 45° South and latitude 45° East?

Ans. 120°.

16. Given in a right-angled triangle the vertex fixed, and

$$\cot^2 a + \cot^2 b = \text{const.};$$

find the envelope of the base.

[See Ex. 6 (b).]

17. Show that the ratio of the cosines of the base angles is equal to the ratio of the sines of the segments of the vertical angle made by the perpendicular drawn from it to the opposite side.

18. Given the vertical angle and ratio of cosines of the base angles; find the envelope of the base.

Ans. A fixed point 90° from the vertex; its position being obtained from Ex. 17.

19. If $A = 36°$ and $B = 60°$, show that—

(1). $a + b + c = 90°$.
(2). The perpendicular on the hypotenuse $= 18°$.
(3). $\cot a - \cot b = 1$.

20. If λ and δ be perpendiculars from a point S on two great circles intersecting at an angle ω, and if the intercepts they make on them from their point of intersection be l and a, show that

$$\cos\omega = \frac{\sin\lambda \tan l + \sin\delta \tan a}{\sin\lambda \tan a + \sin\delta \tan l}.$$

[Professor Haughton, *Educational Times.*]

[Apply the formulæ of Art. 37 to the triangle SPK, where P and K are the poles of the great circles, and eliminate $\sin\omega$.]

21. Using the notation of Ex. 20, show that—

(1). $$\frac{\cos \frac{1}{2}(\omega - \omega')}{\cos \frac{1}{2}(\omega + \omega')} = \frac{\tan \frac{1}{2}\lambda \tan \frac{1}{2} l + \tan \frac{1}{2}\delta \tan \frac{1}{2} a}{\tan \frac{1}{2} l \tan \frac{1}{2}\delta + \tan \frac{1}{2}\lambda \tan \frac{1}{2} a}.$$

(2). $$\frac{\sin \frac{1}{2}(\omega - \omega')}{\sin \frac{1}{2}(\omega + \omega')} = \frac{\tan \frac{1}{2}\delta \tan \frac{1}{2}\lambda - \tan \frac{1}{2} l \tan \frac{1}{2} a}{1 - \tan \frac{1}{2} a \tan \frac{1}{2} l \tan \frac{1}{2}\delta \tan \frac{1}{2}\lambda},$$

ω' being the angle between δ and λ.

Miscellaneous Examples.

In the following Examples the triangle is supposed right-angled (at C), unless otherwise stated.

1. If β be the bisector of the hypotenuse of a right-angled triangle, show that—

$$\sin^2 \beta = \frac{\sin^2 a + \sin^2 b}{4 \cos^2 \frac{c}{2}}.$$



[By Art. 42—

$$\cos a + \cos b = 2 \cos \beta \cos \frac{c}{2}.$$

Squaring we get

$$\cos^2 a + \cos^2 b + 2 \cos c = 2 (1 - \sin^2 \beta)(1 + \cos c); \therefore \text{ \&c.}]$$

2. If ι and η be the internal and external bisectors of C, prove that—

(a) $2 \cot^2 \iota = (\cot a + \cot b)^2$,

(b) $2 \cot^2 \eta = (\cot a - \cot b)^2$. [See p. 55.]

3. Show that—

(a) $\cot^2 \iota + \cot^2 \eta = 2 \cot^2 a + \cot^2 b$,

(b) $\cot^2 \iota - \cot^2 \eta = 2 \cot a \cot b$.

4. If the bisectors of the vertical angle C meet the base in X and Y, show that

$$2 \cos XY = (\cot^2 a - \cot^2 b) \sin \iota \sin \eta.$$

[Cf. p. 57, Ex. 6.]

5. If ϕ be the angle between the bisector of the vertical angle and the perpendicular, show that

$$\tan \phi = \frac{\sin (a - b)}{\sin (a + b)}.$$ [See p. 69, Ex. 11.]

6. In a right-angled triangle, show that

$$\sin(A+B) = \frac{\cos a + \cos b}{1+\cos a \cos b}.$$

[Apply Delambre's formulæ.]

7. Show that

$$\tan S = \cot\frac{a}{2}\cot\frac{b}{2}.$$

[Here $$\tan S = \frac{1+\tan\frac{A+B}{2}}{1-\tan\frac{A+B}{2}}.$$

[Apply Napier's Analogies.]

8. Construct a right-angled triangle, being given the hypotenuse and—

(1). The sum of the base angles;

(2). The difference of the base angles.

[Apply Gauss's formulæ.]

9. Given the hypotenuse and the sum or difference of the sides; construct the triangle. (See Ex. 8.)

10. Given the sum of the sides a and b, and the sum of the base angles; solve the triangle.

[Apply Gauss's formulæ.]

11. Show that

$$\sin\frac{A}{2} = \frac{\sqrt{\sin c + \sin a} + \sqrt{\sin c - \sin a}}{2\sqrt{\sin c}}.$$

[Here, if we divide out by $\sqrt{\sin c}$, and put $\frac{\sin a}{\sin c} = \sin A$, the right-hand side becomes $\frac{1}{2}(\sqrt{1+\sin A}+\sqrt{1-\sin A})$. But $1 \pm \sin A = \left(\sin\frac{A}{2} \pm \cos\frac{A}{2}\right)^2$; therefore, &c.]

12. If a, a', β, β', γ, γ' be the segments of the sides of any spherical triangle made by the perpendiculars from the opposite vertex, show that

$$\cos a \cos\beta \cos\gamma = \cos a' \cos\beta' \cos\gamma'.$$

[For $$\frac{\cos a}{\cos a'} = \frac{\cos a}{\cos b}; \quad \frac{\cos\beta}{\cos\beta'} = \frac{\cos b}{\cos c}; \quad \frac{\cos\gamma}{\cos\gamma'} = \frac{\cos c}{\cos a}.$$

(See p. 99, Ex. 11). Hence, $\frac{\cos a \cos\beta \cos\gamma}{\cos a' \cos\beta' \cos\gamma'} = 1$; therefore, &c.]

13. Using the notation of the preceding Example, show that—

(a) $\sin\alpha \sin\beta \sin\gamma = \sin\alpha' \sin\beta' \sin\gamma'$,

(b) $\tan\alpha \tan\beta \tan\gamma = \tan\alpha' \tan\beta' \tan\gamma'$.

[(a) follows from the formula $\tan p = \tan A \sin a = \tan B \sin a'$.]

14. Find the distance between any two points on the Earth's surface, given their latitudes and longitudes.

If λ and λ' be their latitudes, and l and l' their longitudes, we have

$$\cos\delta = \sin\lambda \sin\lambda' + \cos\lambda \cos\lambda' \cos(l - l').$$

15. The cosines of the base angles of any triangle are proportional to the cosines of angles which the sides make with the arc joining the vertex to the pole of the perpendicular from the vertex on the base.

[For these latter cosines are equal to the sines of the segments of the vertical angle made by the perpendicular.]

16. If δ be the length of the arc through the vertex of an isosceles triangle, dividing the base into segments α and β, prove that

$$\tan\frac{\alpha}{2} \tan\frac{\beta}{2} = \tan\frac{a+\delta}{2} \tan\frac{a-\delta}{2},$$

where a is one of the equal sides of the triangle.

[See Art. 43, or apply Napier's Analogies.]

17. Find a relation connecting the mutual distances of any three places with their latitudes.

[By the aid of Ex. 14. See also Art. 55 (1).]

18. On the Earth considered as a sphere a great circle is drawn, inclined to the Equator at an angle of 30°, and cutting it in points whose longitudes are 0° and 180°. Find the longitudes of the meridians, dividing the great circle into eight equal parts.

19. Two points A and B are taken on two secondaries to a great circle $A'B'$ If θ be the angle between the arcs AB and $A'B'$, its value is found by the equation (*Q. U. I. Exam. Papers*)—

$$\sin A'B' = \frac{\sin AB \cos\theta}{\cos AA' \cos BB'}.$$

$$\left[\frac{\cos\theta}{\cos AA' \cos BB'} = \frac{\sin A}{\cos BB'} = \frac{\sin A}{\sin BC} = \frac{\sin A'B'}{\sin AB}; \text{ therefore, \&c.}\right]$$

CHAPTER V.

OBLIQUE-ANGLED TRIANGLES.

57. In the present Chapter we propose to consider the different cases which occur in the solution of oblique-angled triangles. As in right-angled triangles (Chap. IV.) there are six distinct cases, when we are given three of the parts, and are required to determine the remaining three.

Before proceeding to the general solutions of the triangles, we may notice that from particular values of the given parts the values of the other parts may be made to depend on the solutions of right-angled triangles, as for example—

α. Let the side a be a quadrant.

Here the supplemental triangle is right-angled; therefore, &c.

β. Let the sides a and b be equal.

The triangle is divided into two equal right-angled triangles by the perpendicular to the base passing through the vertex; therefore, &c.

γ. The general solution of the equilateral triangle may be derived by aid of the *logarithmic* equation

$$\frac{\tan \frac{a}{2}}{\tan a} = \cos A, \text{ or } 2 \cos A = \sec^2 \frac{a}{2}.$$

58. The six cases (Art. 57) which present themselves are the following:—

I. Having given three sides,	a, b, c;
II. Having given three angles,	A, B, C;
III. Having given two sides and the included angle,	a, C, b;
IV. Having given two angles and the adjacent side,	A, c, B;
V. Having given two sides and an angle opposite either,	a, b, A;
VI. Having given two angles and a side opposite either,	A, B, a;

but these are immediately resolved into *three* distinct problems by the aid of the polar triangle.

For, when three sides are given, and the angles of the triangle are required, the data applied to the polar triangle transforms the problem into a supplemental problem, viz.—Having given three angles of a triangle, find the sides.

Similarly, Cases III. and IV. are supplemental, likewise V. and VI.

59. When numerical values are assigned to the given parts, in order to ascertain the remaining parts in degrees, minutes, and seconds, we employ formulæ adapted to logarithmic computation.

All the formulæ given in Chap. III. are either in logarithmic form or can immediately be thrown into the desired shape by means of a *subsidiary angle*. For the present we select the logarithmic formulæ, and apply

them to the following numerical examples, while we reserve the remaining formulæ for future discussion. (See Art. 76.)

60. Case I.—*Having given the three sides, a, b, c.* Here we have (Art. 33),

$$\tan\frac{A}{2} = \sqrt{\frac{\sin(s-b)\sin(s-c)}{\sin s \sin(s-a)}}$$

$$= \frac{1}{\sin(s-a)}\sqrt{\frac{\sin(s-a)\sin(s-b)\sin(s-c)}{\sin s}}.$$

The part under the radical, being a symmetric function of the sides, appears on the right-hand side of the equations for determining A, B, C, and when once calculated is utilized in the calculation of each angle.

The logarithmic form of the above equation is

$$L\tan\frac{A}{2} = 10 - L\sin(s-a) + \tfrac{1}{2}\{L\sin(s-a) + L\sin(s-b) + L\sin(s-c) - L\sin s\}.$$

Examples.

(1). Given $a = 46° 24'$, $b = 67° 14'$, $c = 81° 12'$.

$L\sin s = L\sin 97° 25' = 9{\cdot}9963513.$
$L\sin(s-a) = L\sin 51° 1' = 9{\cdot}8906049.$
$L\sin(s-b) = L\sin 30° 11' = 9{\cdot}7013681.$
$L\sin(s-c) = L\sin 16° 13' = 9{\cdot}4460251.$

$L\tan\frac{1}{2}A = 9{\cdot}6302185$; $\therefore A = 46° 13' 30''.$
$L\tan\frac{1}{2}B = 9{\cdot}8194553$; $\therefore B = 66° 50' 20''.$
$L\tan\frac{1}{2}C = 10{\cdot}0747984$; $\therefore C = 99° 49' 10''.$

(2). Given $a = 108° 14'$, $b = 75° 29'$, $c = 56° 37'$.

$$L \sin s = L \sin 120° 10' = 9{\cdot}9367988.$$
$$L \sin (s-a) = L \sin 11° 56' = 9{\cdot}3154947.$$
$$L \sin (s-b) = L \sin 44° 41' = 9{\cdot}8470714.$$
$$L \sin (s-c) = L \sin 63° 33' = 9{\cdot}9519799.$$

$$L \tan \tfrac{1}{2} A = 10{\cdot}2733789 ; \therefore A = 123° 53' 47''.$$
$$L \tan \tfrac{1}{2} B = 9{\cdot}7418022 ; \therefore B = 57° 46' 56''.$$
$$L \tan \tfrac{1}{2} C = 9{\cdot}6368937 ; \therefore C = 46° 51' 51{\cdot}5''.$$

(3). Given $a = 57° 17'$, $b = 20° 39'$, $c = 76° 22'$.

$$L \sin s = L \sin 77° 9' = 9{\cdot}9889849.$$
$$L \sin (s-a) = L \sin 19° 52' = 9{\cdot}5312649.$$
$$L \sin (s-b) = L \sin 56° 30' = 9{\cdot}9211066.$$
$$L \sin (s-c) = L \sin 0° 47' = 8{\cdot}1358104.$$

$$\tan \tfrac{1}{2} A = 9{\cdot}2683336 ; \therefore A = 21° 1' 2''.$$
$$L \tan \tfrac{1}{2} B = 8{\cdot}8784919 ; \therefore B = 8° 38' 46''.$$
$$L \tan \tfrac{1}{2} C = 10{\cdot}6637881 ; \therefore C = 155° 31' 36{\cdot}5''.$$

(4). Given $a = 68° 45'$, $b = 53° 15'$, $c = 46° 30'$.

$$L \sin s = L \sin 84° 15' = 9{\cdot}9978093.$$
$$L \sin (s-a) = L \sin 15° 30' = 9{\cdot}4268988.$$
$$L \sin (s-b) = L \sin 31° 0' = 9{\cdot}7118393.$$
$$L \sin (s-c) = L \sin 37° 45' = 9{\cdot}7869056.$$

$$L \tan \tfrac{1}{2} A = 10{\cdot}0370184 ; \therefore A = 94° 52' 40''.$$
$$L \tan \tfrac{1}{2} B = 9{\cdot}7520779 ; \therefore B = 58° 56' 10''.$$
$$L \tan \tfrac{1}{2} C = 9{\cdot}6770116 ; \therefore C = 50° 50' 52\tfrac{1}{2}''.$$

(5). Given $a = 63° 54'$, $b = 47° 18'$, $c = 53° 26'$.

$$L \sin s = L \sin 82° 19' = 9{\cdot}9960834.$$
$$L \sin (s-a) = L \sin 18° 25' = 9{\cdot}4995840.$$
$$L \sin (s-b) = L \sin 35° 1' = 9{\cdot}7587717.$$
$$L \sin (s-c) = L \sin 28° 53' = 9{\cdot}6839720.$$

$L \tan \frac{1}{2} A = 9{\cdot}9735382$; $\therefore A = 86° 30' 40''$.

$L \tan \frac{1}{2} B = 9{\cdot}7143505$; $\therefore B = 54° 46' 14''$.

$L \tan \frac{1}{2} C = 9{\cdot}7891502$; $\therefore C = 63° 12' 55\frac{1}{2}''$.

61. Case II.—*Having given the three angles A, B, C.* Here we have (Art. 36),

$$\tan \frac{a}{2} = \sqrt{\frac{-\cos S \cos (S-A)}{\cos (S-B) \cos (S-C)}}$$

$$= \cos (S-A) \sqrt{\frac{-\cos S}{\cos (S-A) \cos (S-B) \cos (S-C)}},$$

with similar values for $\tan \frac{b}{2}$ and $\tan \frac{c}{2}$.

Examples.

(1). Given $A = 68° 30'$, $B = 74° 20'$, $C = 83° 10'$.

$L (-\cos S) = L \cos 67° 0' = 9{\cdot}5918780.$

$L \cos (S-A) = L \cos 44° 30' = 9{\cdot}8532421.$

$L \cos (S-B) = L \cos 38° 40' = 9{\cdot}8925365.$

$L \cos (S-C) = L \cos 29° 50' = 9{\cdot}9382576.$

$L \tan \frac{a}{2} = 9{\cdot}8071630$; $\therefore a = 65° 21' 22\frac{1}{2}''$.

$L \tan \frac{b}{2} = 9{\cdot}8464574$; $\therefore b = 70° 9' 9\frac{1}{2}''$.

$L \tan \frac{c}{2} = 9{\cdot}8921785$; $\therefore c = 75° 55' 9''$.

(2). Given $A = 86° 20'$, $B = 76° 30'$, $C = 94° 40'$.

$L (-\cos S) = L \cos 51° 15' = 9{\cdot}7965212.$

$L \cos (S-A) = L \cos 42° 25' = 9{\cdot}8682088.$

$L \cos (S-B) = L \cos 52° 15' = 9{\cdot}7869056.$

$L \cos (S-C) = L \cos 34° 5' = 9{\cdot}9181475.$

$\tan \frac{1}{2} a = 9{\cdot}9798385$; $\therefore a = 87° 20' 28''$.
$\tan \frac{1}{2} b = 9{\cdot}8985353$; $\therefore b = 76° 44' 2\frac{1}{2}''$.
$\tan \frac{1}{2} c = 10{\cdot}0297772$; $\therefore c = 93° 55' 31''$.

(3). Given $A = 96° 45'$, $B = 108° 30'$, $C = 116° 15'$.

$L(-\cos S) = L \cos 19° 45' = 9{\cdot}9750129$.
$L \cos (S - A) = L \cos 64° 0' = 9{\cdot}6418420$.
$L \cos (S - B) = L \cos 52° 15' = 9{\cdot}7869056$.
$L \cos (S - C) = L \cos 44° 30' = 9{\cdot}9532421$.

$\tan \frac{1}{2} a = 9{\cdot}9883536$; $\therefore a = 88° 27' 49''$.
$\tan \frac{1}{2} b = 10{\cdot}1334172$; $\therefore b = 107° 19' 52''$.
$\tan \frac{1}{2} c = 10{\cdot}1997537$; $\therefore c = 115° 28' 13\frac{1}{4}''$.

(4). Given $A = 78° 30'$, $B = 118° 40'$, $C = 93° 20'$.

$L(-\cos S) = L \cos 34° 45' = 9{\cdot}9146852$.
$L \cos (S - A) = L \cos 66° 45' = 9{\cdot}5963154$.
$L \cos (S - B) = L \cos 26° 35' = 9{\cdot}9514757$.
$L \cos (S - C) = L \cos 51° 55' = 9{\cdot}7901493$.

$\tan \frac{1}{2} a = 9{\cdot}8846878$; $\therefore a = 74° 57' 46''$.
$\tan \frac{1}{2} b = 10{\cdot}2398481$; $\therefore b = 120° 8' 49''$.
$\tan \frac{1}{2} c = 10{\cdot}0785217$; $\therefore c = 100° 18' 11\frac{1}{4}''$.

(5). Given $A = 57° 50'$, $B = 98° 20'$, $C = 63° 40'$.

$L(-\cos S) = L \cos 70° 5' = 9{\cdot}5323123$.
$L \cos (S - A) = L \cos 52° 5' = 9{\cdot}7885323$.
$L \cos (S - B) = L \cos 11° 35' = 9{\cdot}9910637$.
$L \cos (S - C) = L \cos 46° 15' = 9{\cdot}8398004$.

$\tan \frac{1}{2} a = 9{\cdot}7449903$; $\therefore a = 58° 8' 19''$.
$\tan \frac{1}{2} b = 9{\cdot}9475217$; $\therefore b = 83° 5' 36''$.
$\tan \frac{1}{2} c = 9{\cdot}7962584$; $\therefore c = 64° 3' 20''$.

62. Case III.—*Having given two sides and the included angle, a, C, b.*

This case is somewhat similar to the corresponding one in *Plane Trig.*, the base angles being calculated from two separate equations, one giving half the sum and the other half the difference of those angles. The formulæ to be employed have been arrived at on p. 43 (Note to Ex. 8). They have, moreover, been demonstrated geometrically from the relations between the sides and angles of certain right-angled triangles. As they are of fundamental importance in the solution of triangles, we shall now proceed by a direct method to obtain the same results.

63. **Napier's Analogies:**

Let $$x = \frac{\sin A}{\sin a} = \frac{\sin B}{\sin b};$$

hence,

$$x = \frac{\sin A \pm \sin B}{\sin a \pm \sin b}. \qquad (1)$$

Now

$$\cos A + \cos B \cos C = \sin B \sin C \cos a = x \sin C \sin b \cos a,$$

and

$$\cos B + \cos C \cos A = \sin C \sin A \cos b = x \sin C \sin a \cos b.$$

By addition,

$$(\cos A + \cos B)(1 + \cos C) = x \sin C \sin (a + b); \qquad (2)$$

by subtraction,

$$(\cos B - \cos A)(1 - \cos C) = x \sin C \sin (a - b). \qquad (3)$$

On taking the positive sign in (1), and dividing by (2), we obtain,

$$\frac{\sin A + \sin B}{\cos A + \cos B} = \frac{\sin a + \sin b}{\sin (a+b)} \cdot \frac{1 + \cos C}{\sin C};$$

or

$$\tan \tfrac{1}{2}(A+B) = \frac{\cos \tfrac{1}{2}(a-b)}{\cos \tfrac{1}{2}(a+b)} \cot \frac{C}{2}. \qquad (\alpha')$$

On taking the negative sign in (1), we obtain, by a similar method,

$$\tan \tfrac{1}{2}(A-B) = \frac{\sin \tfrac{1}{2}(a-b)}{\sin \tfrac{1}{2}(a+b)} \cot \frac{C}{2}. \qquad (\beta')$$

Employing the supplemental formulæ, viz.,

$$\cos a - \cos b \cos c = \sin b \sin c \cos A = \frac{1}{x} \sin B \cos A \sin c,$$

and

$$\cos b - \cos c \cos a = \sin c \sin a \cos B = \frac{1}{x} \sin A \cos B \sin c,$$

we have, as before,

$$x.(\cos a \pm \cos b)(1 \mp \cos c) = \sin c \sin (B \pm A).$$

Dividing each of these equations by results in equation (1) to eliminate x, we have formulæ supplemental to (α') and (β'), viz.,

$$\tan \tfrac{1}{2}(a+b) = \frac{\cos \tfrac{1}{2}(A-B)}{\cos \tfrac{1}{2}(A+B)} \tan \frac{c}{2}, \qquad (\gamma')$$

$$\tan \tfrac{1}{2}(a-b) = \frac{\sin \tfrac{1}{2}(A-B)}{\sin \tfrac{1}{2}(A+B)} \tan \frac{c}{2}. \qquad (\delta')$$

64. **Delambre's (or Gauss's) Analogies:**

By the preceding Article, we have

$$\tan \tfrac{1}{2}(A+B) \tan \frac{C}{2} = \frac{\cos \tfrac{1}{2}(a-b)}{\cos \tfrac{1}{2}(a+b)}.$$

Hence,

$$1 - \tan \tfrac{1}{2}(A+B) \tan \tfrac{1}{2} C = 1 - \frac{\cos \tfrac{1}{2}(a-b)}{\cos \tfrac{1}{2}(a+b)};$$

or

$$\frac{-\cos S}{\cos \tfrac{1}{2}(A+B) \cos \tfrac{1}{2} C} = \frac{2 \sin \tfrac{1}{2} a \sin \tfrac{1}{2} b}{\cos \tfrac{1}{2}(a+b)}; \qquad (1)$$

but

$$-\cos S = \frac{\sin \tfrac{1}{2} a \sin \tfrac{1}{2} b \sin C}{\cos \tfrac{1}{2} c}.$$

[Page 74, Ex. 28.]

Substituting this value in (1), we have at once

$$\cos \tfrac{1}{2}(A+B) \cos \tfrac{1}{2} c = \cos \tfrac{1}{2}(a+b) \sin \tfrac{1}{2} C. \qquad (a)$$

Again,

$$1 + \tan \tfrac{1}{2}(A-B) \tan \tfrac{1}{2} C = 1 + \frac{\sin \tfrac{1}{2}(a-b)}{\sin \tfrac{1}{2}(a+b)}.$$

Hence,

$$\frac{\cos (S-A)}{\cos \tfrac{1}{2}(A-B) \cos \tfrac{1}{2} C} = \frac{2 \sin \tfrac{1}{2} a \cos \tfrac{1}{2} b}{\sin \tfrac{1}{2}(a+b}; \qquad (2)$$

but (p. 75, Ex. 31)

$$\cos (S-A) = \frac{\cos \tfrac{1}{2} b \cos \tfrac{1}{2} c}{\cos \tfrac{1}{2} a} \sin A = \sin \tfrac{1}{2} a \cos \tfrac{1}{2} c \frac{\sin B}{\sin \tfrac{1}{2} b}.$$

Substituting this value in (2), we have

$$\cos \tfrac{1}{2}(A - B) \sin \frac{c}{2} = \sin \tfrac{1}{2}(a + b) \sin \tfrac{1}{2} C. \qquad (\beta)$$

Similarly,

$$\sin \tfrac{1}{2}(A + B) \cos \frac{c}{2} = \cos \tfrac{1}{2}(a \mp b) \cos \tfrac{1}{2} C, \qquad (\gamma)$$

and

$$\sin \tfrac{1}{2}(A - B) \sin \frac{c}{2} = \sin \tfrac{1}{2}(a - b) \cos \tfrac{1}{2} C. \qquad (\delta)$$

65. It will be thus seen that when each of Napier's Analogies is added to or subtracted from unity, the results are easily transformed into Delambre's formulæ.

It is also to be noticed that if the equation

$$\cos \tfrac{1}{2}(A + B) \cos \tfrac{1}{2} c = \cos \tfrac{1}{2}(a + b) \sin \tfrac{1}{2} C$$

be applied to the co-lunar triangle of parts a, $\pi - b$, $\pi - c$, A, $\pi - B$, $\pi - C$, we get

$$\sin \tfrac{1}{2}(A - B) \sin \tfrac{1}{2} c = \sin \tfrac{1}{2}(a - b) \cos \tfrac{1}{2} C.$$

And by a similar method, the equation

$$\sin \tfrac{1}{2}(A + B) \cos \tfrac{1}{2} c = \cos \tfrac{1}{2}(a - b) \cos \tfrac{1}{2} C$$

gives

$$\cos \tfrac{1}{2}(A - B) \sin \tfrac{1}{2} c = \sin \tfrac{1}{2}(a + b) \sin \tfrac{1}{2} C.$$

Remark.—From the three preceding Articles it has been shown that Napier's four Analogies are reducible, by aid of the polar triangle, to *two independent theorems*, involving five parts of a spherical triangle; and that Gauss's four formulæ are, by aid of the colunar triangle, likewise reducible to two independent theorems.

66. In the solution of triangles, CASE III., the formulæ

$$\tan \tfrac{1}{2}(A+B) = \frac{\cos \tfrac{1}{2}(a-b)}{\cos \tfrac{1}{2}(a+b)} \cot \tfrac{1}{2} C,$$

and

$$\tan \tfrac{1}{2}(A-B) = \frac{\sin \tfrac{1}{2}(a-b)}{\sin \tfrac{1}{2}(a+b)} \cot \tfrac{1}{2} C,$$

enable us to calculate $\frac{1}{2}(A+B)$ and $\frac{1}{2}(A-B)$, from which we obtain A and B.

The remaining element c is determined from the equation

$$\cos \tfrac{1}{2}(A+B) \cos \tfrac{1}{2} c = \cos \tfrac{1}{2}(a+b) \sin \tfrac{1}{2} C;$$

or, as will be hereafter shown, it may be determined from the formula

$$\cos c = \cos a \cos b + \sin a \sin b \cos C,$$

without having previously determined A or B.

Examples.

(1). Given $a = 43° 18'$, $b = 19° 24'$, $C = 74° 22'$.

Firstly, determine A and B—

$$L \cos \tfrac{1}{2}(a-b) = L \cos 11° 57' = 9{\cdot}9904848.$$
$$L \cos \tfrac{1}{2}(a+b) = L \cos 31° 21' = 9{\cdot}9314605.$$
$$L \cot \tfrac{1}{2} C \quad = L \cot 37° 11' = 10{\cdot}1199969.$$
$$L \tan \tfrac{1}{2}(A+B) = 10{\cdot}1790212.$$

$$L \sin \tfrac{1}{2}(a-b) = 9{\cdot}3160921.$$
$$L \sin \tfrac{1}{2}(a+b) = 9{\cdot}7162243.$$
$$L \tan \tfrac{1}{2}(A-B) = 9{\cdot}7198647.$$

Therefore,

$$\tfrac{1}{2}(A+B) = 56° 29' 17'', \quad \text{and} \quad \tfrac{1}{2}(A-B) = 27° 41' 0\tfrac{1}{2}''.$$

To determine c—

$$L\cos\tfrac{1}{2}(a+b) = L\cos 31°\ 21' = 9{\cdot}9314605.$$
$$L\cos\tfrac{1}{2}(A+B) = L\cos 56°\ 29'\ 17'' = 9{\cdot}7420263.$$
$$L\sin\tfrac{1}{2}C = L\sin 37°\ 11' = 9{\cdot}7813010.$$

Hence

$$L\cos\tfrac{1}{2}c = 9{\cdot}9707352, \quad \text{or} \quad \tfrac{1}{2}c = 20°\ 48'\ 54\tfrac{1}{4}''.$$

Ans. $A = 84°\ 10'\ 17\tfrac{1}{2}''$, $B = 28°\ 48'\ 17\tfrac{1}{2}''$, $c = 41°\ 35'\ 48\tfrac{1}{2}''$.

(2). $a = 96°\ 24'\ 30''$, $b = 68°\ 27'\ 26''$, $C = 84°\ 46'\ 40''$.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 13°\ 58'\ 32'' = 9{\cdot}9869503.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 82°\ 25'\ 58'' = 9{\cdot}1195505.$$
$$L\cot\tfrac{1}{2}C = L\cot 42°\ 23'\ 20'' = 10{\cdot}0396387.$$
$$L\tan\tfrac{1}{2}(A+B) = 10{\cdot}9070385.$$
$$L\sin\tfrac{1}{2}(a-b) = 9{\cdot}3829313, \quad L\sin\tfrac{1}{2}(a+b) = 9{\cdot}9962011.$$
$$L\tan\tfrac{1}{2}(A-B) = 9{\cdot}4263689.$$
$$\tfrac{1}{2}(A+B) = 82°\ 56'\ 19\tfrac{3}{4}'', \quad \tfrac{1}{2}(A-B) = 14°\ 56'\ 4\ldots$$
$$L\cos\tfrac{1}{2}(A+B) = 9{\cdot}0896545.$$
$$L\sin\tfrac{1}{2}C = 9{\cdot}8287624.$$
$$L\cos\tfrac{1}{2}c = 9{\cdot}8586584, \quad \tfrac{1}{2}c = 43°\ 45'\ 48\tfrac{1}{2}''.$$



Ans. $A = 97°\ 53'\ 0\tfrac{1}{4}''$, $B = 67°\ 59'\ 39\tfrac{1}{4}''$, $c = 87°\ 31'\ 37''$.

(3). $a = 76°\ 24'\ 40''$, $b = 58°\ 18'\ 36''$, $C = 116°\ 30'\ 28''$.

$$L\cos\tfrac{1}{2}(a-b) = L\cos\ 9°\ 3'\ 2'' = 9{\cdot}9945590.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 67°\ 21'\ 38'' = 9{\cdot}5853827.$$
$$L\cot\tfrac{1}{2}C = L\cot 58°\ 15'\ 14'' = 9{\cdot}7914976.$$
$$L\tan\tfrac{1}{2}(A+B) = 10{\cdot}2006739.$$
$$L\sin\tfrac{1}{2}(a-b) = 9{\cdot}1967450, \quad L\sin\tfrac{1}{2}(a+b) = 9{\cdot}9651760.$$
$$L\tan\tfrac{1}{2}(A-B) = 9{\cdot}0230666.$$
$$\tfrac{1}{2}(A+B) = 57°\ 47'\ 23\tfrac{3}{4}'', \quad \tfrac{1}{2}(A-B) = 6°\ 1'\ 11\tfrac{1}{2}''.$$

$$L\cos\tfrac{1}{2}(A+B) = 9{\cdot}7267476$$
$$L\sin\tfrac{1}{2}C \quad = 9{\cdot}9296171.$$
$$\tfrac{1}{2}c = 52^\circ\ 6'\ 43\tfrac{1}{2}''.$$

Ans. $A = 63^\circ\ 48'\ 35\frac{1}{4}''$, $B = 51^\circ\ 46'\ 12\frac{1}{4}''$, $c = 104^\circ\ 13'\ 27''$.

(4). $a = 86^\circ\ 18'\ 40''$, $b = 45^\circ\ 36'\ 20''$, $C = 120^\circ\ 46'\ 30''$.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 20^\circ\ 21'\ 10'' = 9{\cdot}9720032.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 65^\circ\ 57'\ 30'' = 9{\cdot}6100219.$$
$$L\cot\tfrac{1}{2}C \quad = L\cot 60^\circ\ 23'\ 15'' = 9{\cdot}7546296.$$
$$\tfrac{1}{2}(A+B) = 52^\circ\ 36'\ 4\tfrac{3}{4}''.$$

$$L\sin\tfrac{1}{2}(a-b) = 9{\cdot}5413288, \quad L\sin\tfrac{1}{2}(a+b) = 9{\cdot}9605894.$$
$$\tfrac{1}{2}(A-B) = 12^\circ\ 12'\ 49''.$$
$$L\sin\tfrac{1}{2}C \quad = 9{\cdot}9392132.$$
$$L\cos\tfrac{1}{2}(A+B) = 9{\cdot}7834444.$$
$$\tfrac{1}{2}c = 54^\circ\ 19'\ 35\tfrac{3}{4}''.$$

Ans. $A = 64^\circ\ 48'\ 53\frac{3}{4}''$, $B = 40^\circ\ 23'\ 15\frac{3}{4}''$, $c = 108^\circ\ 39'\ 11\frac{1}{2}''$.

(5). $a = 88^\circ\ 24'$, $b = 56^\circ\ 48'$, $C = 128^\circ\ 16'$.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 15^\circ\ 48' = 9{\cdot}9832735.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 72^\circ\ 36' = 9{\cdot}4757304.$$
$$L\cot\tfrac{1}{2}C \quad = L\cot 64^\circ\ 8' = 9{\cdot}6856120.$$
$$\tfrac{1}{2}(A+B) = 57^\circ\ 20'\ 27\tfrac{1}{4}''.$$

$$L\sin\tfrac{1}{2}(a-b) = 9{\cdot}4350161, \quad L\sin\tfrac{1}{2}(a+b) = 9{\cdot}9796578.$$
$$\tfrac{1}{2}(A-B) = 7^\circ\ 52'\ 36\tfrac{1}{4}''.$$
$$L\cos\tfrac{1}{2}(A+B) = 9{\cdot}7321029.$$
$$L\sin\tfrac{1}{2}C \quad = 9{\cdot}9541517.$$
$$\tfrac{1}{2}c = 60^\circ\ 5'\ 26''.$$

Ans. $A = 65^\circ\ 13'\ 3\frac{1}{2}''$, $B = 49^\circ\ 27'\ 51''$, $c = 120^\circ\ 10'\ 52''$.

(6). $a = b = 32° 23' 57''$, $C = 66° 49' 17''$.

(*Science and Art Exam. Papers.*)

$L \cos \frac{1}{2}(a - b) = 10.$

$L \cos \frac{1}{2}(a + b) = L \cos 32° 23' 57'' = 9{\cdot}9265152.$

$L \cot \frac{1}{2} C = L \cot 33° 24' 38\frac{1}{2}'' = 10{\cdot}1806890.$

$\frac{1}{2}(A + B) = 60° 53' 2''.$

$L \sin a = 9{\cdot}7290144.$

$L \sin \frac{1}{2} C = 9{\cdot}7408650.$

$\frac{1}{2} c = 17° 9' 35\frac{1}{2}''.$

Ans. $A = B = 60° 53' 2''$, $c = 34° 19' 11''$.

67. Case IV.—*Having given two angles and the adjacent side* (A, B, c).

By *Napier's Analogies:*

$$\tan \tfrac{1}{2}(a + b) = \frac{\cos \frac{1}{2}(A - B)}{\cos \frac{1}{2}(A + B)} \tan \tfrac{1}{2} c,$$

and

$$\tan \tfrac{1}{2}(a - b) = \frac{\sin \frac{1}{2}(A - B)}{\sin \frac{1}{2}(A + B)} \tan \tfrac{1}{2} c;$$

from which we may determine $\frac{1}{2}(a + b)$ and $\frac{1}{2}(a - b)$, and thence a and b.

The remaining part C is determined by the equation

$$\cos \tfrac{1}{2}(a + b) \sin \tfrac{1}{2} C = \cos \tfrac{1}{2}(A + B) \cos \tfrac{1}{2} c.$$

It is evident that the value of C thus found is ambiguous,* inasmuch as the angle is obtained from the *sine;* and the values are, therefore, C and $180 - C$. The equation has, however, been selected as an example, to show that an

* By using formula γ (or δ), p. 113, ambiguity is avoided.

ambiguity may sometimes be got rid of by attending to the relations the parts of a triangle bear to each other; and in the following numerical examples the ambiguity is removed by remembering that the greater side is opposite to the greater angle.

Examples.

(1). $A = 68° 40'$, $B = 56° 20'$, $c = 84° 30'$.

$$L\cos\tfrac{1}{2}(A-B) = L\cos\ 6° 10' = 9\ 9974797.$$
$$L\cos\tfrac{1}{2}(A+B) = L\cos 62° 30' = 9{\cdot}6644056.$$
$$L\tan\tfrac{1}{2}c \quad = L\tan 42° 15' = 9{\cdot}9582465.$$
$$\tfrac{1}{2}(a+b) = 62° 55' 9''.$$
$$L\sin\tfrac{1}{2}(A-B) = 9{\cdot}0310890, \quad L\sin\tfrac{1}{2}(A+B) = 9{\cdot}9479289.$$
$$\tfrac{1}{2}(a-b) = 6° 16' 39''.$$
$$L\cos\tfrac{1}{2}(a+b) = 9{\cdot}6582472.$$
$$L\cos\tfrac{1}{2}c \quad = 9{\cdot}8693597.$$
$$\tfrac{1}{2}C = 48° 39' 31\tfrac{3}{4}''.$$

Ans. $a = 69° 11' 48''$, $b = 56° 38' 30''$, $C = 97° 19' 3\tfrac{1}{2}''$.

(2). $A = 31° 34' 26''$, $B = 30° 28' 12''$, $c = 70° 2' 3''$.

$$L\cos\tfrac{1}{2}(A-B) = L\cos\ 0° 33'\ 7'' = 9{\cdot}\ 9999799.$$
$$L\cos\tfrac{1}{2}(A+B) = L\cos 31°\ 1'\ 19'' = 9{\cdot}\ 9329656.$$
$$L\tan\tfrac{1}{2}c \quad = L\tan 35°\ 1'\ 1\tfrac{1}{2}'' = 9{\cdot}\ 8455023.$$
$$\tfrac{1}{2}(a+b) = 39° 16' 4\tfrac{1}{4}''.$$
$$L\sin\tfrac{1}{2}(A-B) = 7{\cdot}9837459. \quad L\sin\tfrac{1}{2}(A+B) = 9{\cdot}7121160.$$
$$\tfrac{1}{2}(a-b) = 0° 45' 1''.$$
$$L\cos\tfrac{1}{2}(a+b) = 9{\cdot}8888507.$$
$$L\cos\tfrac{1}{2}c \quad = 9{\cdot}9132738.$$
$$\tfrac{1}{2}C = 65° 1' 55''.$$

Ans. $a = 40° 1' 5\tfrac{1}{4}''$, $b = 38° 31' 3\tfrac{1}{4}''$, $C = 130° 3' 50''$.

(3). $A = 130° 5' 22{\cdot}41''$, $B = 32° 26' 6{\cdot}41''$, $c = 51° 6' 11{\cdot}6''$.

(*Science Siz. Exam. Papers.*)

$$L \cos \tfrac{1}{2}(A - B) = L \cos 48° 49' 38'' = 9{\cdot}8184449.$$
$$L \cos \tfrac{1}{2}(A + B) = L \cos 81° 15' 44{\cdot}41'' = 9{\cdot}1815879.$$
$$L \tan \tfrac{1}{2} c = L \tan 25° 33' 5{\cdot}8'' = 9{\cdot}6795022.$$

$$\tfrac{1}{2}(a + b) = 64° 14' 7''.$$

$$L \sin \tfrac{1}{2}(A - B) = 9{\cdot}8766380, \quad L \sin \tfrac{1}{2}(A + B) = 9{\cdot}9949301.$$

$$\tfrac{1}{2}(a - b) = 20° 0' 22''.$$

$$L \cos \tfrac{1}{2}(a + b) = 9{\cdot}6381663.$$
$$L \cos \tfrac{1}{2} c = 9{\cdot}9552944.$$

$$\tfrac{1}{2} C = 18° 22' 43''.$$

Ans. $a = 84° 14' 29''$, $b = 44° 13' 45''$, $C = 36° 45' 26''$.

(4). $A = 96° 46' 30''$, $B = 84° 30' 20''$, $c = 126° 46'$.

$$L \cos \tfrac{1}{2}(A - B) = L \cos 6° 8' 5'' = 9{\cdot}9975058.$$
$$*L \text{ cos supplement of } \tfrac{1}{2}(A + B) = L \cos 89° 21' 35'' = 8{\cdot}0482011.$$
$$L \tan \tfrac{1}{2} c = L \tan 63° 23'' = 10{\cdot}3000526.$$

Supplement of $\tfrac{1}{2}(a + b) = 89° 40' 38''$.

$$L \sin \tfrac{1}{2}(A - B) = 9{\cdot}0288420, \quad L \sin \tfrac{1}{2}(A + B) = 9{\cdot}9999729.$$

$$\tfrac{1}{2}(a - b) = 12° 2' 20''.$$

$$L \text{ cos supplement of } \tfrac{1}{2}(a + b) = 7{\cdot}7506455, \quad \cos \tfrac{1}{2} c = 9{\cdot}6512966.$$

$$\tfrac{1}{2} C = 62° 44' 6\tfrac{5}{8}''.$$

Ans. $a = 102° 21' 42''$, $b = 78° 17' 2''$, $C = 125° 28' 13\tfrac{1}{4}''$.

* When $\tfrac{1}{2}(A + B)$ exceeds a quadrant, by substituting its supplement in the equation, a value is obtained for the supplement of $\tfrac{1}{2}(a + b)$. See *Napier's Analogies*, and *Gauss's Formulæ*, from which it also appears that $\tfrac{1}{2}(A + B)$ and $\tfrac{1}{2}(a + b)$ are of the same affection.

(5). $A = 84° 30' 20''$, $B = 76° 20' 40''$, $c = 130° 46$.

$$L \cos \tfrac{1}{2}(A - B) = L \cos 4° 4' 50'' = 9{\cdot}9988977.$$
$$L \cos \tfrac{1}{2}(A + B) = L \cos 80° 25' 30'' = 9{\cdot}2209927.$$
$$L \tan \tfrac{1}{2} c = L \tan 65° 23' = 10{\cdot}3389566.$$

$$\tfrac{1}{2}(a + b) = 85° 37' 50\tfrac{1}{4}''.$$

$$L \sin \tfrac{1}{2}(A - B) = 8{\cdot}8522289, \quad L \sin \tfrac{1}{2}(A + B) = 9{\cdot}9939071.$$
$$\tfrac{1}{2}(a - b) = 8° 57' 2''.$$

$$L \cos \tfrac{1}{2}(a + b) = 8{\cdot}8818753.$$
$$L \cos \tfrac{1}{2} c = 9{\cdot}6196622.$$

$$\tfrac{1}{2} C = 65° 25' 46\tfrac{3}{4}''.$$

Ans. $a = 94° 34' 52\tfrac{1}{4}''$, $b = 76° 40' 48\tfrac{1}{4}''$, $C = 130° 51' 33\tfrac{1}{2}''$.

68. Case V.—*Having given two sides and the angle opposite either* (a, b, A).

The angle B is found from the formula

$$\sin B = \frac{\sin b}{\sin a} \sin A.$$

For the angle C,

$$\tan \tfrac{1}{2} C = \frac{\cos \tfrac{1}{2}(a - b)}{\cos \tfrac{1}{2}(a + b)} \cot \tfrac{1}{2}(A + B).$$

For the side c,

$$\tan \tfrac{1}{2} c = \frac{\cos \tfrac{1}{2}(A + B)}{\cos \tfrac{1}{2}(A - B)} \tan \tfrac{1}{2}(a + b).$$

Since B has been found from its sine, it may have two values, viz. B and $180 - B$, and the triangle in the general case will admit of two solutions.

This is known as the *ambiguous case*, a full discussion of which is given in Arts. 70–75.

The following numerical examples will serve to elucidate how two different triangles may be constructed when three parts are given :—

Examples.

1). $a = 42° 45'$, $b = 47° 15'$, $A = 56° 30'$.

$$L \sin a = L \sin 42° 45' = 9{\cdot}8317423.$$

$$L \sin b = L \sin 47° 15' = 9{\cdot}8658868.$$

$$L \sin A = L \sin 56° 30' = 9{\cdot}9211066.$$

$$L \sin B = 9{\cdot}9552511.$$

Hence

$$B = 64° 26' 4'', \quad \text{or} \quad 115° 33'' 56''.$$

Firstly—take the value of B less than a right angle.

$$L \cos \tfrac{1}{2}(b + a) = L \cos 45° 0' = 9{\cdot}8494850.$$

$$L \cos \tfrac{1}{2}(b - a) = L \cos 2° 15' = 9{\cdot}9996650;$$

and

$$L \cot \tfrac{1}{2}(B + A) = L \cot 60° 28' 2'' = 9{\cdot}7532216.$$

Hence

$$L \tan \tfrac{1}{2} C = 9{\cdot}9034016;$$

or

$$\tfrac{1}{2} C = 38° 40' 47\tfrac{1}{2}''.$$

$$L \cos \tfrac{1}{2}(B + A) = L \cos 60° 28' 2'' = 9{\cdot}6927772.$$

$$L \cos \tfrac{1}{2}(B - A) = L \cos 3° 58' 2'' = 9{\cdot}9989581.$$

$$L \tan \tfrac{1}{2}(b + a) = L \tan 45° 0' 0'' = 10{\cdot}0000000.$$

$$\tfrac{1}{2} c = 26° 17' 39''.$$

Secondly—take the value of B greater than a right angle. Then

$$L \cot \tfrac{1}{2}(B + A) = L \cot 86° 1' 58'' = 8{\cdot}8410586;$$

and by aid of the logarithms in the first case we have

$$\tfrac{1}{2} C = 5° 35' 50\tfrac{1}{4}''.$$

Again,

$$L\cos\tfrac{1}{2}(B+A) = L\cos 86^\circ\ 1'\ 58'' = 8{\cdot}8400167.$$

$$L\cos\tfrac{1}{2}(B-A) = L\cos 29^\circ\ 31'\ 58'' = 9{\cdot}9395561.$$

Hence

$$\tfrac{1}{2}c = 4^\circ\ 32'\ 47\tfrac{1}{8}''.$$

Ans.—

$$B = 64^\circ\ 25'\ 4'',\quad C = 77^\circ\ 21'\ 35'',\quad c = 52^\circ\ 35'\ 18'';$$

or

$$B = 115^\circ\ 33'\ 56'',\quad C = 11^\circ\ 11'\ 40\tfrac{1}{2}'',\quad c = 9^\circ\ 5'\ 34\tfrac{1}{4}''.$$

(2). $a = 62^\circ\ 15'\ 24'',\quad b = 103^\circ\ 18'\ 47'',\quad A = 53^\circ\ 42'\ 38''.$

$$L\sin a = L\sin 62^\circ\ 15'\ 24'' = 9{\cdot}9469638.$$

$$L\sin b = L\sin 76^\circ\ 41'\ 13'' = 9{\cdot}9881693.$$

$$L\sin A = L\sin 53^\circ\ 42'\ 38'' = 9{\cdot}9063552.$$

$$B = 62^\circ\ 24'\ 24\tfrac{3}{8}'',\quad \text{or}\quad 117^\circ\ 35'\ 35\tfrac{1}{8}''.$$

Firstly—

$$L\cos\tfrac{1}{2}(b+a) = L\cos 82^\circ\ 47'\ 5\tfrac{1}{2}'' = 9{\cdot}0989736.$$

$$L\cos\tfrac{1}{2}(b-a) = L\cos 20^\circ\ 31'\ 41\tfrac{1}{2}'' = 9{\cdot}9715077.$$

$$L\cot\tfrac{1}{2}(B+A) = L\cot 58^\circ\ 3'\ 31\tfrac{3}{8}'' = 9{\cdot}7947983.$$

$$\tfrac{1}{2}C = 77^\circ\ 51'\ 35\tfrac{3}{8}''.$$

$$L\cos\tfrac{1}{2}(B+A) = L\cos 58^\circ\ 3'\ 31\tfrac{3}{8}'' = 9{\cdot}7234966.$$

$$L\cos\tfrac{1}{2}(B-A) = L\cos 4^\circ\ 20'\ 53\tfrac{3}{8}'' = 9{\cdot}9987482.$$

$$L\tan\tfrac{1}{2}(b+a) = L\tan 82^\circ\ 47'\ 5\tfrac{1}{2}'' = 10{\cdot}8975738.$$

$$\tfrac{1}{2}c = 76^\circ\ 34'\ 47\tfrac{3}{4}''.$$

Secondly—

$$L\ \cot\tfrac{1}{2}(B+A) = L\cot 85^\circ\ 39'\ 6\tfrac{3}{8}'' = 8{\cdot}8810181.$$

$$\tfrac{1}{2}C = 29^\circ\ 33'\ 5\tfrac{1}{8}''.$$

$$L\cos\tfrac{1}{2}(B+A) = L\cos 85^\circ\ 39'\ 6\tfrac{3}{8}'' = 8{\cdot}8797663.$$

$$L\cos\tfrac{1}{2}(B-A) = L\cos 31^\circ\ 56'\ 28\tfrac{3}{4}'' = 9{\cdot}9286983.$$

$$\tfrac{1}{2}c = 35^\circ\ 12'\ 43''.$$

Ans.—

$$B = 62^\circ\ 24'\ 24\tfrac{4}{8}'',\quad C = 155^\circ\ 43'\ 11\tfrac{1}{8}'',\quad c = 153^\circ\ 9'\ 35\tfrac{1}{2}'';$$

or

$$B = 117^\circ\ 35'\ 35\tfrac{1}{8}'',\quad C = 59^\circ\ 6'\ 10\tfrac{3}{4}'',\quad c = 70^\circ\ 25'\ 26''.$$

(3). $a = 52° 45' 20''$, $b = 71° 12' 40''$, $A = 46° 22' 10''$.

$$B = 59° 24' 15\tfrac{3}{4}'', \text{ or } 120° 35' 44\tfrac{1}{4}''.$$

Firstly—

$$L\cos\tfrac{1}{2}(b+a) = L\cos 61° 59' = 9{\cdot}6718468.$$
$$L\cos\tfrac{1}{2}(b-a) = L\cos 9° 13' 40'' = 9{\cdot}9943430.$$
$$L\cot\tfrac{1}{2}(B+A) = L\cot 52° 53' 12\tfrac{7}{8}'' = 9{\cdot}8788970.$$
$$\tfrac{1}{2}C = 57° 49' 57\tfrac{3}{4}''.$$
$$L\cos\tfrac{1}{2}(B+A) = L\cos 52° 53' 12\tfrac{7}{8}'' = 9{\cdot}7805983.$$
$$L\cos\tfrac{1}{2}(B-A) = L\cos 6° 31' 2\tfrac{7}{8}'' = 9{\cdot}9971842.$$
$$L\tan\tfrac{1}{2}(b+a) = L\tan 61° 59' = 10{\cdot}2740209.$$
$$\tfrac{1}{2}c = 48° 46' 39\tfrac{2}{3}''.$$

Secondly—

$$L\cot\tfrac{1}{2}(B+A) = L\cot 83° 28' 57\tfrac{1}{8}'' = 9{\cdot}0578349.$$
$$\tfrac{1}{2}C = 13° 29' 57\tfrac{3}{8}''.$$
$$L\cos\tfrac{1}{2}(B+A) = L\cos 83° 28' 57\tfrac{1}{8}'' = 9{\cdot}0550190.$$
$$L\cos\tfrac{1}{2}(B-A) = L\cos 37° 6' 47\tfrac{1}{8}'' = 9{\cdot}9017013.$$
$$\tfrac{1}{2}c = 14° 58' 35\tfrac{1}{4}''.$$

Ans.—

$$B = 59° 24' 15\tfrac{3}{4}'', \quad C = 115° 39' 55\tfrac{1}{2}'', \quad c = 97° 33' 18\tfrac{4}{3}'',$$

or

$$B = 120° 35' 44\tfrac{1}{4}'', \quad C = 26° 59' 55\tfrac{1}{3}'', \quad c = 29° 57' 10\tfrac{1}{2}''.$$

(4). $a = 48° 45' 40''$, $b = 67° 12' 20''$, $A = 42° 20' 30''$.

$$B = 55° 39' 57'', \text{ or } 124° 20' 3''.$$

Firstly—

$$L\cos\tfrac{1}{2}(b+a) = L\cos 57° 59' 0'' = 9{\cdot}7244118.$$
$$L\cos\tfrac{1}{2}(b-a) = L\cos 9° 13' 20'' = 9{\cdot}9943498.$$
$$L\cot\tfrac{1}{2}(B+A) = L\cot 49° 0' 13\tfrac{1}{2}'' = 9{\cdot}9391057.$$
$$\tfrac{1}{2}C = 58° 17' 9''.$$
$$L\cos\tfrac{1}{2}(B+A) = L\cos 49° 0' 13\tfrac{1}{2}'' = 9{\cdot}8169102.$$
$$L\cos\tfrac{1}{2}(B-A) = L\cos 6° 39' 43\tfrac{1}{2}'' = 9{\cdot}9970576.$$
$$L\tan\tfrac{1}{2}(b+a) = L\tan 57° 59' 0'' = 10{\cdot}2029297.$$
$$\frac{c}{2} = 46° 34' 4\tfrac{1}{3}''.$$

Secondly—

$$L \cot \tfrac{1}{2}(B + A) = L \cot 83° 20' 16\tfrac{1}{2}'' = 9{\cdot}0674506.$$

$$\tfrac{1}{2} C = 12° 16' 7\tfrac{1}{2}''.$$

$$L \cos \tfrac{1}{2}(B + A) = L \cos 83° 20' 16\tfrac{1}{2}'' = 9{\cdot}0645081.$$

$$L \cos \tfrac{1}{2}(B - A) = L \cos 40° 59' 46\tfrac{1}{2}'' = 9{\cdot}8778046.$$

$$\tfrac{1}{2} c = 13° 48' 40''.$$

Ans.—

$$B = 55° 39' 57'', \quad C = 116° 34' 18'', \quad c = 93° 8' 9\tfrac{3}{5}'';$$

or

$$B = 124° 20' 3'', \quad C = 24° 32' 15'', \quad c = 27° 37' 20''.$$

(5). $a = 46° 20' 45''$, $b = 65° 18' 15''$, $A = 40° 10' 30''$.

$$B = 54° 6' 19'', \text{ or } 125° 53' 41''.$$

Firstly—

$$L \cos \tfrac{1}{2}(b + a) = L \cos 55° 49' 30'' = 9{\cdot}7495218.$$

$$L \cos \tfrac{1}{2}(b - a) = L \cos 9° 28' 45'' = 9{\cdot}9940291.$$

$$L \cot \tfrac{1}{2}(B + A) = L \cot 47° 8' 24\tfrac{1}{2}'' = 9{\cdot}9675258.$$

$$\tfrac{1}{2} C = 58° 27' 43''.$$

$$L \cos \tfrac{1}{2}(B + A) = L \cos 47° 8' 24\tfrac{1}{2}'' = 9{\cdot}8326414.$$

$$L \cos \tfrac{1}{2}(B - A) = L \cos 6° 57' 54\tfrac{1}{2}'' = 9{\cdot}9967831.$$

$$L \tan \tfrac{1}{2}(b + a) = L \tan 55° 49' 30'' = 10{\cdot}1681548.$$

$$\tfrac{1}{2} c = 45° 15' 53''.$$

Secondly—

$$L \cot \tfrac{1}{2}(B + A) = L \cot 83° 2' 5\tfrac{1}{2}'' = 9{\cdot}0869538.$$

$$\tfrac{1}{2} C = 12° 6' 26\tfrac{2}{3}''.$$

$$L \cos \tfrac{1}{2}(B + A) = L \cos 83° 2' 5\tfrac{1}{2}'' = 9{\cdot}0837368.$$

$$L \cos \tfrac{1}{2}(B - A) = L \cos 42° 51' 35\tfrac{1}{2}'' = 9{\cdot}8651156.$$

$$\tfrac{1}{2} c = 13° 41' 37''.$$

Ans.—

$$B = 54° 6' 19'', \quad C = 116° 55' 26'', \quad c = 90° 31' 46'.$$

$$B = 125° 53' 41'', \quad C = 24° 12' 53\tfrac{1}{3}'', \quad c = 27° 23' 14''.$$

69. CASE VI.—*Having given two angles and a side opposite one of them* (A, B, a).

This Case reduces, by aid of the polar triangle, to the preceding, and the direct solution may be therefore obtained by means of formulæ supplemental to those employed in Art. 68, viz.,

$$\sin b = \frac{\sin B}{\sin A} \sin a,$$

$$\tan \tfrac{1}{2} C = \frac{\cos \tfrac{1}{2}(a-b)}{\cos \tfrac{1}{2}(a+b)} \cot \tfrac{1}{2}(A+B),$$

$$\tan \tfrac{1}{2} c = \frac{\cos \tfrac{1}{2}(A+B)}{\cos \tfrac{1}{2}(A-B)} \tan \tfrac{1}{2}(a+b).$$

The same ambiguities will thus present themselves.

Examples.

(1). $a = 59° 28' 27''$, $A = 66° 7' 20''$, $B = 52° 50' 20''$.

$b = 48° 39' 16''$, or $131° 20' 44''$.

Firstly—take the value of b less than a right angle.

$$L \cos \tfrac{1}{2}(a-b) = L \cos 5° 24' 35\tfrac{1}{2}'' = 9{\cdot}9980612.$$
$$L \cos \tfrac{1}{2}(a+b) = L \cos 54° 3' 51\tfrac{1}{2}'' = 9{\cdot}7685470.$$
$$L \cot \tfrac{1}{2}(A+B) = L \cot 59° 28' 50'' = 9{\cdot}7704854.$$

$$\tfrac{1}{2} C = 45.$$

$$L \cos \tfrac{1}{2}(A+B) = L \cos 59° 28' 50'' = 9{\cdot}7057190.$$
$$L \cos \tfrac{1}{2}(A-B) = L \cos 6° 38' 30'' = 9{\cdot}9970756.$$
$$L \tan \tfrac{1}{2}(a+b) = L \tan 54° 3' 51\tfrac{1}{2}'' = 10{\cdot}1397643.$$

$$\tfrac{1}{2} c = 35° 11' 50\tfrac{3}{4}''.$$

The second value of b is inadmissible (see foot note, p. 119), and therefore there is only one solution.

Ans.—

$$b = 48^\circ\ 39'\ 16'', \quad C = 90^\circ, \quad c = 70^\circ\ 23'\ 41\tfrac{1}{2}''.$$

(2). $a = 53^\circ\ 18'\ 20$, $A = 79^\circ\ 30'\ 45''$, $B = 46^\circ\ 15'\ 15''$.

$$b = 36^\circ\ 5'\ 34\tfrac{3}{4}'', \text{ or } 143^\circ\ 54'\ 25\tfrac{1}{4}''.$$

Firstly—

$$L\cos\tfrac{1}{2}(a-b) = L\cos 8^\circ\ 36'\ 22\tfrac{1}{2}'' = 9{\cdot}9950822.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 44^\circ\ 41'\ 57\tfrac{1}{4}'' = 9{\cdot}8517528.$$
$$L\cot\tfrac{1}{2}(A+B) = L\cot 62^\circ\ 53' \quad = 9{\cdot}7093488.$$
$$\tfrac{1}{2}C = 35^\circ\ 27'\ 47\tfrac{1}{2}''.$$

$$L\cos\tfrac{1}{2}(A+B) = L\cos 62^\circ\ 53' \quad = 9{\cdot}6587780.$$
$$L\cos\tfrac{1}{2}(A-B) = L\cos 16^\circ\ 37'\ 45'' \quad = 9{\cdot}9814457.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 44^\circ\ 41'\ 57\tfrac{1}{4}'' = 9{\cdot}9954404.$$
$$\tfrac{1}{2}c = 25^\circ\ 12'\ 28\tfrac{1}{2}''.$$

[No ambiguity, as in Ex. 1.]

Ans.—

$$b = 36^\circ\ 5'\ 34\tfrac{3}{4}'', \quad C = 70^\circ\ 55'\ 35'', \quad c = 50^\circ\ 24'\ 57'.$$

(3). $a = 46^\circ\ 45'\ 30''$, $A = 73^\circ\ 11'\ 18''$, $B = 61^\circ\ 18'\ 12''$;

therefore $b = 41^\circ\ 52'\ 34\tfrac{3}{4}''$, or $138^\circ\ 7'\ 25\tfrac{1}{4}''$.

Firstly—

$$L\cos\tfrac{1}{2}(a-b) = L\cos 2^\circ\ 26'\ 23\tfrac{1}{4}'' = 9{\cdot}9996061.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 44^\circ\ 19'\ 6\tfrac{3}{4}'' = 9{\cdot}8545894.$$
$$L\cot\tfrac{1}{2}(A+B) = L\cot 67^\circ\ 14'\ 45'' = 9{\cdot}6226494.$$
$$\tfrac{1}{2}C = 30^\circ\ 21'\ 23\tfrac{1}{4}''.$$

$$L\cos\tfrac{1}{2}(A+B) = L\cos 67^\circ\ 14'\ 45'' = 9{\cdot}5874618.$$
$$L\cos\tfrac{1}{2}(A-B) = L\cos 5^\circ\ 56'\ 33'' = 9{\cdot}9976600.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 44^\circ\ 19'\ 6\tfrac{3}{4}'' = 9{\cdot}9896683.$$
$$\tfrac{1}{2}c = 20^\circ\ 47'\ 32''.$$

[Only one solution.]

Ans.—

$$b = 41^\circ\ 52'\ 34\tfrac{3}{4}'', \quad C = 60^\circ\ 42'\ 46\tfrac{1}{2}'', \quad c = 41^\circ\ 35'\ 4''.$$

(4). $a = 42° 15' 20''$, $A = 46° 30' 40''$, $B = 36° 20' 20''$.

$b = 33° 18' 47\frac{1}{2}''$, or $146° 41' 12\frac{1}{2}''$.

Firstly—

$L \cos \frac{1}{2}(a - b) = L \cos 4° 28' 16\frac{1}{4}'' = 9{\cdot}9986763.$

$L \cos \frac{1}{2}(a + b) = L \cos 37° 47' 3\frac{3}{4}'' = 9{\cdot}8978042.$

$L \cot \frac{1}{2}(A + B) = L \cot 41° 25' 30'' = 10{\cdot}0543373.$

$\frac{1}{2} C = 55° 1' 37\frac{1}{3}''.$

$L \cos \frac{1}{2}(A + B) = L \cos 41° 25' 30'' = 9{\cdot}8749585.$

$L \cos \frac{1}{2}(A - B) = L \cos 5° 5' 10'' = 9{\cdot}9982866.$

$L \tan \frac{1}{2}(a + b) = L \tan 37° 47' 3\frac{3}{4}'' = 9{\cdot}8894377.$

$\frac{1}{2} c = 30° 16' 3''.$

[Only one solution.]

Ans.—

$b = 33° 18' 47\frac{1}{2}''$, $C = 110° 3' 14\frac{2}{3}''$, $c = 60° 32' 6''$.

(5). $a = 34° 30'$, $A = 61° 29' 30''$, $B = 24° 30' 30''$.

$b = 15° 30' 30\frac{1}{2}''$, or $164° 29' 29\frac{1}{2}''$.

$L \cos \frac{1}{2}(a - b) = L \cos 9° 29' 44\frac{3}{4}'' = 9{\cdot}9940081.$

$L \cos \frac{1}{2}(a + b) = L \cos 25° 0' 15\frac{1}{4}'' = 9{\cdot}9572609.$

$L \cot \frac{1}{2}(A + B) = L \cot 43° = 10{\cdot}0303441.$

$\frac{1}{2} C = 49° 24' 29\frac{1}{4}''.$

$L \cos \frac{1}{2}(A + B) = L \cos 43° = 9{\cdot}8641275.$

$L \cos \frac{1}{2}(A - B) = L \cos 18° 29' 30'' = 9{\cdot}9769777.$

$L \tan \frac{1}{2}(a + b) = L \tan 25° 0' 15\frac{1}{4}'' = 9{\cdot}6687563.$

$\frac{1}{2} c = 19° 46' 56''.$

[Only one solution.]

Ans.—

$b = 15° 30' 30\frac{1}{2}''$, $C = 98° 48' 58\frac{1}{2}''$, $c = 39° 33' 52''$.

(6). $a = 42° 15' 20''$, $A = 36° 20' 20''$, $B = 46° 30' 40''$;

therefore $b = 55° 25' 2\frac{1}{2}''$, or $124° 34' 57\frac{1}{2}''$.

Firstly—take the value of b less than a right angle.

$$L\cos\tfrac{1}{2}(a-b) = L\cos\ 6°\ 34'\ 51\tfrac{1}{4}'' = 9{\cdot}9971289.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 48°\ 50'\ 11\tfrac{1}{4}'' = 9{\cdot}8183648.$$
$$L\cot\tfrac{1}{2}(A+B) = L\cot 41°\ 25'\ 30'' = 10{\cdot}0543373.$$
$$\tfrac{1}{2}C = 59°\ 41'\ 13\tfrac{3}{4}''.$$
$$L\cos\tfrac{1}{2}(A+B) = L\cos 41°\ 25'\ 30'' = 9{\cdot}8749585.$$
$$L\cos\tfrac{1}{2}(A-B) = L\cos\ 5°\ 5'\ 10'' = 9{\cdot}9982866.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 48°\ 50'\ 11\tfrac{1}{4}'' = 10{\cdot}0583343.$$
$$\tfrac{1}{2}c = 40°\ 43'\ 43\tfrac{1}{8}''.$$

Secondly—take the value of b greater than a right angle.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 41°\ 9'\ 48\tfrac{3}{4}'' = 9{\cdot}8766992.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 83°\ 25'\ 8\tfrac{3}{4}'' = 9{\cdot}0592074.$$
$$\tfrac{1}{2}C = 82°\ 20'\ 57\tfrac{1}{2}''.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 83°\ 25'\ 8\tfrac{3}{4}'' = 10{\cdot}9379216.$$
$$\tfrac{1}{2}c = 81°\ 17'\ 13\tfrac{1}{2}''.$$

Ans.—$b = 55°\ 25'\ 2\tfrac{1}{2}''$, $C = 119°\ 22'\ 27\tfrac{1}{2}''$, $c = 81°\ 27'\ 26\tfrac{1}{4}''$;
or $b = 124°\ 34'\ 57\tfrac{1}{2}''$, $C = 164°\ 41'\ 55''$, $c = 162°\ 34'\ 27''$.

(7). $a = 59°\ 28'\ 27''$, $A = 52°\ 50'\ 20''$, $B = 66°\ 7'\ 20''$;
therefore $b = 81°\ 15'\ 15''$, or $98°\ 44'\ 45''$.

Firstly—take the value of b less than a right angle.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 10°\ 53'\ 24'' = 9{\cdot}9921078.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 70°\ 21'\ 51'' = 9{\cdot}5263918.$$
$$L\cot\tfrac{1}{2}(A+B) = L\cot 59°\ 28'\ 50'' = 9{\cdot}7704854.$$
$$\tfrac{1}{2}C = 59°\ 51'\ 54''.$$
$$L\cos\tfrac{1}{2}(A+B) = L\cos 59°\ 28'\ 50'' = 9{\cdot}7057190.$$
$$L\cos\tfrac{1}{2}(A-B) = L\cos\ 6°\ 38'\ 30'' = 9{\cdot}9970756.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 70°\ 21'\ 51'' = 10{\cdot}4475887.$$
$$\tfrac{1}{2}c = 55°\ 5'\ 25\tfrac{1}{4}''.$$

Secondly—take the value of b greater than a right angle.

$$L\cos\tfrac{1}{2}(a-b) = L\cos 19°\ 38'\ 9'' = 9{\cdot}9739306.$$
$$L\cos\tfrac{1}{2}(a+b) = L\cos 79°\ 6'\ 36'' = 9{\cdot}2762872.$$
$$\tfrac{1}{2}C = 71°\ 12'\ 29\tfrac{1}{2}''.$$
$$L\tan\tfrac{1}{2}(a+b) = L\tan 79°\ 6'\ 36'' = 10{\cdot}7158207.$$
$$\tfrac{1}{2}c = 69°\ 22'\ 43''.$$

Ans.—$b = 81°\ 15'\ 15''$, $C = 119°\ 43'\ 48''$, $c = 110°\ 10'\ 50\tfrac{1}{2}''$.
or $b = 98°\ 44'\ 45''$, $C = 142°\ 24'\ 55''$, $c = 138°\ 45'\ 26''$.

70. **Discussion of the Ambiguous Case.**—Before entering upon the general discussion of the ambiguous case of spherical triangles, it will be necessary for the student to be familiar with the analogous ambiguity *in plano*, a complete discussion of which is generally given in works on Plane Trigonometry.

The reasoning is somewhat more complex in the case of spherical triangles, as there are nine distinct cases to be considered, in any of which an ambiguity in the construction of the triangle may present itself. It has been seen in the examples of Arts. 68 and 69, that from the given parts two real solutions were obtained, from which we infer that there can be constructed two distinct triangles satisfying the given conditions.

We shall now examine in detail the variety of cases which may arise when '*two sides and the angle opposite one of them are given*,' leaving the supplemental case '*having given two angles and the side opposite one of them*' to be reasoned out in an analogous manner.

71. *Let the given parts be denoted by A, a, b.*

The discussion resolves itself thus into three groups, each containing three separate cases:—

Case I.—When $A = \frac{1}{2}\pi$ and b is equal to, less than, or greater than $\frac{1}{2}\pi$;

Case II.—When $A < \frac{1}{2}\pi$ and b is equal to, less than, or greater than $\frac{1}{2}\pi$;

Case III.—When $A > \frac{1}{2}\pi$ and b is equal to, less than, or greater than $\frac{1}{2}\pi$;

the solution in any case being impossible, evanescent, unique, ambiguous, or indeterminate, according to the value of a.

72. Case I.—$A = \frac{1}{2}\pi$.

For the consideration of the three cases in Art. 71, take any great circle AC (fig. 29) equal to the given side b, and at right angles to the great circle ABX.




Fig. 29.

Firstly.—Let $b = \frac{1}{2}\pi$; and with C as pole, and a radius equal to a, describe a circle. Now if a be less or greater than a quadrant, this circle being concentric with ABX, the triangle is *impossible*. If a be a quadrant, the circles

are coincident, and the triangle is *indeterminate* with the given parts A, a, b, each quadrants.

Secondly.—Let $b < \frac{1}{2}\pi$. With C as pole (fig. 29), and a radius a, describe a circle. Since AC and $A'C$ are unequal, this circle may be situated relatively to ABX in five ways—(1) When the circumference falls wholly within ABX; (2) When it touches ABX at A; (3) When it cuts ABX in the points B and X; (4) When it touches ABX at A'; (5) When it falls without ABX. These different positions are drawn on fig. 29, where it will be seen that the points A and A' are diametrically opposite on ABX; also that B and X are equidistant from the points A and A'.

The following results are therefore geometrically manifest:—

If $a < b$,	the triangle is impossible:	see circle 1.
If $a = b$,	„ „ evanescent:	„ 2.
If $a > b$ and $< \pi - b$,	two identical solutions:	„ 3.
If $a = \pi - b$,	the triangle is evanescent:	„ 4.
If $a > \pi - b$,	„ „ impossible:	„ 5.

73. *Thirdly.*—Let $b > \frac{1}{2}\pi$. Let A' be the vertex of the triangle to be constructed with the given parts, and let $A'C = b$. With C as pole, describe a circle of radius a which, as already explained, may assume any of the five positions (fig. 29), according to the magnitude of a. Hence—

If $a < \pi - b$,	the triangle is impossible:	see circle 1.
If $a = \pi - b$,	„ „ evanescent:	„ 2.
If $a > \pi - b$ and $< b$,	two identical solutions:	„ 3.
If $a = b$,	the triangle is evanescent:	„ 4.
If $a > b$,	„ „ impossible:	„ 5.

74. Case II.—$A < \frac{1}{2}\pi$.

Draw an arc AC (fig. 30) equal to b, and making the given angle A with the great circle $B_1B_2B_3$..... With C




Fig. 30.

as pole, and a radius a, describe a circle. The position of this circle, as in Case I., will depend entirely on the value of the side a; and eight positions, with respect to $B_1B_2B_3$, ... are shown on fig. 30, viz., circles 0, 1, 2, &c. The circles numbered 0 and 7 respectively have no point in common

with $B_1B_2B_3$, . . . which we may regard as the base of the triangle under consideration.

We shall discuss the different cases under the heading of this Article—firstly, when $b = \frac{1}{2}\pi$; secondly, when $b < \frac{1}{2}\pi$; thirdly, when $b > \frac{1}{2}\pi$.

Firstly.—Let $b = \frac{1}{2}\pi$. Then C, the vertex of the triangle, is the middle point of the arc AA', and we have—

If $a < A$, the triangle is impossible: see circle 0.
If $a = A$, „ „ unique: see circle 1.
If $a > A$ and $< b$, „ „ ambiguous: see circle 2.
If $a = b = \frac{1}{2}\pi$, „ „ evanescent: see circles 3 & 5.
If $a > b$, „ „ impossible: see circles 6 & 7.

Secondly.—Let $b < \frac{1}{2}\pi$. The following results are evident from fig. 30:—

If $\sin a < \sin b \sin A$, the triangle is impossible.
(See circles 0 and 7.)
If $\sin a = \sin b \sin A$, „ „ unique.
(See circle 1.)
If $\sin a > \sin b \sin A$ and $a < b$, „ „ ambiguous.
(See circle 2.)
If $a = b$, „ „ unique.
(See circle 3.)
If $a > b$, and $< \pi - b$, „ unique.
(See circle 4.)
If $a = \pi - b$, „ evanescent.
(See circle 5.)
If $a > \pi - b$, „ impossible.
(See circles 6 and 7.)



Thirdly.—Let $b > \frac{1}{2}\pi$. Let $A'C$ (fig. 30) be the given side b, and A' the given angle ($= A$).

Proceeding as before, we have—

If $\sin a < \sin b \sin A$, (See circle 0.)	the triangle is	impossible.
If $\sin a = \sin b \sin A$, (See circle 1.)	„ „	unique.
If $\sin a > \sin b \sin A$ and $a < \pi - b$, (See circle 2.)	„ „	ambiguous.
If $a = \pi - b$, (See circle 3.)	„ „	unique.
If $a > \pi - b$, and $< b$, (See circle 4.)	„ „	unique.
If $a = b$, (See circle 5.)	„ „	evanescent.
If $a > b$, (See circles 6 and 7.)	„ „	impossible.

75. In the previous Articles, the general cases of ambiguity when A is equal to, or less than, a quadrant, and a and b any values between 0° and π, have been discussed. The following additional cases are left as exercises for the reader when A is obtuse.

Examples.

1. Given $A > \frac{1}{2}\pi$ and $b < \frac{1}{2}\pi$, the triangle is impossible if $a < b$.

2. In the same case the triangle is unique, when a lies between b and $\pi - b$.

3. Given A and b, as in Ex. 1, the triangle is ambiguous when a lies between $\pi - b$ and the greater value of $\sin^{-1}(\sin b \sin A)$.

4. Given A obtuse and b acute, the triangle is impossible if a is greater than the greater value of $\sin^{-1}(\sin b \sin A)$.

5. Given A and b both obtuse, the triangle is impossible when $a < b$.

6. Given A and b both obtuse; find the limiting values of a giving rise to an ambiguity.

Ans. $a > b$ and $< \sin^{-1}(\sin b \sin A.)$

7. What values assigned to the given parts give rise to an *indeterminate* construction?

*8. If a, b, c are each $< \frac{1}{2}\pi$, the greater angle alone *may* exceed $\frac{1}{2}\pi$.

[Let $a > b > c$; then both $\cos b$ and $\cos c > \cos a$, and therefore $> \cos c \cos a$, or $\cos a \cos b$; therefore, &c., Art. 26 (1).]

*9. If a alone $> \frac{1}{2}\pi$, A *must* exceed $\frac{1}{2}\pi$.

[Apply Art. 26 (1).]

*10. If a and b are each $> \frac{1}{2}\pi$, and $c < \frac{1}{2}\pi$; prove that—

(1) the greatest angle (A) must be $> \frac{1}{2}\pi$;

(2) B may be $> \frac{1}{2}\pi$;

(3) C must be $< \frac{1}{2}\pi$.

[Proof as before.]

*11. If $\cos a$, $\cos b$, $\cos c$ are all negative; then $\cos A$, $\cos B$, $\cos C$ are all necessarily negative.

12. In a spherical triangle of the five products

$\cos a \cos A$, $\cos b \cos B$, $\cos c \cos C$, $\cos a \cos b \cos c$, $-\cos A \cos B \cos C$,

one is negative, the other four being positive.—(Cotterill).

[Apply Examples 8, 9, 10, 11.]

76. **The Subsidiary Angle.**—It has been seen that, when two sides and the included angle of a triangle were given, the value of the third side was made to depend on the values ascertained for the base angles. It is likewise evident, in Cases V. and VI. of this Chapter, that the values

* *Educational Times*, vol. xli., reprint, p. 89.

of two of the parts depended on the value or values determined for the remaining part. An error in this latter will, consequently, cause an error in the values of parts found from it.

By means of the subsidiary angle, each of the parts required are determined from the data in question in a manner independent of one another.

(1) In CASE III., Art. 62, c may be found in terms of a, b, C, from a formula adapted to logarithmic computation: thus—

$$\cos c = \cos a \cos b + \sin a \sin b \cos C$$

$$= \cos b (\cos a + \sin a \tan b \cos C).$$

Let $$\tan \theta = \tan b \cos C. \qquad (1)$$

Hence

$$\cos c = \cos b (\cos a + \sin a \tan \theta) = \cos b \cos (a - \theta) \sec \theta;$$

or $$\frac{\cos b}{\cos c} = \frac{\cos \theta}{\cos (a - \theta)}. \qquad (2)$$

The subsidiary angle θ is found from (1), and c can therefore be calculated from (2).

NOTE.—It is evident, from (2), that θ and $a - \theta$ are the segments of a, made by the perpendicular to it from the opposite angle A. (Cf. Chap. IV., Ex. 11.) The solution of the triangle is thus shown to be equivalent to the solutions of these right-angled triangles.

(2) In CASE IV., Art. 67, C may be found directly from the given parts A, B, c, in a form adapted to logarithmic computation: thus—

$$\cos C = -\cos A \cos B + \sin A \sin B \cos c$$

$$= \cos B(-\cos A + \sin A \tan B \cos c).$$

Let $$\cot \theta = \tan B \cos c.$$

Hence

$$\cos C = \cos B(-\cos A + \sin A \cot \theta) = \cos B \sin(A-\theta) \operatorname{cosec} \theta;$$

or $$\frac{\cos B}{\cos C} = \frac{\sin \theta}{\sin(A-\theta)};$$ therefore, &c.

It is obvious that the subsidiary angle θ is in this case a segment of the angle A made by the perpendicular from A on the opposite side. (Cf. Chap IV., Ex. 17.)

Examples.

1. Having given a, b, A, adapt the formula

$$\cot a \sin b = \cot A \sin C + \cos b \cos C$$

to logarithmic computation for the angle C.

Ans. $\tan a : \tan b = \cos \theta : \cos(C - \theta.)$

2. What is the geometrical interpretation of θ in Ex. 1 ?

[Cf. Art. 44 (2).]

3. Given, as before, a, b, A; show how to find c in a logarithmic form.

Ans. $\cos a : \cos b = \cos(c - \theta) : \cos \theta$, where $\tan \theta = \tan b \cos A$.

[Art. 26 (1.)]

4. Prove the result given in Ex. 3 geometrically.

5. Having given A, B, a; construct geometrically the subsidiary angles required—(1) to calculate C; (2) to calculate c.

Ans. Draw a perpendicular from C to c, and take—(1) the segment of C adjacent to a; (2) the segment on c adjacent to a.

6. Having given $a = 84° 30'$, $b = 56° 46'$, $C = 73° 52'$; calculate c by means of the subsidiary angle.

By aid of Art. 76, we have

$$L \tan \theta = L \tan b + L \cos C - 10$$

$$= 9{\cdot}6274648;$$

therefore $\theta = 22° 58' 54''$.

Again, $$L \cos c = L \cos b + L \cos (a - \theta) - L \cos \theta$$

$$= 9{\cdot}4531418;$$

therefore $c = 73° 30' 28''$.

7. Having given $a = 71° 35' 16''$, $b = 40° 33' 12''$, $C = 103° 57' 20\frac{1}{4}''$; calculate c by means of the subsidiary angle.

As in Ex. 6,

L tan supplement of $\theta = 9{\cdot}3146412$; therefore $\theta = 168° 20' 23''$.

$L \cos c = 8{\cdot}9600566$; therefore $c = 84° 46'$.

8. Having given $A = 54° 40'$, $B = 75° 20'$, $c = 40° 0' 4''$; calculate C by means of a subsidiary angle.

By Art. 76,

$$L \cot \theta = L \tan B + L \cos c - 10;$$

therefore $\theta = 18° 51' 48''$.

Again, $$L \cos C = L \cos B + L \sin (A - \theta) - L \sin \theta$$

$$= 9{\cdot}6609932;$$

therefore $C = 62° 44'$.

9. Having given $A = 56° 58'$, $B = 39° 26'$, $c = 68° 48'$; calculate C by means of the subsidiary angle.

Ans. $\theta = 73° 26' 16''$, $C = 103° 12' 25''$.

EXAMINATION PAPERS.

[The following Examples have been taken chiefly from University Calendars, Science and Art Examination Papers, and Mathematical Journals. They are arranged in sets, without any attempt at graduation. Answers are given in almost all cases, and hints to the solution are given where any difficulty is presented.]

I.

1. Being given of a triangle the base and difference of base angles, and that the sum of its sides is 180°, construct it.

2. Prove the following formula for an equilateral triangle:—

$$\log \sin \frac{A}{2} + \log \cos \frac{a}{2} + \log 2 = 0.$$

3. Find $\cos x$, in terms of a and b, from the equations

$$\begin{aligned} \cos a &= \cos \theta \cos x, \\ \cos b &= \cos \phi \cos x, \\ \cos (\theta + \phi) &= \cos a \cos b. \end{aligned}$$

(*Science and Art Hon.*)

4. Given the base, sum of sides, and one base angle; find the other parts of the triangle.

(*Science and Art Hon.*)

5. If an angle of a triangle be equal to (or supplemental to) the opposite side, show that

$$1 - \sec^2 a - \sec^2 b - \sec^2 c + 2 \sec a \sec b \sec c = 0.$$

6. Find the locus of the intersection of equal tangents to two small circles.

7. If each leg of a tripod be inclined at an angle ϕ to each of the other legs; find the angle between a leg and the horizontal plane, the legs being supposed equal to each other.

8. Given that $\tan x = \tan a \cos b$,

and $\tan y = \tan b \cos a$;

show that $\sec (x \pm y) = \sec a \sec b \pm \tan a \tan b$.

(*Science and Art. Hon.*)

9. Given $a = 84° 30'$, $b = 46° 10'$, $C = 75° 46'$;

calculate A, B, and c.

10. Given $a = 86° 45'$ $b = 74° 35'$, $c = 63° 40'$;

calculate A, B, and C.

III.

1. Given the base c of a triangle, and that

$$\tan \tfrac{1}{2} a \tan \tfrac{1}{2} b = \tan^2 \frac{\beta}{2};$$

find $a - b$, β being the bisector of the base.

2. If $C = A + B$, show that

$$1 - \cos a - \cos b + \cos c = 0.$$

3. If A denote one of the angles of an equilateral triangle, and A' an angle of its polar triangle, show that

$$\cos A \cos A' = \cos A + \cos A'.$$

Explain the result when $A = 60°$.

(*Science and Art Exam. Papers.*)

4. Express in terms of C and c,

$$\frac{\cos a \cos B - \cos b \cos A}{\sin a - \sin b}.$$

5. If four points A, B, C, D on a great circle be joined to a point P on the sphere, show that

$$\begin{vmatrix} \cot AB & \cot APB & 1 \\ \cot AC & \cot APC & 1 \\ \cot AD & \cot APD & 1 \end{vmatrix} = 0.$$

6. Using the notation of Ex. 5, show that if cot AB, cot AC, cot AD are in Arithmetical Progression, cot APB, cot APC, cot APD are also in Arithmetical Progression, and hence *a harmonic row always subtends a harmonic pencil at any point on the sphere.*

7. The sum of the sides of a spherical polygon is less than 2π.

8. Prove the following formula for the reduction of the parts of a spherical triangle:—

$$(\sec a \sin b \cos A - \sin c)^2 + (\sec a \cos b - \cos c)^2 (1 - \operatorname{cosec}^2 a \sin^2 A) = \tan^2 a \cos^2 B \cos^2 C.$$

(*Educational Times*, vol. xxxvii. p. 69.)

9. Given $a = 72° 44'$, $b = 36° 22'$, $C = 56° 12'$; calculate A, B and c.

10. Given, in a right-angled triangle, $a = 86° 45'$, $b = 108° 20'$; calculate A, B, and c.

III.

1. If two small circles on a sphere touch each other, show that the great circle joining their poles passes through their point of contact.

2. Prove that

$$\cos A = \frac{\cos a \sin b - \sin a \cos b \cos C}{\sin c};$$

and hence

$$\cos A + \cos B = \frac{2 \sin (a + b) \sin^2 \tfrac{1}{2} C}{\sin c}.$$

(Cf. p. 45, Ex. 1.)

3. Show that the arcs joining the middle points of the sides of the colunar triangles of a given spherical triangle are equal to the corresponding angles of the chordal triangle.

4. Show that a triangle can be constructed having sides $90° - \alpha$, $90° - \beta$, $90° - \gamma$, and angles $90 - \frac{1}{2} a$, $90 - \frac{1}{2} b$, $90 - \frac{1}{2} c$, where α, β, γ are the arcs joining the middle points of the sides of a triangle whose sides are a, b, c.

5. Find the locus of the middle points of the sides of a triangle, having given the base and the sum of the three angles.

6. Show that

$$\Sigma \sin^2 a \cot XY = 0,$$

where XY is the intercept made by the bisectors of the angle A on the side a of a spherical triangle.

7. Deduce the analogue *in plano* of Ex. 6.

8. Having given the sides of a triangle, each $= 60°$; find the segments of the bisectors of the angles.

9. In a right-angled triangle, being given

$$A = 47° \ 45', \quad B = 63° \ 30';$$

calculate the sides.

10. Given $a = 75° \ 12', \quad b = 63° \ 27', \quad c = 52° \ 23';$

calculate A, B, and C.

IV.

1. Having given the relation

$$\tan \tfrac{1}{2} a \tan \tfrac{1}{2} b = \tan^2 \tfrac{1}{2} c ;$$

show that

$$\cos \beta = \frac{1 - \sin^2 c \cos^2 \theta}{1 - \sin^2 c \sin^2 \theta},$$

where β is the bisector of the side $2c$ drawn from the opposite vertex, and θ the angle it makes with that side.

(*Dublin Univ. Exam. Papers.*)

2. Show that the sum of the three arcs joining the middle points of the sides of the three colunar triangles of a given triangle is equal to two right angles.

3. If
$$\sin^2 \frac{c}{2} = \sin^2 \frac{a}{2} + \sin^2 \frac{b}{2},$$

prove that $C = A + B$.

4. Given the hypotenuse of a right-angled triangle; prove that the difference of the sides is a maximum when their sum is equal to a quadrant.

5. Find the relation connecting the angles of a triangle if

$$1 + \cos a + \cos b + \cos c = 0.$$

6. Prove that

$$\frac{\cos (B - C)}{\cos (A - C)} = \frac{\tan \tfrac{1}{2} a - \tan \tfrac{1}{2} b \cos C - \tan \tfrac{1}{2} c \cos B}{\tan \tfrac{1}{2} b - \tan \tfrac{1}{2} a \cos C - \tan \tfrac{1}{2} c \cos A}.$$

7. The great circle through the poles of two small circles is perpendicular to the great circle through their points of intersection.

8. If a and a' are the segments of the bisector of the side a of a triangle, show that

$$\frac{\sin a}{\sin a'} = 2 \cos \frac{a}{2}.$$

9. Given $A = 58° 16'$, $B = 41° 28'$, $c = 75° 54'$;

calculate a, b, and C.

10. Given $a = 47° 45'$, $c = 69° 30'$, $C = 90°$;

calculate A, B, and b.

*

V.

1. If a, b, c, d be the sides of a spherical quadrilateral, δ and δ' its diagonals, and if a and c, b and d, δ and δ' intersect at angles θ, ϕ, ψ, respectively, show that

$$\sin a \sin c \cos\theta + \sin b \sin d \cos\phi + \sin\delta \sin\delta' \cos\psi = 0.$$

2. Find the locus of a point P, whose distances from three fixed points A, B, C are connected by the relation

$$\cos AP + \cos BP + \cos CP = \text{const.}$$

3. If α, α'; β, β'; γ, γ', be the segments of the perpendiculars to the sides of a spherical triangle drawn from the opposite vertices, show that

$$\tan\alpha \tan\alpha' = \tan\beta \tan\beta' = \tan\gamma \tan\gamma'.$$

4. Using the notation of Ex. 3, show that

$$\frac{\cos(\alpha+\alpha')}{\cos\alpha \cos\alpha'} = \frac{\cos(\beta+\beta')}{\cos\beta \cos\beta'} = \frac{\cos(\gamma+\gamma')}{\cos\gamma \cos\gamma'}.$$

5. Simplify the expression

$$\frac{\cos A \cot a + \cos B \cot b}{\cot a \cot b - \cos A \cos B}.$$

6. If the bisectors of the angles of a triangle meet at O, show that the angle AOB is supplemental to the angle COD, where OD is a perpendicular from O on BC.

7. Using the notation of Ex. 6, show that—

(1) $\tan AO \cos BOC = \tan BO \cos COA = \tan CO \cos AOB$;

(2) $\sin BOC : \sin COA : \sin AOB = \cos\frac{1}{2}A : \cos\frac{1}{2}B : \cos\frac{1}{2}C$.

8. The angle in a semicircle is a right angle; what is the analogous theorem on the sphere?

9. Given $\quad a = 85° 16', \quad b = 63° 24', \quad C = 76° 12';$

calculate A, B, and c.

10. In a right-angled triangle, given $c = 48° 45'$, and $A = 21° 27'$; calculate a, B, and b.

VI.

1. Two places are situated on the same parallel of latitude λ; find the difference of the distances sailed over by two ships passing between them, one keeping to the great circle course, the other to the parallel; the difference of longitude of the places being $2l$.

2. Having given the base c of a spherical triangle, and the ratio $\sin \frac{1}{2} a : \sin \frac{1}{2} b$, find the locus of the vertex.

3. If α, β, γ be the sides of the triangle formed by joining the middle points of the sides b, c; c, a; a, b, respectively, of a given triangle, show that

$$\sin \alpha : \sin \beta : \sin \gamma :: \operatorname{cosec} \frac{a}{2} : \operatorname{cosec} \frac{b}{2} : \operatorname{cosec} \frac{c}{2}.$$

4. Using the notation of the preceding Example, show that

$$\tan \alpha \sin a = \tan \beta \sin b = \tan \gamma \sin c.$$

5. If the sides of a spherical triangle be each 60°; show that the circles described, each having a vertex for pole, and passing through the middle points of the sides which meet at it, have the sides of the supplemental triangle for common tangents.

6. If the bisectors of the angles of a triangle meet at O; show that

$$\frac{\sin BOC}{\sin AO} : \frac{\sin COA}{\sin BO} : \frac{\sin AOB}{\sin CO} = \sin a : \sin b : \sin c.$$

7. Using the notation of Ex. 6; show that

$$\sin^2 AO : \sin^2 BO : \sin^2 CO = \frac{\sin (s-a)}{\sin a} : \frac{\sin (s-b)}{\sin b} : \frac{\sin (s-c)}{\sin c}.$$

8. If the base of a triangle be divided into four equal parts, which subtend angles $\theta_1, \theta_2, \theta_3, \theta_4$ at the vertex; show that

$$\frac{\sin (\theta_1 + \theta_2)}{\sin (\theta_3 + \theta_4)} = \frac{\sin \theta_1 \sin \theta_3}{\sin \theta_2 \sin \theta_4}.$$

9. Given $A = 86° 44'$, $B = 58° 32'$, $c = 79° 40'$;

calculate a, b, and C.

10. Given $A = 46° 45'$, $c = 75° 40'$, $C = 90°$;

calculate a, b, and B.

VII.

1. If a great circle cuts a fixed great circle at a constant angle ω, find the locus of its pole.

2. Show from Ex. 1, Paper v., that the perpendiculars of a spherical triangle are concurrent.

3. Given three fixed points A, B, C; find the locus of a fourth point P on the sphere, such that

$$l \cos AP + m \cos BP + n \cos CP = \text{const.},$$

where l, m, n are given quantities.

4. If β be the bisector of the base c of a triangle, show that

$$\sin \beta = \frac{\sqrt{\sin^2 a + \sin^2 b + 2 \sin a \sin b \cos C}}{2 \cos \frac{1}{2} c}.$$

5. Deduce the corresponding equation for a plane triangle.

6. A spherical quadrilateral is circumscribed to a small circle; show that the sum of one pair of opposite sides is equal to the sum of the other pair.

7. In a spherical triangle, prove that—

(1) If three sides are acute, two angles are acute;

(2) If one side is acute and one side is not acute, one angle is obtuse and two are acute;

(3) If two sides are obtuse and one acute, one angle is obtuse;

(4) If two sides are obtuse and the other is not acute, all the angles are obtuse.

(*Educational Times.*)

8. Prove the relations—

(1) $\cos^2 \frac{1}{2} c = \cos^2 \frac{1}{2} (a - b) \cos^2 \frac{1}{2} C + \cos^2 \frac{1}{2} (a + b) \sin^2 \frac{1}{2} C$;

(2) $\sin^2 \frac{1}{2} c = \sin^2 \frac{1}{2} (a - b) \cos^2 \frac{1}{2} C + \sin^2 \frac{1}{2} (a + b) \sin^2 \frac{1}{2} C$.

9. Given $a = 68° \ 16'$, $b = 53° \ 24'$, $C = 86° \ 40'$;

calculate A, B, and c.

10. Given $a = 38° \ 40'$ $c = 73° \ 30'$, $C = 90°$;

calculate A, B, and b.

VIII.

1. From a point P on a small circle, in which is inscribed an equilateral triangle ABC, arcs of great circles PA, PB, PC are drawn; prove that

$$\sin \tfrac{1}{2} AP + \sin \tfrac{1}{2} BP = \sin \tfrac{1}{2} CP,$$

and give the corresponding property *in plano*.

2. Prove *Legendre's Theorem* from the formula

$$\cos^2 \tfrac{1}{2} A = \frac{\sin s \sin (s-a)}{\sin b \sin c}.$$

3. Prove *Legendre's Theorem* from each of the formulæ for $\sin \frac{1}{2} A$ and $\tan \frac{1}{2} A$, respectively, in terms of the sides.

4. Deduce the area of a plane triangle, in terms of the base and base angles, from the formulæ

$$\sin^2 \frac{a}{2} = \frac{-\cos S \cos (S-A)}{\sin B \sin C}, \text{ \&c., \&c.}$$

5. From the *rule of sines*, p. 37, find the area of a plane triangle in terms of the base and base angles.

6. Prove directly, employing the solution on p. 35, that

$$\frac{\tan^2 \dfrac{A}{2} - \tan^2 \dfrac{A'}{2}}{\tan^2 \dfrac{A'}{2}} = \frac{bc}{3r^2}, \text{ nearly.}$$

7. If A and A', B and B', C and C', denote the pairs of opposite angles of a complete quadrilateral, prove the relation

$$\cos B \cos B' \sim \cos C \cos C' = \sin A \sin A' \cos AA'.$$

8. Using the notation of Ex. 7, show that for a plane quadrilateral—

(α) $\cos B \cos B' \sim \cos C \cos C' = \sin A \sin A'$;

(β) $\sin A \sin A' \sim \sin B \sin B' = \sin C \sin C'$.

9. Given $a = 87° 27'$, $b = 63° 19'$, $c = 74° 40'$;

calculate A, B, and C.

10. Given $a = 38° 50'$, $B = 47° 44'$, $C = 90°$;

calculate A, b, and c.

IX.

1. If A and A', B and B', C and C', denote the pairs of opposite angles of a quadrilateral, and δ_1, δ_2, δ_3, the diagonals, prove that the (*algebraic*) sum

$$\sin A \sin A' \cos \delta_1 + \sin B \sin B' \cos \delta_2 + \sin C \sin C' \cos \delta_3 = 0.$$

2. If two diagonals of a spherical quadrilateral are each quadrants, the third diagonal is also a quadrant.

(*Dub. Univ. Exam. Papers.*)

3. The intersections of the corresponding sides of a triangle and its polar triangle lie on a great circle.

4. Prove generally that if three great circles be drawn from the vertices of a triangle ABC through any point O, intersecting the opposite sides of the triangle respectively at angles θ, ϕ, ψ,

$$\sin a \sin AO \cos \theta + \sin b \sin BO \cos \phi + \sin c \sin CO \cos \psi = 0.$$

5. In a plane triangle ABC three right lines AP, BP, CP are drawn, cutting the opposite sides respectively in X, Y, Z; show that

$$a\,AP \cos X + b\,BP \cos Y + c\,CP \cos Z = 0.$$

6. Using the notation of Ex. 4 for an equilateral spherical triangle, show that

$$\sin AO \cos \theta + \sin BO \cos \phi + \sin CO \cos \psi = 0.$$

7. If ϕ denote the perpendicular from the angle A on the opposite side a, and β and γ the segments of the base made by it, prove the relation

$$\sin^2 p \cos \beta \cos \gamma - \sin p \sin (\beta + \gamma) \cot A - \sin \beta \sin \gamma = 0;$$

and hence show that of all spherical triangles described on the same base, and having equal altitudes, the vertical angle is a maximum when the triangle is isosceles—(W. Nicolls).

8. If a, b, c, d be the sides of a cyclic spherical quadrilateral, δ, one of the diagonals, is given by the equation

$$\sin^2 \tfrac{1}{2}\delta = \frac{(\sin \tfrac{1}{2}a \sin \tfrac{1}{2}d + \sin \tfrac{1}{2}b \sin \tfrac{1}{2}c)(\sin \tfrac{1}{2}a \sin \tfrac{1}{2}c + \sin \tfrac{1}{2}b \sin \tfrac{1}{2}d)}{\sin \tfrac{1}{2}a \sin \tfrac{1}{2}b + \sin \tfrac{1}{2}c \sin \tfrac{1}{2}d}.$$

9. Given $A = 69°\ 56'$, $B = 46°\ 36'$, $c = 76°\ 28''$;
calculate a, b, and C.

10. Given $a = 45°\ 30'$, $B = 63°\ 45'$, $C = 90°$;
calculate A, b, and c.

X.

1. Given $A + a$, $B + b$, and $C + c$; solve the triangle.

(*Dub. Univ. Exam. Papers.*)

2. If O be the mid-point of an equilateral triangle ABC, and P any point on the surface of the sphere, show that

$$(\tan PO \tan AO)^2 (\cos AP + \cos BP + \cos CP)^2$$
$$= 4 [\cos^2 AP + \cos^2 BP + \cos^2 CP - \cos BP \cos CP - \cos CP \cos AP - \cos AP \cos BP].$$

3. The perpendiculars of a triangle are concurrent; hence show from the supplemental figure that if two diagonals of a quadrilateral are quadrants, the third is also a quadrant.

4. Using the notation of Ex. 2, and each side of the triangle being a quadrant, show that

$$\cos AP + \cos BP + \cos CP = \sqrt{3} \cos OP.$$

5. Construct a triangle right-angled at C, being given the angle A and the sum or difference of the sides which contain it.

6. Given the base of a spherical triangle, and the length of the arc joining the middle points of the sides; find the locus of the vertex.

7. If the angle A of a right-angled triangle be given, show that the difference of the sides which contain it is a maximum when their sum is a quadrant.

8. Given the base of a triangle, and the locus of the vertex a great circle; construct the triangle, having a maximum or minimum sum of sides.

9. Given $a = 64° 30'$, $b = 48° 12'$, $C = 90°$; calculate the remaining parts.

10. Solve the triangle when

$$a = 56° 48', \quad b = 42° 26', \quad C = 83° 12'.$$

XI.*

1. Find the locus of a point P, such that

$$\sin PX + \sin PY = \text{const.},$$

where PX and PY are secondaries to two given great circles.

2. Find the locus of a point P, such that

$$l \sin PX \pm m \sin PY, = \text{const.},$$

where PX and PY are secondaries to two given great circles and l and m constants.

3. Find the locus of a point P, such that

(α) $\sin PX + \sin PY + \sin PZ = \text{const.}$;

(β) $l \sin PX \pm m \sin PY \pm n \sin PZ = \text{const.}$,

where X, Y, Z are the feet of the perpendiculars from P on the sides of a spherical triangle.

4. In a right-angled triangle prove that—

(α) The tangents of the segments of the hypotenuse made by the perpendicular from the right angle, and the sine of the perpendicular are in geometrical progression.

(β) The tangents of the hypotenuse, either side, and the adjacent segment of the hypotenuse made by the perpendicular, are also in geometrical progression.

5. Construct a right-angled triangle, having given the hypotenuse, and the locus of the vertex a given great circle.

6. Having given the base and the sum of the sides, show that the product of the sines of the perpendiculars from the extremities of the base on the bisector of the external vertical angle is constant.

7. Having given the base and the difference of the sides, show that the product of the sines of the perpendiculars from the extremities of the base on the bisector of the internal vertical angle is constant.

8. If p_1 and p_2 denote the perpendiculars from the base angles on the bisector of the external vertical angle, and p_3 and p_4 those on the bisector of the internal vertical angle, prove the relation

$$\sin p_1 \sin p_2 + \sin p_3 \sin p_4 = \sin a \sin b.$$

9. Solve the triangle, having given $A = 29° 33'$, $B = 87° 21'$, and $C = 90°$.

10. Having given $a = 62° 56'$, $b = 39° 28'$, $C = 75° 18'$, solve the triangle.

* This Paper and part of the preceding were communicated to us by Dr. Traill, F.T.C.D.

XII.*

1. If β be the bisector of the base AB of a triangle ABC, and if β' be the bisector of the base BA' of the colunar BCA', show that

$$(\alpha)\ \sin^2\beta\cos^2\frac{c}{2} - \sin^2\beta'\sin^2\frac{c}{2} = \sin a\sin b\cos C;$$

$$(\beta)\ \sin^2\beta\cos^2\frac{c}{2} + \sin^2\beta'\sin^2\frac{c}{2} = \tfrac{1}{2}(\sin^2 a + \sin^2 b).$$

2. Show that if $E = A + B + C - \pi$, then

$$\sin\tfrac{1}{2}E\sin(A - \tfrac{1}{2}E)\sin(B - \tfrac{1}{2}E)\sin(C - \tfrac{1}{2}E) = N^2.$$

3. Show that

$$\tan^2\frac{c}{2} = \frac{\tan^2\frac{1}{2}a - 2\tan\frac{1}{2}a\tan\frac{1}{2}b\cos C + \tan^2\frac{1}{2}b}{1 + 2\tan\frac{1}{2}a\tan\frac{1}{2}b\cos C + \tan^2\frac{1}{2}a\tan^2\frac{1}{2}b}.$$

4. Show that

$$\Sigma\tan\frac{a}{2}\frac{\sin(B - C)}{\sin B\sin C} = 0.$$

5. If the base AB of a triangle be divided into segments α and β by an arc θ drawn from the vertex C, show that

$$(\alpha)\quad \cos\theta\sin c = \cos a\sin\alpha + \cos b\sin\beta.$$

If θ be the bisector of the vertical angle, show that

$$(\beta)\quad \cos\theta = \frac{\sin(a + b)}{\sin a\cos\alpha + \sin b\cos\beta}.$$

6. If the internal bisector of the angle C of a triangle meets the opposite side at an angle θ, and divides it into segments α and β adjoining the angles A and B, respectively, and if the external bisector of C makes an angle θ with the opposite side, show that

$$(1)\quad \sin\alpha\sin\beta = \frac{\sin a\sin b\sin^2 c}{\sin^2 a + \sin^2 b + 2\sin a\sin b\cos c};$$

$$(2)\quad \sin\theta = \frac{\sin\frac{1}{2}C}{\sin c}\sqrt{\sin^2 a + \sin^2 b + 2\sin a\sin b\cos c};$$

$$(3)\quad \cos\theta\cos\theta' = \frac{\sin(A + B)\sin(A - B)}{2\sin C}.$$

* The greater part of this Paper was communicated to us by Mr. W. S. M'Cay, F.T.C.D.

7. Being given the angle A of a right-angled triangle, show that when $c - b$ is a maximum,

$$\cos A = \cos^2 a.$$

8. Given $A = 56° 58'$, $B = 39° 26'$, $c = 68° 48'$;
calculate the remaining parts.

9. Given $A = 54° 40'$, $B = 75° 20'$, $C = 62° 44'$
solve the triangle.

10. Given $a = 84° 30'$, $b = 56° 46'$, $C = 73° 52'$;
calculate the remaining parts.

11. Given $a = 40° 45'$, $A = 51° 50'$, $C = 90°$;
solve the triangle.

12. Given $a = 51° 20'$, $A = 62° 12'$, $C = 90°$;
solve the triangle.

13. Given $A = 81° 54'$, $B = 67° 12'$, $c = 87° 42'$;
solve the triangle.

14. Given $A = 67° 37'$, $B = 39° 19'$, $c = 84° 46'$;
solve the triangle.

15. Given $A = 63° 40'$, $B = 87° 10'$, $C = 59° 46'$;
solve the triangle.

ANSWERS TO EXAMINATION PAPERS.

II.

1. The sum of the base angles is given by Art. 24.

2. Apply formula for $\sin \frac{1}{2} A$ (Art. 33).

3. x is the perpendicular from the vertex on the hypotenuse of a right-angled triangle, the sides of which are a and b, and the hypotenuse $\theta + \phi$; hence

$$\sin x = \frac{\sin a \sin b}{\sin (\theta + \phi)}. \qquad \text{[Page 50, Ex. 2.]}$$

4. The remaining parts may be determined by aid of *Napier's Analogies*, or geometrically from fig. 22, p. 70.

5. Suppose $a = A$; then, from Art. 26,

$$\sec a = \sec b \sec c - \tan b \tan c;$$

and therefore

$$\tan^2 b \tan^2 c = (\sec a - \sec b \sec c)^2;$$

therefore, &c.

6. A great circle perpendicular to that through the poles of the small circles, and dividing it so that the ratio of the cosines of its segments is equal to the ratio of the cosines of radii of the small circles—in other words, the great circle through the intersections of the small circles.

7. If θ be the inclination to the horizon,

$$\cos \theta = \frac{2}{\sqrt{3}} \sin \frac{\phi}{2}.$$

8. x and y are the segments of the base of a triangle made by the perpendicular from the vertex, the sides a and b being equal to the opposite angles; therefore, &c.

9. $A = 95° \ 56' \ 17{\cdot}87''$, $B = 46° \ 7' \ 15{\cdot}53''$, $c = 75° \ 56' \ 27''$.

10. $A = 94° \ 3' \ 50{\cdot}28''$, $B = 74° \ 23' \ 45{\cdot}16''$, $C = 63° \ 33' \ 42{\cdot}48''$.

III.

1. We have

$$\frac{1-\tan\frac{1}{2}a\tan\frac{1}{2}b}{1+\tan\frac{1}{2}a\tan\frac{1}{2}b}=\frac{1-\tan^2\beta}{1+\tan^2\beta}.$$

nce

$$\frac{\cos\frac{1}{2}(a+b)}{\cos\frac{1}{2}(a-b)}=\cos\beta=\frac{\cos a+\cos b}{2\cos\frac{1}{2}c}; \qquad \text{(Art. 42.)}$$

l therefore $$\cos^2\tfrac{1}{2}(a-b)=\cos\tfrac{1}{2}c.$$

2. Use *Delambre's Analogies*. See also p. 75, Ex. 29.

3. See p. 33, Ex. 8.

4. Substitute for $\cos A$ and $\cos B$ in terms of the sides, and the expression uces to

$$\frac{\cos C+\cos c}{\sin c}.$$

5. Apply the formulae of Art. 37 to the triangles APB, APC, APD, and eliminate $\sin A$ and $\cos A$.

6. This appears at once from the determinant of Ex. 5.

7. See Art. 19 (1).

8. $(\sec a\sin b\cos A-\sin c)^2+(\sec a\cos b-\cos c)^2(1-\operatorname{cosec}^2 a\sin^2 A)$

$$=\left(\frac{\sin b\cos A-\cos a\sin c}{\cos a}\right)^2+\left(\frac{\cos b-\cos a\cos c}{\cos a}\right)^2\left(1-\frac{\sin^2 A}{\sin^2 a}\right)$$

$$=\left(\frac{\sin b\sin c\cos A-\cos a\sin^2 c}{\cos a\sin c}\right)^2+\left(\frac{\sin^2 c\sin^2 a\cos^2 B}{\cos^2 a}\right)\left(1-\frac{\sin^2 A}{\sin^2 a}\right)$$

$$=\left(\frac{\cos a-\cos b\cos c-\cos a\sin^2 c}{\cos a\sin c}\right)^2+\tan^2 a\cos^2 B(\sin^2 c-\sin^2 C)$$

$$=\frac{\cos^2 c(\cos a\cos c-\cos b)^2}{\sin^2 c\cos^2 a}+\tan^2 a\cos^2 B(\sin^2 c-\sin^2 C)$$

$$=\cos^2 c\tan^2 a\cos^2 B+\tan^2 a\cos^2 B(\sin^2 c-\sin^2 C).$$

$$=\tan^2 a\cos^2 B\cos^2 C.$$

9. $A=107°\ 36'\ 9{\cdot}4''$, $B=36°\ 17'\ 20{\cdot}6''$, $c=56°\ 21'\ 30''$.

10 $A=86°\ 54'\ 52{\cdot}7''$, $B=108°\ 18'\ 20{\cdot}9''$, $c=101°\ 1'\ 18{\cdot}5''$.

III.

1. The great circle is the shortest distance between the poles. See Art. 3.

2. In the value of $\cos A$ [p. 30 (2)] substitute the value of $\cos c$ [p. 30 (1)], and reduce.

3. By Ex. 14, p. 11, the radii of the sphere drawn to the middle points of the sides of a colunar are parallel to the sides of the chordal triangle; and hence the angle between them, which is the arc joining the middle points of the sides of the colunar, is equal to the corresponding angle of the chordal triangle.

4. See p. 67, Ex. 4.

5. The locus is the circle LM, fig. 10, p. 21.

6. This follows from the value of cot XY given, p. 56, Ex. 6.

7. See Casey's "Sequel to Euclid," sec. VIII. Ex. 17—

$$\frac{a^2}{XY} + \frac{b^2}{X'Y'} + \frac{c^2}{X''Y''} = 0.$$

8. If α and α' are the segments, we have

$$\cos(\alpha + \alpha') = \frac{1}{\sqrt{3}}, \text{ and } \sin\alpha = \sqrt{3}\sin\alpha';$$

therefore, &c.

9. $a = 41° 17' 48''$, $b = 52° 55' 47{\cdot}5''$, $c = 63° 4' 17''$.

10. $A = 91° 24' 19{\cdot}5''$, $B = 67° 39' 45{\cdot}8''$, $C = 54° 59' 22{\cdot}26''$.

IV.

1. Using the given relation, we have

$$\frac{1-\tan\frac{1}{2}a\tan\frac{1}{2}b}{1+\tan\frac{1}{2}a\tan\frac{1}{2}b}=\frac{1-\tan^2\frac{1}{2}c}{1+\tan^2\frac{1}{2}c};\quad\therefore\quad\frac{\cos\frac{1}{2}(a+b)}{\cos\frac{1}{2}(a-b)}=\cos c.$$

Again (Art. 42),

$$\cos\beta=\frac{\cos a+\cos b}{2\cos c}=\cos^2\tfrac{1}{2}(a-b);$$

and

$$\cos\theta=\frac{\cos a-\cos c\cos\beta}{\sin c\sin\beta}=\frac{\cos a-\cos b}{2\sin c\sin\beta}=-\frac{\sin\frac{1}{2}(a-b)\sin\frac{1}{2}(a+b)}{\sin c\sin\beta}.$$

Substituting, we get the required result.

2. See Ex. 3, Paper III.

3. This follows at once from Exs. 8 and 9, Art. 33. See also Ex. 29, p. 75.

4. For $\quad 2\cos c=2\cos a\cos b=\cos(a+b)+\cos(a-b);$

therefore

$$\cos(a-b)=2\cos c-\cos(a+b);$$

therefore, &c.

5. See Ex. 29, p. 75.

6. Substitute for $\tan\frac{1}{2}a$, $\tan\frac{1}{2}b$, $\tan\frac{1}{2}c$, from Ex. 27, p. 74.

7. Proof as *in plano*.

8. If θ and ϕ be the segments of the angle B, made by the bisector of the side b, we have

$\sin a:\sin a'=\sin\theta\sin c:\sin\phi\sin\frac{1}{2}a$. But $\sin\theta:\sin\phi=\sin a:\sin c$. (Ex. 5, p. 38); therefore, &c.

9. $a=58°\ 33'\ 37{\cdot}48''$, $b=41°\ 47'\ 39{\cdot}62''$, $C=104°\ 48'\ 3{\cdot}58''$.

10. $A=52°\ 12'\ 36{\cdot}7''$, $B=65°\ 41'\ 37{\cdot}2''$, $b=58'\ 36'\ 37{\cdot}8''$.

V.

1. Cf. Examples 24 and 25, pp. 73, 74.

2. $\cos AP + \cos BP = 2 \cos \frac{1}{2} c \cos \beta$, where β is the arc joining P to the middle point of AB. But

$$2 \cos \tfrac{1}{2} c = \frac{\sin \alpha}{\sin \alpha'}$$

(Ex. 8, Paper IV.) Hence

$$\sin \alpha \cos \beta + \sin \alpha' \cos CP = \text{const.} = \sin (\alpha + \alpha') \cos OP,$$

[See Art. 40, p. 58.]

where O is the intersection of the bisectors of the sides of ABC. Hence OP is constant, and therefore the locus is a circle, having O for pole.

3. Apply formula (2), Art. 44.

4. This follows at once from Ex. 3.

5. Reduces to tan c.

6. Proof as *in plano*.

7. By aid of Ex. 6 and Art. 44.

8. $C = A + B$.

9. $A = 91° \ 55' \ 57{\cdot}3$, $B = 63° \ 43' \ 41{\cdot}3''$, $c = 75° \ 33' \ 8''$.

10. $a = 15° \ 56' \ 30''$, $b = 46° \ 42' \ 12{\cdot}4''$, $B = 75° \ 28' \ 35''$.

VI.

1. If 2θ be the arc of the great circle joining the two places, $r \sin\theta = r' \sin l$ = half chord of arc, r', being the radius of the parallel of latitude, is equal to $r \cos\lambda$, where r is the radius of the earth. Hence great circle course $= 2r\theta = 2r \sin^{-1}(\cos\lambda \sin l)$; but the parallel course $= 2r'l = 2rl \cos\lambda$. Hence difference $= 2r[l \cos\lambda - \sin^{-1}(\cos\lambda \sin l)]$.

2. $\sin\frac{1}{2}a : \sin\frac{1}{2}b$ as the chords of the sides a and b. Hence we are given the base and the ratio of the sides of the chordal triangle, and the locus of *its* vertex is therefore a sphere, and hence (Art. 10) the locus of the vertex of the *spherical* triangle is a circle.

3. In fig. 14, p. 21, the arc $YZ = a$. Hence

$$\sin a \sin XYZ = \sin E, \text{ since } ZE = 90°;$$

but

$$\sin E \cos\tfrac{1}{2}c = \sin\tfrac{1}{2}a \sin X, \text{ and } \sin X : \sin Y = \sin\tfrac{1}{2}b : \sin\tfrac{1}{2}a.$$

Hence

$$\sin\alpha = \frac{\sin\frac{1}{2}b}{\cos\frac{1}{2}c}; \text{ and similarly } \sin\beta = \frac{\sin\frac{1}{2}a}{\cos\frac{1}{2}c};$$

therefore, &c.

4. With the preceding Example use Ex. 4, p. 67.

5. If the sides CA and CB be produced to meet the circles again in X and Y, $CX = CY = 90°$; $\therefore$ C is the pole of XY; therefore, &c.

6. See Ex. 6, Paper V.

8. Use the relations given in Examples 1 and 2, p. 37.

9. $a = 81° 46' 6{\cdot}7''$, $b = 57° 43' 43{\cdot}76''$, $C = 82° 56' 4{\cdot}4''$.

10. $a = 44° 53' 9{\cdot}4''$, $b = 69° 32' 55''$, $B = 75° 15' 22''$.

VII.

1. A small circle having the same poles as the fixed circle, and a spherical radius equal to ω.

2. Apply the relation of Paper V. Ex. 1, to a quadrilateral $ABCP$, where P is inside the triangle P; and it follows that if the angle between AP and BC is right, and also that between BP and AC, then the angle between CP and AB will be right also.

3. See Ex. 2, Paper V., and apply the theorem of Art. 40.

4. Reduces to the expression of Art. 42.

5. $$2\beta = \sqrt{a^2 + b^2 + 2ab\cos C};$$

or the corresponding equation, Ex. 14, p. 52.

6. Proof as *in plano*.

7. Let a, b, c, and therefore A, B, C, be in descending order of magnitude; then

$$\tan\tfrac{1}{2}(A+B) = \frac{\cos\frac{1}{2}(a-b)}{\cos\frac{1}{2}(a+b)}\cot\tfrac{1}{2}C;$$

and as $\cos\frac{1}{2}(a-b)$ and $\cos\frac{1}{2}C$ are essentially positive, $\tan\frac{1}{2}(A+B)$ and $\cos\frac{1}{2}(a+b)$ have the same sign; therefore, &c.

8. These relations are easily deduced from Napier's Analogies.

9. $A = 74°\ 2'\ 53{\cdot}1''$, $B = 56°\ 11'\ 54{\cdot}9''$, $c = 74°\ 41'\ 4''$.

10. $A = 40°\ 39'\ 50{\cdot}5''$, $B = 76°\ 17'\ 19{\cdot}6''$, $b = 68°\ 40'\ 9{\cdot}2''$.

VIII.

1. By aid of the property, Ex. 6, p. 28,

$$\sin \tfrac{1}{2} a \sin \tfrac{1}{2} AP + \sin \tfrac{1}{2} a \sin \tfrac{1}{2} BP = \sin \tfrac{1}{2} a \sin \tfrac{1}{2} CP,$$

where a is the side of the equilateral triangle; therefore, &c.

2. Expanding the terms on the right-hand side of the equation, and neglecting the powers of the radius higher than the second; remembering also that

$$\frac{1}{1-n} = (1-n)^{-1} = 1 + n \text{ nearly};$$

if n be very small compared with unity.

Hence
$$\cos^2 \tfrac{1}{2} A = \frac{s(s-a)}{bc} \left[1 - \frac{s^2 + (s-a)^2 - b^2 - c^2}{6r^2}\right]$$
$$= \frac{s(s-a)}{bc} \left[1 - \frac{(s-b)(s-c)}{3r^2}\right] = \cos^2 \tfrac{1}{2} A' - \frac{\Delta^2}{3bcr^2}.$$

Again, let $\frac{1}{2} A = \frac{1}{2} A' + \theta$; then $\cos^2 \frac{1}{2} A = \cos^2 \frac{1}{2} A' - \theta \sin A'$ nearly; hence

$$\theta \sin A' = \frac{\Delta^2}{3bcr^2}, \quad i.e.\ 2\theta = \frac{\Delta}{3r^2};$$

but 2θ is the difference between A and A'; therefore, &c.

(Cf. pp. 34, 35.)

3. The method in either case is similar to that given in the preceding Example. See also pp. 34, 35.

4. Squaring the expression for $\sin \frac{1}{2} a$, given in Art. 33, and dividing by the product of the analogous expressions for $\sin \frac{1}{2} b$ and $\sin \frac{1}{2} c$, since $\cos(S-A)$, $\cos(S-B)$, $\cos(S-C)$ become *in plano*, respectively, $\sin A$, $\sin B$, $\sin C$, we get,

$$\frac{a^2}{c} = \frac{\sin^2 A}{\sin B \sin C}.$$

Hence
$$\frac{a^2 \sin B \sin C}{\sin(B+C)} = bc \sin A = 2\Delta.$$

5. See Ex. 1.

6. See Ex. 3.

7. Draw the required figure, and join AA'. Denote the angles BAA' and $BA'A$, respectively, by x and x', Then we have—

(1) $\cos B = -\cos x \cos x' + \sin x \sin x' \cos AA'$;

(2) $\cos B' = -\cos(A-x)\cos(A'-x') + \sin(A-x)\sin(A'-x')\cos AA'$;

(3) $\cos C = \cos x \cos(A'-x') + \sin x \sin(A'-x')\cos AA'$;

(4) $\cos C' = \cos x' \cos(A-x) + \sin x' \sin(A-x)\cos AA'$.

Multiply equations (1) and (2), and subtract from the result the product of equations (3) and (4); therefore, &c.

8. (α) follows at once from Ex. 7. (β) We have

$\Sigma(\cos B \cos B' - \cos C \cos C') = \Sigma \sin A \sin A'$; therefore, &c.

9. $A = 94° 56' 36{\cdot}6''$, $B = 63° 0' 24{\cdot}6''$, $C = 74° 6' 20{\cdot}6''$.

10. $A = 54° 47' 53{\cdot}6''$, $b = 34° 36' 11''$, $c = 50° 7' 13{\cdot}2''$.

IX.

1. By aid of Paper VIII. Ex. 7, we obtain

$\Sigma(\sin A \sin A' \cos \delta_1) \equiv \Sigma(\cos B \cos B' - \cos C \cos C') \equiv 0$.

2. It is evident, in the preceding Example, that if δ_1 and δ_2 are quadrants, each term in that equation vanishes. Hence $\cos \delta_3 = 0$; therefore, &c.

3. The great circle on which they lie is the polar of the intersection of the perpendiculars of the triangle. (See Note, p. 23.)

4. This follows from Paper V. Ex. 1. See also the preceding solution, Ex. 3.

5. An obvious particular case of the preceding Example, and which may be verified directly as follows:—Draw perpendiculars from A and P to the base BC; the distance between their feet is $AP\cos X$; then

$$\Sigma\, aAP\cos X = 2\Sigma(BP^2 - CP^2) - 2\Sigma(b^2 - c^2);$$

i. e. $$aAP\cos X + bBP\cos Y + cCP\cos Z = 0.$$

6. In Ex. 4, let $a = b = c$; therefore, &c.

7. The left-hand side of the equation reduces to

$\cos a - \cos\beta\cos\gamma\cos^2 p - \sin b \sin c \cos A$; since $\beta + \gamma = a$,

and $\sin b \sin c \sin A = \sin a \sin p$.

Again, $\cos a - \sin b \sin c \cos A = \cos b \cos c$.

Hence the given expression becomes

$$\cos b \cos c - \cos\beta \cos\gamma \cos^2 p, \text{ or } \cos\beta \cos\gamma \cos^2 p - \cos\beta \cos\gamma \cos^2 p.$$

[Art. 44 (1).]

Again, $$\cot A = \frac{\cos a - \cos\beta \cos\gamma \cos^2 p}{\sin a \sin p}.$$

Now A is a maximum when $\cot A$ is a minimum; *i. e.* when $\cos\beta \cos\gamma$ is a maximum; but

$$2\cos\beta \cos\gamma = \cos(\beta+\gamma) + \cos(\beta-\gamma) = \cos a + \cos(\beta-\gamma).$$

From this result it follows that the vertical angle is a maximum when the perpendicular from it on the base bisects the base; therefore, &c.

[Cf. TOWNSEND's *Mod. Geom.*, vol. i., Ex. 7, p. 47.]

8. For $\sin\frac{1}{2}a$, $\sin\frac{1}{2}b$, &c., substitute the chords of the arcs a, b, &c., to which they are proportional, and the expression reduces to that which gives a diagonal of a plane cyclic quadrilateral in terms of its sides.

[See TODHUNTER's *Plane Trig.*, Art. 254.]

9. $a = 66°\ 19'\ 44{\cdot}17''$, $b = 45°\ 6'\ 32{\cdot}75''$, $C = 94°\ 21'\ 40{\cdot}3''$.

10. $A = 51°\ 3'\ 4{\cdot}2''$, $b = 55°\ 20'\ 23{\cdot}35''$, $c = 66°\ 30'\ 30{\cdot}6''$.

X.

1.* $A+a=\alpha$, $B+b=\beta$, $C+c=\gamma$, $A-a=\theta$, $B-b=\phi$, $C-c=\psi$;

$$\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c}; \quad \therefore \frac{\tan\frac{1}{2}\theta}{\tan\frac{1}{2}\alpha} = \frac{\tan\frac{1}{2}\phi}{\tan\frac{1}{2}\beta} = \frac{\tan\frac{1}{2}\psi}{\tan\frac{1}{2}\gamma}; \qquad (1)$$

but $$\frac{\cos\frac{1}{2}(A+B)}{\cos\frac{1}{2}(a+b)} = \frac{\sin\frac{1}{2}C}{\cos\frac{1}{2}c} = \frac{\cos\frac{1}{2}(\pi-C)}{\cos\frac{1}{2}c}; \qquad \text{(GAUSS.)}$$

therefore

$$\tan\tfrac{1}{4}(\theta+\phi)\tan\tfrac{1}{4}(\beta+\alpha) = \tan\tfrac{1}{4}(\pi-\gamma)\tan\tfrac{1}{4}(\pi-\psi). \qquad (2)$$

Again,

$$\frac{\sin\frac{1}{2}(A-B)}{\sin\frac{1}{2}(a-b)} = \frac{\cos\frac{1}{2}C}{\sin\frac{1}{2}c} = \frac{\sin\frac{1}{2}(\pi-C)}{\sin\frac{1}{2}c}; \quad \therefore \frac{\tan\frac{1}{4}(\theta-\phi)}{\tan\frac{1}{4}(\alpha-\beta)} = \frac{\tan\frac{1}{4}(\pi-\gamma)}{\tan\frac{1}{4}(\pi-\psi)}. \qquad (3)$$

* We are indebted to Mr. T. A. Finch for the first solution we give of this question; and to Mr. Robert Russell for an independent method. Mr. Russell remarks that the equations used by him to solve the triangle are likewise applicable to the more general problem—"Given *any three* of the six quantities $A \pm a$, $B \pm b$, $C \pm c$; solve the triangle."

(2) and (3) multiplied together give the ratio $\cos\frac{1}{2}\theta : \cos\frac{1}{2}\phi$, and from (1) we have the ratio $\tan\frac{1}{2}\theta : \tan\frac{1}{2}\phi$; and hence θ, ϕ, ψ are known; therefore, &c.

Cor.—Given $A-a$, $B-b$, and $C-c$; find the remaining parts.

Otherwise thus:—

$$\frac{\sin A}{\sin a} = \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c}; \text{ therefore } \frac{\sin A + \sin a}{\sin A - \sin a} = \ldots = \ldots$$

Hence
$$\frac{\tan\frac{1}{2}(A+a)}{\tan\frac{1}{2}(A-a)} = \frac{\tan\frac{1}{2}(B+b)}{\tan\frac{1}{2}(B-b)} = \ldots$$

Let
$$\cot\tfrac{1}{2}(A+a) = \iota \sin X,$$

and
$$\cot\tfrac{1}{2}(A-a) = \iota \sin x, \quad \text{where } \iota = \sqrt{-1}.$$

Hence
$$\frac{\sin X}{\sin x} = \frac{\sin Y}{\sin y} = \frac{\sin Z}{\sin z};$$

from which the question reduces to—Having given the three angles X, Y, Z of a spherical triangle; find the sides.*

We have thus—
$$\cos x = \frac{\cos X + \cos Y \cos Z}{\sin Y \sin Z};$$

or
$$\operatorname{cosec}\tfrac{1}{2}(A-a) = \frac{\operatorname{cosec}\frac{1}{2}(A+a) + \operatorname{cosec}\frac{1}{2}(B+b)\operatorname{cosec}\frac{1}{2}(C+c}{-\cot\frac{1}{2}(B+b)\cot\frac{1}{2}(C+c)};$$

or
$$\sin\tfrac{1}{2}(A-a) = -\frac{\sin\frac{1}{2}(A+a)\cos\frac{1}{2}(B+b)\cos\frac{1}{2}(C+c)}{\sin\frac{1}{2}(A+a) + \sin\frac{1}{2}(B+b)\sin\frac{1}{2}(C+c)}.$$

2. Let $AO = BO = CO = R$, and let $PO = \delta$, and the angle $POC = \theta$. Then we have—

$$\cos AP = \cos R \cos\delta + \sin R \sin\delta \cos(120-\theta). \quad (1)$$

$$\cos BP = \cos R \cos\delta + \sin R \sin\delta \cos(120+\theta). \quad (2)$$

$$\cos CP = \cos R \cos\delta + \sin R \sin\delta \cos\theta. \quad (3)$$

Hence, by addition,

$$\cos AP + \cos BP + \cos CP = 3\cos R \cos\delta. \quad (4)$$

* Mr. Robert Russell, *Quarterly Journal of Mathematics*, vol. xx.

Multiply (2) by ω and (3) by ω^2, where $\omega^3 = 1$, and add; then

$$\cos AP + \omega \cos BP + \omega^2 \cos CP$$
$$= \sin R \sin \delta \{\cos (120 - \theta) + \omega \cos (120 + \theta) + \omega^2 \cos \theta\}. \quad (5)$$

Similarly,

$$\cos AP + \omega^2 \cos BP + \omega \cos CP$$
$$= \sin R \sin \delta \{\cos (120 - \theta) + \omega^2 \cos (120 + \theta) + \omega \cos \theta\}. \quad (6)$$

Multiplying equations (5) and (6), we have, on reduction,

$$\cos^2 AP + \cos^2 BP + \cos^2 CP - \cos BP \cos CP - \ldots = \tfrac{3}{4} \sin^2 R \sin^2 \delta;$$

or by the aid of (4) $= \frac{1}{4} (\cos AP + \cos BP + \cos CP)^2 \tan^2 R \tan^2 \delta$.

NOTE.—To simplify the right-hand side of the product of (5) and (6), throw the part within the brackets into the determinant form—

$$\begin{vmatrix} 1 & \cos \theta & \cos (120 + \theta) \\ 1 & \cos (120 - \theta) & \cos \theta \\ 1 & \cos (120 + \theta) & \cos (120 - \theta) \end{vmatrix};$$

and remembering that

$$\cos \theta + \cos (120 + \theta) + \cos (120 - \theta) = 0,$$

the reduction presents no difficulty.

(*Educational Times.*)

3. The right angles on the primitive figure are transformed into the diagonals of the supplemental figure; therefore, &c.

4. It has been seen in Paper X., Ex. 2, that

$$\cos AP + \cos BP + \cos CP = 3 \cos AO \cos OP.$$

In this instance, $\cos AO = \dfrac{1}{\sqrt{3}}$;

therefore, &c.

5. $\cos A = \dfrac{\tan b}{\tan c}$; therefore $\dfrac{\tan c + \tan b}{\tan c - \tan b} = \dfrac{\sin (c + b)}{\sin (c - b)} = \text{given}.$

Hence if $c + b$ be known, $c - b$ is determined, and therefore c and b.

6. See fig. 10, p. 21. The arc $LM = 2\,XY$. Hence the angle AOB is known, and consequently the triangle AOB is completely determined; and since $OA + OC = \pi$, the locus of the vertex is a circle, of which O is the pole.

7. This follows at once from Ex. 5, where it is shown that sin $(c + b)$ varies as sin $(c - b)$ when A is given.

8. From A, one of the extremities of the given base, let fall AP perpendicular to the given great circle, and produce AP to Q, so that $AP = PQ$. Join Q to B, the other extremity of the given base. Let BQ cut the given circle in C. Then ABC and its colunar ABC' are the required triangles. C' being diametrically opposite to C, is on the given circle.

9. $A = 70° \ 25' \ 33\frac{1}{2}''$, $B = 51° \ 5' \ 47''$, $c = 73° \ 19' \ 28\frac{1}{2}''$.

10. $A = 70° \ 22' \ 16{\cdot}1''$, $B = 49° \ 25' \ 12{\cdot}3''$ $c = 61° \ 54' \ 6''$.

XI.

1. Form a spherical triangle by joining the poles of the given great circles to the point P, and it is obvious that of this triangle we are given the base and the sum of the cosines of the sides; therefore, &c. (See p. 60.)

2. By the method of the previous exercise. See also Ex. 1, p. 60.

3. See Paper VII. Ex. 3.

4. By the ordinary formulæ for right-angled triangles.

5. From the extremities A and B of the base let fall secondaries AX and BY on the given locus. Then

$\cos c = \cos a \cos b = \cos AX \cos CX \cos BY \cos CY$; but AX, BY, and XY, are given arcs. Hence it is required to divide a given arc XY at a point C, such that cos CX cos CY may be given.

6. The product is evidently

$$\sin a \sin b \cos^2 \tfrac{1}{2} C = \frac{\sin a \sin b \sin s \sin (s-c)}{\sin a \sin b} = \sin s \sin (s - c),$$

which is given.

7. The product is

$$\sin a \sin b \sin^2 \tfrac{1}{2} C = \sin (s - a) \sin (s - b);$$

therefore, &c.

8. $\sin p_1 \sin p_2 = \sin a \sin b \cos^2 \frac{1}{2} C$; $\sin p_3 \sin p_4 = \sin a \sin b \sin^2 \frac{1}{2} C$; therefore, &c. (See Exs. 6 and 7.)

9. $a = 29° \ 26' \ 30''$, $b = 84° \ 37' \ 15''$, $c = 85° \ 19' \ 2''$.

10. $A = 82° \ 24' \ 33{\cdot}4''$ $B = 45° \ 2' \ 9{\cdot}88''$, $c = 60° \ 20' \ 10{\cdot}5''$.

XII.

1. For by Art. 42, p. 60, we have

$$\cos a + \cos b = 2 \cos \tfrac{1}{2} c \cos \beta,$$

$$\cos a - \cos b = 2 \sin \tfrac{1}{2} c \cos \beta':$$

squaring each of these equations, and subtracting the results—(α) follows at once; when the results are added, we obtain (β).

2. See p. 49 (12).

3. This equation is an expression for the base in terms of two sides and the angle included between them, and must, therefore, be identical with the form

$$\cos c = \cos a \cos b + \sin a \sin b \cos C.$$

The identity of these equations will be at once made evident by the substitution of

$$\frac{1 - \tan^2 \tfrac{1}{2} a}{1 + \tan^2 \tfrac{1}{2} a} \text{ for } \cos a, \text{ etc.,}$$

and

$$\frac{2 \tan \tfrac{1}{2} a}{1 + \tan^2 \tfrac{1}{2} a} \text{ for } \sin a, \text{ etc.}$$

4. Substitute the value of $\tan \frac{1}{2} a$ in terms of the angles (p. 49).

5. Apply Art. 40, p. 58.

6. *Firstly*—By the rule of sines

$$\frac{\sin a}{\sin b} = \frac{\sin (c - a)}{\sin a} = \sin c \cot a - \cos c ;$$

hence

$$\cot a = \frac{\sin b \cos c + \sin a}{\sin b \sin c} ;$$

i. e.

$$\sin^2 a = \frac{\sin^2 b \sin^2 c}{\sin^2 a + \sin^2 b + 2 \sin a \sin b \cos c} .$$

Similarly,

$$\sin^2 \beta = \frac{\sin^2 a \sin^2 c}{\sin^2 a + \sin^2 b + 2 \sin a \sin b \cos c} ;$$

therefore, &c.

Secondly—

$$\sin^2 a \sin^2 \theta = \sin^2 b \sin^2 \tfrac{1}{2} C.$$

Substitute the value already found for $\sin^2 a$; therefore, &c.

Thirdly—Multiply together the results in Exs. 9 and 10, pp. 47 and 48.

7. When $c - b$ is a maximum (by Paper x. Ex. 7), $c = 90 - b$; hence

$$\cos c = \sin b = \cos a \cos b, \quad i.e. \quad \cos a = \tan b = \cot c.$$

Again, $\cos A = \tan b \cot c$, and is therefore equal to $\cos^2 a$.

8. $a = 53° 24' 11{\cdot}78''$, $b = 37° 27' 55{\cdot}34''$, $C = 76° 47' 34{\cdot}84''$.

9. $a = 36° 9' 12{\cdot}78''$, $b = 44° 23' 39{\cdot}54''$, $c = 40° 0' 3{\cdot}8''$.

10. $A = 94° 16' 17\frac{1}{4}''$, $B = 56° 55' 38\frac{3}{4}''$, $c = 73° 30' 28''$.

11. $b = 42° 37' 43\frac{1}{4}''$, $c = 56° 7' 29\frac{1}{2}''$, $B = 54° 39' 26{\cdot}4''$;

or, $b = 137° 22' 16\frac{3}{4}''$, $c = 123° 52' 30\frac{1}{2}''$, $B = 125° 20' 33{\cdot}6'$.

12. $b = 41° 12' 54\frac{1}{2}''$, $c = 61° 57' 58\frac{1}{2}''$, $B = 48° 17' 8\frac{1}{4}''$,

or, $b = 138° 47' 5\frac{1}{2}''$, $c = 118° 2' 1\frac{1}{2}''$, $B = 131° 42' 51\frac{3}{4}''$;

13. $a = 82° 38' 40{\cdot}8''$, $b = 67° 6' 44{\cdot}48''$, $C = 88° 58' 12{\cdot}38''$.

14. $a = 71° 35' 16''$, $b = 40° 33' 12''$, $C = 76° 2' 39{\cdot}72''$.

15. $a = 57° 7' 7{\cdot}42''$, $b = 69° 22' 20{\cdot}4''$, $c = 54° 3' 18{\cdot}2''$.

END OF PART I.



